


DRAFT - Hillsburgh Dam

Town of Erin
Environmental Assessment - Natural Environment Report



Prepared for:
Town of Erin

Project Number:
AA12-137A

Date:
March 15, 2016

ABOUD & ASSOCIATES INC.
Consulting Arborists • Ecologists • Landscape Designers

591 Woolwich Street . Guelph . Ontario . N1H 3Y5 . T:519.822.6839 . F:519.822.4052 . info@aboudtng.com . www.aboudtng.com



EXECUTIVE SUMMARY

ENVIRONMENTAL ASSESSMENT - NATURAL ENVIRONMENT REPORT

About & Associates Incorporated (AA) was retained by Triton Engineering Services Limited (Triton) on behalf of the Town of Erin to complete the natural heritage component of a Schedule B Municipal Class Environmental Assessment (EA). The EA is being completed in order to determine the best option to ensure the long term safety of the Hillsburgh Dam, Bridge and Pond.

The Hillsburgh Dam is an earthen berm located within the community of Hillsburgh, part of the Town of Erin, Wellington County. The water held back by the Dam creates the Hillsburgh Pond, an approximately 9.0 ha, open body of water, with associated wetland areas. The river system of the Hillsburgh pond is the West Credit River, a cold water river system.

The study area for the Natural Environment report is 77.05 ha, centered on the Hillsburgh Dam and includes lands upstream and downstream of the dam. Natural features within the study area include Provincially Significant Wetlands, meadows, open water communities and forests.

The natural heritage studies of the EA characterized and mapped the significant natural features within the study area, identified potential constraints and analyzed proposed alternatives. Species at Risk (SAR), Fish Habitat, Significant Wildlife Habitat, Rare Species, Landscape-level Features, and a Provincially Significant Wetland were identified in the study area. These findings were considered as part of the Analysis of Alternatives.

The Four Alternatives assessed to determine the preferred Alternative are:

- Alternative A – Do Nothing
- Alternative B – Rehabilitate Hillsburgh Dam and Reconstruct Station Street Bridge
- Alternative C (Option 1): Rehabilitate Station Street Bridge and Decommission Dam
- Alternative C (Option 2): Rehabilitate Station Street Bridge, Decommission Dam and Construct Offline Pond
- Alternative D (Option 1): Reconstruct Station Street Bridge and Decommission Dam
- Alternative D (Option 2): Reconstruct Station Street Bridge, Decommission Dam and Construct Offline Pond

The Analysis of Alternatives identified the potential and actual impacts of each proposed EA Alternative with respect to the existing natural heritage features in the study area and surrounding landscape. The analysis concluded that there are two preferred alternatives: Alternative C - Option 2 and Alternative D - Option 2. These alternatives have the least negative impacts to the existing natural heritage features. Both of these also provide benefits to cold water Fish and Fish Habitat.

The next preferred alternative is Alternative B. If Alternative B is selected, mitigation measures should be considered to minimize impacts to aquatic habitat through the creation of a fish-bypass to allow fish passage and bottom draw dam design to minimize thermal impacts to the downstream watercourse. The least preferred alternatives are Alternative C -Option 1 and Alternative D - Option 1.

Under Alternative A (Do nothing), no new impacts are anticipated under the current conditions. However if failure of the dam and/or bridge occurs, significant negative impacts are anticipated to all natural environment criteria, the extent of which is unknown.

For the selected Alternative, measures should be implemented to protect the natural environment during construction and mitigate short and long-term impacts of the overall ecological integrity of the area.

Opportunities to enhance the natural environment as part of protection and mitigation measures should be considered following selection of the overall preferred Alternative.

DRAFT

Table of Contents

1.0	INTRODUCTION.....	1
1.1	STUDY AREA.....	1
1.2	EXISTING LAND USE.....	1
1.3	EXISTING REGULATIONS.....	2
1.3.1	<i>Provincial Policy Statement.....</i>	2
1.3.2	<i>Endangered Species Act, 2007.....</i>	3
1.3.3	<i>Fisheries Act, 1985.....</i>	3
1.3.4	<i>Credit Valley Conservation.....</i>	4
1.3.5	<i>Wellington County Official Plan.....</i>	4
1.3.6	<i>Town of Erin Official Plan.....</i>	4
1.4	CREDIT RIVER FISHERIES MANAGEMENT PLAN.....	5
1.5	CONSULTATION AND COMMENTS.....	6
1.6	TERMS OF REFERENCE.....	7
2.0	METHODS.....	8
2.1	BACKGROUND REVIEW.....	8
2.2	VEGETATION.....	8
2.2.1	<i>Ecological Land Classification.....</i>	8
2.2.2	<i>Botanical Inventory.....</i>	9
2.3	PROVINCIALY SIGNIFICANT WETLANDS.....	9
2.4	WILDLIFE.....	10
2.4.1	<i>Amphibians (Anurans).....</i>	10
2.4.2	<i>Breeding Birds.....</i>	10
2.4.3	<i>Snakes.....</i>	11
2.4.4	<i>Salamanders.....</i>	11
2.4.5	<i>Turtles.....</i>	11
2.4.6	<i>Winter Wildlife.....</i>	11
2.4.7	<i>Migratory Birds.....</i>	12
2.4.8	<i>Incidental Wildlife Observations.....</i>	12
2.5	SIGNIFICANT WILDLIFE HABITAT.....	12
2.6	SAR HABITAT ASSESSMENT.....	12
2.7	AQUATIC HABITAT ASSESSMENT.....	12
3.0	EXISTING CONDITIONS.....	15
3.1	BACKGROUND REVIEW.....	15
3.1.1	<i>Natural Heritage Information Centre - Species at Risk.....</i>	15
3.1.2	<i>Ontario Breeding Bird Atlas.....</i>	15
3.1.3	<i>Ontario Reptile and Amphibian Atlas.....</i>	16
3.1.4	<i>Atlas of the Mammals of Ontario.....</i>	16
3.1.6	<i>Credit Valley Conservation.....</i>	16
3.1.7	<i>Ministry of Natural Resources and Forestry.....</i>	17
3.2	VEGETATION.....	18
3.2.2	<i>Botanical Inventory.....</i>	19
3.3	PROVINCIALY SIGNIFICANT WETLANDS.....	20
3.3.1	<i>Boundary Review.....</i>	20
3.3.2	<i>Wetland Characteristics.....</i>	20
3.4	WILDLIFE.....	20
3.4.1	<i>Amphibians (Anurans).....</i>	20
3.4.2	<i>Breeding Birds.....</i>	22

3.4.3	<i>Snakes</i>	24
3.4.4	<i>Salamanders</i>	25
3.4.5	<i>Turtles</i>	25
3.4.6	<i>Winter Wildlife</i>	27
3.4.7	<i>Migratory Birds</i>	27
3.4.8	<i>Incidental Wildlife Observations</i>	27
3.4.9	<i>Species Listed under the Endangered Species Act</i>	28
3.5	SIGNIFICANT WILDLIFE HABITAT.....	29
3.6	SAR HABITAT ASSESSMENT.....	30
3.6.1	<i>Vegetation</i>	30
3.6.2	<i>Wildlife</i>	31
3.6.3	<i>Fish</i>	32
3.7	AQUATIC HABITAT ASSESSMENT.....	32
3.8	LANDSCAPE EVALUATION.....	35
3.8.1	<i>Ecoregion</i>	35
3.8.2	<i>Surficial Geology and Groundwater</i>	35
3.8.3	<i>Connectivity and Existing Natural Features</i>	35
3.8.4	<i>Aquatic Habitat</i>	35
3.8.5	<i>Rare Features</i>	36
4.0	SUMMARY OF NATURAL HERITAGE CONSTRAINTS	37
4.1	SUMMARY OF EXISTING CONDITIONS.....	37
4.1.1	<i>Vegetation</i>	37
4.1.2	<i>Wetlands</i>	37
4.1.3	<i>Wildlife</i>	38
4.1.4	<i>Significant Wildlife Habitat (SWH)</i>	39
4.1.5	<i>Species with Conservation Designation Habitat Assessment</i>	39
4.1.6	<i>Aquatic Habitat Assessment</i>	39
4.2	SUMMARY OF SIGNIFICANT FEATURES.....	41
5.0	ASSESSMENT OF ALTERNATIVES, AND IMPACTS TO THE NATURAL ENVIRONMENT	44
5.1	ALTERNATIVES CONSIDERED FOR THE HILLSBURGH DAM AND POND.....	44
5.2	EVALUATION OF IMPACTS.....	45
5.3	SUMMARY OF IMPACTS TO THE NATURAL ENVIRONMENT.....	47
5.3.1	<i>Alternative A - Do Nothing</i>	47
5.3.2	<i>Alternative B - Rehabilitate Hillsburgh Dam and Reconstruct Station Street Bridge</i>	47
5.3.3	<i>Alternative C - Rehabilitate Station Street Bridge and Decommission the Hillsburgh Dam</i>	48
5.3.4	<i>Alternative D - Reconstruct Station Street Bridge, Decommission Dam</i>	50
6.0	CONCLUSION	53
5.0	REFERENCES	55

List of Tables

Table 1. NHIC Species at Risk Records	15
Table 2. Ecological Land Classification	18
Table 3. Summary of Amphibian Observations (2015)	21
Table 4. Regionally or Locally Significant Breeding Bird Species	24
Table 5. Turtle Habitat Results.....	25
Table 6. Incidental Species with Conservation Designation Observations	28
Table 7. Confirmed Significant Wildlife Habitat.....	30
Table 8. Summary of Significant Features	41
Table 9: Comparison of Alternatives.....	46

List of Figures

Figure 1	Site Location
Figure 2	Property Access and CVC Botanical Study Area
Figure 3	Ecological Land Classifications
Figure 4	Wildlife Habitat Targets & Survey Locations
Figure 5	Aquatic Habitat Assessments
Figure 6	Landscape Evaluation
Figure 7	Significant Wildlife Observations and Significant Wildlife Habitat

List of Appendices

Appendix 1	Communications
Appendix 2	Terms of Reference and Approval
Appendix 3	Site Investigation Details
Appendix 4	Ecological Land Classification – Community Descriptions
Appendix 5	Ecological Land Classification Data Sheets
Appendix 6	Vascular Plant List
Appendix 7	Anuran Survey Results and Call Codes
Appendix 8	Breeding Bird Results and Evidence Codes
Appendix 9	Marsh Breeding Bird Results
Appendix 10	Snake Transect Survey Results
Appendix 11	Turtle Basking Survey Results
Appendix 12	Winter Wildlife Results
Appendix 13	Migratory Bird Survey
Appendix 14	Shorebird Habitat Assessment
Appendix 15	Incidental Wildlife Observations
Appendix 16	Significant Wildlife Habitat Assessment
Appendix 17	Species at Risk Habitat Assessment
Appendix 18	Aquatic Habitat Description
Appendix 19	Background Wildlife List
Appendix 20	Fish Records
Appendix 21	Curriculum Vitae

Glossary of Terms

BBS: Breeding Bird Survey

CC: coefficient of Conservatism

COSSARO: Committee on the status of Species at Risk Ontario

COSEWIC: Committee on the Status of Endangered Wildlife in Canada

CRFMP: The Credit River Fisheries Management Plan

CVC: Credit Valley Conservation

CVC Species of Conservation Concern: CVC ranking of Species based on Conservation Concern Status, from Tier 1 to Tier 5.

DFO: Department of Fisheries and Oceans Canada

EA: Environmental Assessment

ELC: Ecological Land classification

END: Endangered Species

ESA: Endangered Species Act

ESSMP: Erin Service and Settlement Master Plan

G-Rank: Conservation Status of Species at the global Level

LIO: Land Information Ontario

MMP: Marsh Monitoring Protocol

MNRF: Ministry of Natural Resources and Forestry

NHIC: Natural Heritage Information Center

NRVIS: Natural Resources and Values Information System

OBBA: Ontario Breeding Bird Atlas

OMA: Ontario Mammal Atlas

ORAA: Ontario Reptile and Amphibian Atlas

OP: Official Plan

OWES: Ontario Wetland Evaluation System

PPS: Provincial Policy Statement

PIF: Partners in Flight

PSW: Provincially Significant Wetland

SAR: Species at Risk

SARA: Species at Risk Act

SC: Special Concern Species

Species with conservation designation: All species listed under SARA, COSEWIC, ESA and/or an S1-S3 provincial designation.

S-Rank: Conservation Status of Species at the Provincial Level

SWH: Significant Wildlife Habitat

THR: Threatened Species

VASCAN: Database of Vascular Plants of Canada

1.0 Introduction

Aboud & Associates Incorporated (AA) was retained by Triton Engineering Services Limited (Triton) on behalf of the Town of Erin to complete the natural heritage component of a Schedule B Municipal Class Environmental Assessment (EA). The EA is being completed in order to determine the best option to ensure the long term safety of the Hillsburgh Dam and associated bridge with due consideration for the natural environment, transportation, socio-economic impacts, constructability and cost. The natural heritage existing condition component of the EA focuses on characterizing the existing natural features within the study area, mapping significant natural features and identifying potential constraints.

1.1 Study Area

The study area is 77.05 ha and located in the community of Hillsburgh - Town of Erin (*Figure 1*). It is centered on the Hillsburgh Dam and includes lands upstream and downstream of the dam, extending south downstream to Wellington Road 22 and upstream along the two upstream tributaries of the West Credit River. The study area is entirely within the jurisdiction of Credit Valley Conservation (CVC), and almost entirely within CVC's Regulation Limit. Natural features within the study area include wetlands, meadows, open water communities and forests.

The Hillsburgh Dam is an earthen berm located within the community of Hillsburgh, part of the Town of Erin, Wellington County. The water held back by the Dam creates the Hillsburgh Pond, an approximately 9.0 ha, open body of water. The Dam supports a section of Station Street Road, a two lane municipal road that crosses the West Credit River by way of the Dam and associated bridge (Structure #2064); the latter being the main outflow for the Hillsburgh pond.

1.2 Existing Land Use

The individual properties that comprise the study area are a combination of private property, and public property (Town of Erin, Credit Valley Conservation land and County of Wellington). The majority of the study area is natural or naturalized land, containing a diversity of ecosystems, including the Provincially Significant West Credit River Wetland Complex, the West Credit River, Significant Woodlands, and open water communities. In addition to the Hillsburgh Pond, there are two other aquatic communities resulting from impoundment of water behind the dams on the West Credit River, and a number of smaller dug offline ponds throughout the study area (*Figure 2*).

The watercourse within the study area is a mixture of cold and warm water systems, containing both cold and warm water species of fish, although the entire watercourse is managed as a Coldwater Fishery within the limits of the study area (CRFMP 2001; ESSMP 2011; Pers. Comm. T. Slaght, 2014). The Elora-Cataract Trailway crosses through the study area and is the main access route into the different sections of the study area.

Within or adjacent the study area are a number of private residential properties comprised of dwellings, driveways and associated landscaping and yard maintenance. Residential areas were not included in the Ecological Land Classification (ELC) surveys and flora species within residential gardens and yards are not included in flora species lists.

Access to specific properties within the study area was requested by Triton through letter and door to door communication with landowners. Due to private property restrictions, large sections

of the study area were not accessible, and could only be assessed from the edge of property, aerial photo interpretation and through background resources. The extents of lands accessed as part of the current investigation are shown in *Figure 2*.

1.3 Existing Regulations

1.3.1 Provincial Policy Statement

The *Provincial Policy Statement* ([PPS] OMMHA 2014) provides policy direction on matters of provincial interest related to land use planning and development.

The PPS states that:

“Natural features and areas shall be protected for the long term.”

And that:

“The diversity and connectivity of natural features in an area, and the long-term ecological function and biodiversity of natural heritage systems, should be maintained, restored or, where possible, improved, recognizing linkages between and among natural heritage features and areas, surface water features and ground water features.”

Under the PPS, development and site alteration are not permitted in:

- a) significant wetlands;
- b) significant woodlands;
- c) significant valleylands;
- d) significant wildlife habitat;
- e) significant areas of natural and scientific interest; and
- f) coastal wetlands,

unless it has been demonstrated that there will be no negative impacts on the natural features or their ecological functions.

The PPS (2014) also states that:

1. Development and site alteration is not permitted in fish habitat, habitat of endangered species and threatened species except in accordance with provincial and federal requirements.
2. Development and site alteration is not permitted on adjacent lands to the natural heritage features and areas identified above, unless the ecological function of the adjacent lands has been evaluated and it has been demonstrated that there will be no negative impacts on the natural features or on their ecological functions.
3. Development and site alteration is restricted in or near sensitive surface water features and sensitive ground water features in order to protect the hydrologic functions of the feature. Mitigation and/or alternative development approaches may be required in order to protect, improve or restore sensitive surface water features, sensitive ground water features, and their hydrologic functions.

1.3.2 Endangered Species Act, 2007

The provincial Endangered Species Act, 2007 (ESA) provides protection to species designated as Threatened or Endangered on the Species at Risk in Ontario list (MNR 2015a). The habitat of species at risk is also generally protected under the ESA. Protected habitat is habitat identified as essential for life processes including: breeding, rearing, feeding, hibernation and migration.

The ESA (Subsection 9(1)) states that:

“No person shall,

- (a) kill, harm, harass, capture or take a living member of a species that is listed on the Species at Risk in Ontario List as an extirpated, endangered or threatened species;*
- (b) possess, transport, collect, buy, sell, lease, trade or offer to buy, sell, lease or trade,*
 - (i) a living or dead member of a species that is listed on the Species at Risk in Ontario List as an extirpated, endangered or threatened species,*
 - (ii) any part of a living or dead member of a species referred to in subclause (i),*
 - (iii) anything derived from a living or dead member of a species referred to in subclause (i); or*
- (c) sell, lease, trade or offer to sell, lease or trade anything that the person represents to be a thing described in subclause (b) (i), (ii) or (iii).*

Clause 10(1)(a) of the ESA also states that:

“No person shall damage or destroy the habitat of a species that is listed on the Species at Risk in Ontario list as an endangered or threatened species.”

An authorization or permit between the proponent and the Minister of Natural Resources and Forestry is required to authorize activities that would otherwise be prohibited by subsection 9(1) and 10(1) of the ESA.

1.3.3 Fisheries Act, 1985

The study area contains fish bearing waters in the form of open water, rivers and wetlands. These areas and the fish within are protected under the Federal Fisheries Act, 1985. The Fisheries Act provides protection for the sustainability and ongoing productivity of Canada's recreational, commercial and Aboriginal fisheries.

Section 35 (1) of the Fisheries Act States that:

“No person shall carry on any work, undertake activity that results in serious harm to fish that are part of a commercial, recreational or Aboriginal fishery, or fish that support such a fishery”

The Fisheries Act requires that projects and activities avoid causing serious harm to fish and fish habitat unless authorized to do so by the Department of Fisheries and Oceans Canada (DFO). This applies to work conducted in or near waterbodies that support recreational, commercial and Aboriginal fisheries. Within the context of the Hillsburgh EA, any proposed actions that could impact fish or fish habitat would need to be assessed for compliance with the Fisheries Act. If it is determined that proposed actions will cause serious harm to fish that cannot be mitigated for then a Fisheries Act Authorization would be required.

1.3.4 Credit Valley Conservation

The Study Area is located within the jurisdiction of CVC and contained within the CVC Regulation Limit from two regulated features: the West Credit River Wetland Complex (PSW), and the West Credit River.

CVC's *Policies are regulated under the Administration of the Development, Interference with Wetlands and Alterations to Shorelines and Watercourses Regulation* (Ontario Regulation 160/06).

Interference with a wetland or watercourse; or development within a regulated area is generally not permitted. Interference with a wetland or watercourse, or development may be permitted within a regulated area if, the control of flooding, erosion, dynamic beaches, pollution or the conservation of land will not be affected.

CVC may permit development or site alteration where impacts have been addressed through an environmental assessment, comprehensive environmental study or technical report (CVC 2010a).

1.3.5 Wellington County Official Plan

The *Wellington County Official Plan* (County of Wellington 2013, Section 5.5.4) states that, "woodlands over 4 ha and plantations over 10ha are considered to be significant by the County, and are included in the Greenlands System". Section 5.4 of the *Official Plan* (Section 5 – The Greenlands System) specifies that within the Greenlands System, areas with greater sensitivity or significance are identified and protected as 'Core Greenlands', and include Provincially Significant Wetlands. The *Wellington County Official Plan* (2013) shows that the study area contains 'Greenlands' and 'Core Greenlands'.

Section 5.5 of the *Official Plan* states that *Significant Woodlands* "will be protected from development or site alterations which would negatively impact the woodlands or their ecological function".

Section 5.4 of the OP states that *Core Greenlands* include: provincially significant wetlands, all other wetlands; habitat of endangered or threatened species and fish habitat; and hazardous lands.

Development and site alteration is not permitted in Provincially Significant Wetlands, in the habitat of threatened or endangered species, or fish habitat except in accordance with provincial and federal requirements.

1.3.6 Town of Erin Official Plan

The Town of Erin Official Plan (2012) encourages the protection and enhancement of natural heritage features, including the protection, preservation and enhancement of significant natural features such as rivers, streams, valley lands, wetlands, flood plains, headwaters, environmentally significant features, wildlife and fish habitats and lands with ecological functions. Ponds, lakes and reservoirs and natural links are also afforded protection from development or site alteration which would have negative impacts.

Lands designated under the OP as *Core Greenlands* are protected from any development or site alteration which would have a significant negative impact on the *Core Greenland* or their ecological function.

Fish Habitat is recognized as important under the OP fisheries policies. Fisheries are afforded protection through the maintenance of groundwater and surface water inflows, maintaining or establishing tree cover over rivers and streams, providing public access to fishery resources, and minimizing or eliminating negative thermal impacts to the fishery. The naturalization of watercourse corridors is also encouraged under the OP.

1.4 Credit River Fisheries Management Plan

The Credit River Fisheries Management Plan (CRFMP) was a joint project between the MNRF and the CVC, along with other government and non-government partners. The goal of the CRFMP is “to have a healthy aquatic ecosystem that provides long-term benefits to help satisfy society’s need for a high-quality environment, wholesome food, employment and income, recreational activities, and cultural heritage.” The CRFMP guides the protection and enhancement of the Credit River and provides fisheries management objectives for specific species of fish, as well as management objectives for specific management zones within the Credit River.

The Credit River watershed covers 871 km², eventually flowing into Lake Ontario. The watershed encompasses portions of nine municipalities, supports a human population of over 500 000 and has a diverse land use of urban, rural, agricultural and natural areas. Land use in the Credit River watershed has intensified with increasing urban growth. Limiting impacts to the watercourse and fish populations from growth and development is a key objective of the CRFMP.

The Credit River system provides high quality fishing opportunities for anglers, with the cold waters of the upper watershed offering fly fishing opportunities for Brook Trout (*Salvelinus fontinalis*) and Brown Trout (*Salmo trutta*). The systems contain at least 57 fish species, with new legal and illegal introduction continuing to increase the number of species. Biodiversity of the systems has increased with the introduction of sports fishing species such as, Brown Trout, Chinook (*Oncorhynchus tshawytscha*) and Coho Salmon (*Oncorhynchus kisutch*). Native sports fish such as Largemouth Bass (*Micropterus salmoides*) and Northern Pike (*Esox Lucius*) have experienced range expansion within the watershed through introductions and through the alteration and creation of habitats such as ponds and lakes. Undesirable invasive species, such as the Common Carp (*Cyprinus carpio*) and Round Goby (*Neogobius melanostomus*) have been introduced into the watershed (CRFMP 2002).

The CRFMP divides the watershed into three sub-watershed zones, consisting of the upper watershed, middle watershed and lower watershed. The upper watershed is on or above the Niagara Escarpment, and is primarily in rural landscapes. The watercourses of the Upper Watershed are of higher quality than the lower watersheds and have been retained in relatively natural conditions, with large riparian buffers and limited alterations to stream morphology. Base flow in these areas is provided through springs and groundwater discharge. Water is generally cold and of high quality.

Fisheries Management zones within the Credit River have been developed based on habitat, thermal conditions and fish community composition for specific stream sections within the watershed. The current EA study area is entirely within the Coldwater Fish Habitat Management Zone. The management zone extends upstream to the headwaters of the tributaries and extends downstream, approximately 20 km beyond the study area to the community of Inglewood. The fish communities in the management zone are comprised primarily of fish

species intolerant of water temperatures that exceed 22°C and commonly found only in groundwater-rich areas. Sport fish species common to the cold water communities include Brook Trout and Brown Trout.

Brook Trout and resident Brown Trout are the primary target management species within the Upper Watershed Coldwater Management Zone. Within the management zone, Coldwater construction timing windows must be adhered to, even if specific water bodies, such as the Hillsburgh Pond contain warm water fish species.

The CRFMP identifies risks and impacts to fisheries and recognizes online ponds as an ongoing concern within the watershed. Impacts associated with dams and ponds include; thermal warming, siltation, flooding, erosion, nutrient enrichment and pollution, and fish passage barriers. The Hillsburgh Dam and Pond are specifically cited as having known negative impacts to the management of the Coldwater fishery of the upper Credit River. The Management Plan recommends dam mitigation or removal in order to alleviate the impacts to fish communities. Removal of the dam would allow the watercourse to re-naturalize over time, permit fish passage, and create additional cold water habitat. In the absence of dam removal, potential dam mitigation measures addressed in the Management Plan include fishways (fish ladders), and bottom draw outlets. Fishways would allow fish passage and mitigate the impact of fish barriers. Bottom draw systems are designed to release water from near the bottom of a pond or body of water behind a dam where the open water community is colder than water drawn from the top, and therefore reducing thermal heating of the downstream watercourses.

The CRFMP guides decision-making and management of the Credit River and associated fisheries. Any actions regarding the Hillsburgh Dam and Pond should be examined with respect to the management objectives and target species of the Management Plan.

1.5 Consultation and Comments

A pre-consultation meeting was held on September 24, 2014 that included representatives from the CVC (T. Slaughter, J. Wong, and J. Clayton), MNRF (R. Whalen, D. Ryan), Town of Erin (L. Van Wyck), Triton Engineering Services Limited (C. Clark) and Aboud & Associates Inc. (S. Aboud, R. Hamelin). A summary of the meeting and follow-up minutes relevant to the natural heritage investigation are provided below; detailed Meeting Minutes are provided in *Appendix 1*.

1. Triton reviewed the project history, from temporary work to the present need for a permanent solution involving the completion of a Municipal Class Environmental Assessment (Class EA).
2. Aboud & Associates presented the proposed study area with respect to the Natural Heritage investigation. Modifications to the study area were recommended by MNRF and CVC.
3. MNRF and CVC reviewed available data for the project including:
 - a. Fish collection records;
 - b. Presence of invasive Round Goby;
 - c. Water temperature records;
 - d. Known Brook Trout spawning upstream and downstream;

- e. Known ground water seeps throughout the system, but no specific location;
 - f. CVC considers the Banded Killifish (*Fundulus diaphanous*) and Slimy Sculpin (*Cottus cognatus*), as important species due to the rarity in the watershed.
4. MNRF indicated that there are no known Species at Risk (SAR) in the study area.
 5. CVC stated that the main criteria when evaluating EA options will be; 1) Flood hazard reduction 2) Sediment/Erosion impact reduction.
 6. Triton reminded the group of potential property access limitations due to private property.
 7. General discussion on how the potential option could affect the existing PSW wetland complex.

On December 19, 2014 the CVC provided a letter, Re: Class Environmental Assessment Study to Triton Engineering Services (*Appendix 2*). The letter offered preliminary comments and recommendations for the Class EA study.

1.6 Terms of Reference

Based on the above regulations and policies (Section 1.3), the Credit River Fisheries Management Plan (Section 1.4), and communication with regulatory authorities, a proposed Terms of Reference (ToR) for the EA was developed and submitted to the CVC and MNRF on December 8th 2014. Comments regarding the proposed ToR were received from CVC on December 17, 2014 and from the MNRF on January 26, 2015. Follow up comments and request for clarifications were sent to CVC on January 6th, 2015. Response from CVC was received on January 23, 2015.

Based on comments received from the CVC, additional wildlife surveys (e.g. Snakes, Turtles, Salamander, Bat Maternity Roost and West Virginia White Surveys) were added to the EA study requirements. Correspondence with the MNRF was conducted to identify potential SAR within the study area and to determine the appropriate survey protocols. A letter was sent to CVC on April 10, 2015 to outline the SAR surveys and methods that would be completed as part of the EA. The appropriateness of the additional studies was confirmed by CVC on April 13, 2015 (Pers. comm., T. Slaught, 2015).

The Terms of Reference, CVC and MNRF comments, and final changes in study requirements, including the SAR surveys and methods are provided in *Appendix 2*.

2.0 Methods

2.1 Background Review

A background information review was conducted of both biological and physical features within the vicinity of the study area. The following resources were consulted as part of this review:

1. Fisheries and Oceans Canada (DFO), Online mapping (accessed: 2015)
2. Ministry of Natural Resources and Forestry (MNRF), Guelph District
3. Ministry of Natural Resources and Forestry (MNRF), Peterborough District
4. Natural Heritage Information Centre (NHIC) database (accessed: 2015)
5. Ontario Reptile and Amphibian Atlas (Ontario Nature 2015a)
6. Ontario Reptile and Amphibian Atlas Interactive map (Ontario Nature 2015b)
7. Ontario Mammal Atlas (1994)
8. Atlas of the Breeding Birds of Ontario, 2001-2005
9. Credit River Fisheries Management Plan (CVC 2002)
10. Credit Valley Conservation Terrestrial Monitoring Program Report 2005-2009
11. Wellington County Official Plan, May 6, 1999 (Last Revision March 9, 2015)
12. Credit Valley Conservation Authority Hillsburgh Dam Terrestrial and Aquatic Monitoring Data (provided by CVC, February, 2015)
13. Credit Valley Conservation Authority Regulation Mapping (accessed 2015)
14. Region of Wellington Significant species list (2008)
15. Peel Region Natural Areas Inventory Vol. 1 (2011)
16. County of Wellington Official Plan (2006, last revision 2015)
17. Town of Erin Official Plan (2012b)
18. Town of Erin Servicing and Settlement Master Plan (2015)
19. Town of Erin Servicing and Settlement Master Plan - Phase 1 Environmental Component (2011)

2.2 Vegetation

2.2.1 Ecological Land Classification

Ecological Land Classification (ELC) field investigations were completed from May 13th to September 25th, 2015. Detailed survey dates and weather information are provided in *Appendix 3*. Surveys were completed by qualified ecologists, Ryan Hamelin, OMNRF Certified in Ecological Land Classification and Cheryl-Anne Ross, OMNRF Certified in Ecological Land Classification. Vegetation communities within the study area were characterized and delineated through field investigation, following the Ecological Land Classification (ELC) system for Southern Ontario 1st approximation; community codes used generally follow the 2nd approximation (Lee, et al., 1998, 2008). Boundaries of ELC communities were mapped using aerial images and field observations (*Figure 3*). Detailed descriptions of each ELC community are provided in *Appendix 4*, and digitized ELC data sheets are provided in *Appendix 5*.

Identified ELC communities were cross referenced with the NHIC Ontario Plant Community List (NHIC 2015) to determine the presence of rare plant communities (S3-S1). The Subnational, or Provincial, Ranks (S Rank) are assigned by the Ontario Ministry of Natural Resources and Forestry (MNRF) Natural Heritage Information Centre (NHIC) in order to help assign protection priorities.

2.2.2 Botanical Inventory

2.2.2.1 *About & Associates*

Concurrent with ELC evaluations, the subject lands were systematically searched in order to provide a comprehensive three season botanical inventory. Detailed survey dates and weather information are provided in *Appendix 3*.

Identified vascular plant species were compared to provincial and federal SAR lists (COSARO, SARA) provincial ranks (NHIC 2015), global ranks, CVC list of Species of Conservation Concern Status (CVC 2010b) and Significant Plants of Wellington County (Dougan & Associates 2009), in order to assess federal, provincial, regional and local conservation status of each species. English colloquial names and scientific binomials of plant species generally follow VASCAN (VASCAN 2015).

Identification of environmentally sensitive plant species was completed based on assignment of a coefficient of conservatism value (CC) for each native species (Oldham, et al., 1995). The value of CC, ranging from 0 (low) to 10 (high), is based on a species' tolerance of disturbance and fidelity to specific natural habitat parameters. Species with a CC value of 9 or 10 generally exhibit a high degree of fidelity to a narrow range of habitat parameters. These species may be more sensitive to environmental changes (Mortarello et. al., 2010).

A list of all identified plant species is provided in *Appendix 6*. The list provides botanical name, common name, provincial rarity rank (S-rank), global rarity rank (G-rank), provincial SAR status, federal SAR status, CVC Species of Conservation Status (CVC 2010b), Local rarity/significance within Wellington county (Dougan & Associates 2009), coefficient of conservatism (CC) and coefficient of wetness (CW). Plant species that could only be identified to genus (*Carex sp.*, *Crataegus sp.*) were not assigned the above information.

2.2.2.2 *Credit Valley Conservation Authority*

In addition to surveys completed by About & Associate, data from previous botanical surveys conducted by CVC were also compiled and are provided in *Appendix 6*. CVC data was collected within the study area from 2008 to 2013. Much of the data was gathered from properties within the study area that were not accessible during this EA study due to a lack of landowner permission. The current study includes properties not surveyed as part of the CVC botanical studies. As a result of these differences in property access, the combined data for the two different sources provides a more complete inventory of the study area.

2.3 Provincially Significant Wetlands

The Provincially Significant West Credit Wetland Complex is partly within the study area and comprises a large portion of the natural feature upstream and downstream of the Hillsburgh Dam. The wetland was first evaluated under the Ontario Wetland Evaluation System (OWES) in 1995 by the MNRF, with updates in 2005 (NRVIS 2010). A copy of the wetland evaluation data and scoring record was obtained from the Guelph District MNRF office and reviewed in order to

determine the presence of potentially significant features.

The mapped wetland boundary was obtained from the Land Information Ontario (LIO) online database. The accuracy of the boundary was confirmed through a combination of desktop analysis and field surveys, conducted concurrent with ELC evaluations by Ryan Hamelin, OMNRF Certified in OWES. Detailed survey dates and weather information are provided in *Appendix 3*. The wetland boundary was established where vegetation was comprised of 50% wetland and 50% upland species, and where soils displayed hydric conditions (e.g. presence of mottles and/or gleys), per the *Ontario Wetland Evaluation System* (2013). Due to property access restrictions, it was not possible to confirm the accuracy of the entire wetland boundary within the study area.

2.4 Wildlife

2.4.1 Amphibians (Anurans)

Evening point count surveys to detect breeding calls of anurans (frog and toad) were conducted by Cheryl-Anne Ross, Wildlife Ecologist and Ryan Hamelin, Terrestrial and Wetland Ecologist, in accordance with the *Marsh Monitoring Program Participants Handbook for Surveying Amphibians* (Bird Studies Canada 2008). Three surveys were completed, in accordance with the recommended windows for the spring and early summer, in order to maximize the chances of detecting all potential species. Surveys coincided with optimum weather conditions for anuran breeding activity and detection of calls, i.e. suitable temperature relative to each survey window, humid or damp but not raining, and low wind. Call Level Codes were applied to each species detected per area of suitable habitat, and numbers of individuals were counted or estimated, where applicable. The surveys took place on April 15, May 28 and June 24, 2015. The point Count Locations are illustrated on *Figure 4*; Survey Results and Call Level Code descriptions are provided in *Appendix 7*. Detailed survey dates and weather information are provided in *Appendix 3*.

2.4.2 Breeding Birds

Breeding Bird Surveys were conducted through 10 minute point counts positioned approximately 250m apart within the study area where access was permitted by Cheryl-Anne Ross, Wildlife Ecologist. The highest observed level of breeding evidence was used to assign breeding status (i.e. confirmed, possible, probable or observed) to each species, as per the *Ontario Breeding Bird Atlas: Guide for Participants* (Bird Studies Canada 2001). Marsh Breeding Bird Surveys were completed following each Breeding Bird Survey, at point count locations where habitat was also conducive to marsh birds; methods followed the *Marsh Monitoring Protocols* (Bird Studies Canada 2008). Marsh Breeding Bird Surveys included five minutes of passive listening; five minutes of playing a callback tape of target species, and five further minutes of passive listening, all other bird species observed during the survey were also recorded, including incidental species and aerial foragers.

As per the OBBA and MMP recommendations, two surveys were performed during the peak breeding season for the bulk of species in Southern Ontario (May 24 to July 10), and were spaced at least 10 days apart in order to determine presumed permanent territories through territorial singing males. The two surveys took place on the mornings of June 11 and July 9, 2015. The point Count Locations, including Marsh Breeding Bird Stations are illustrated on *Figure 4*, breeding bird survey results and breeding evidence codes are provided in *Appendix 8*, marsh breeding bird survey results are provided in *Appendix 9*. Detailed survey dates and weather information are provided in *Appendix 3*.

2.4.3 Snakes

Visual encounter and active hand search surveys occurred between April and May 2015 in all candidate habitats identified during initial ELC screening and site visit by Cheryl-Anne Ross, Wildlife Ecologist and Ryan Hamelin, Terrestrial and Wetland Ecologist. Three surveys two weeks apart, were undertaken and included flipping any natural or naturalized cover, identified in the project location. Surveys were undertaken on sunny days when air temperatures were between 8°C and 25°C and on overcast days when air temperatures were above 15°C. Surveys followed pre-determined transects, traversing areas of suitable habitat for both Eastern Ribbonsnake (*Thamnophis sauritus*) and Milksnake (*Lampropeltis Triangulum*). Transect locations are illustrated on *Figure 4*, survey results are provided in *Appendix 10*. Surveys generally followed methods outlined in the Milksnake Survey Protocol (MNRF 2013). Detailed survey dates and weather information are provided in *Appendix 3*.

2.4.4 Salamanders

Visual surveys for candidate vernal pools were undertaken in early April by Cheryl-Anne Ross, Wildlife Ecologist, to determine the presence or absence of candidate habitat for salamander species that may occur in the study area. These surveys were conducted to determine the possible presence of Jefferson Salamander (*Ambystoma jeffersonianum*) habitat within the study area. Since no vernal pools were identified, further visual inspections were not required. Detailed survey dates and weather information are provided in *Appendix 3*.

2.4.5 Turtles

Five basking surveys in candidate habitats within the project location were conducted by Cheryl-Anne Ross, Wildlife Ecologist and Ryan Hamelin, Terrestrial and Wetland Ecologist, in 2015 following the MNRF Guelph District Blanding's survey protocol (2012). Basking surveys, including overwintering (late March-early April) and summer habitat (late April-June 15), were conducted at all waterbodies and wetlands with open water. Locations of candidate habitat are illustrated on *Figure 4*.

All shorelines and potential basking sites in the project location were surveyed from the sunlit side using 8x power binoculars and a stationary 50x maximum power spotting scope. Between late March and early May, surveys were conducted between 9am and 5pm (When temperatures were between 6°C and 10°C, surveys occurred on sunny days with no wind. When temperatures were between 10°C and 25°C, surveys were conducted between 9am and noon on sunny days). Between late May and early June turtles are less reliably found late in the day, as a result surveys occurred between 9am and 12pm. Survey Methods generally followed the MNRF - Guelph District Blanding's Turtle (*Emydoidea blandingii*) Protocol (MNRF 2012). Survey results are provided in *Appendix 11*. Detailed survey dates and weather information are provided in *Appendix 3*.

2.4.6 Winter Wildlife

A Winter Wildlife Survey was undertaken on February 25th 2015 by Ryan Hamelin, Terrestrial and Wetland Ecologist and Matt Isles, Wildlife Ecologist. Detailed survey dates and weather information are provided in *Appendix 3*. Wildlife sightings and evidence such as tracks, scat, vocalizations and markings were used to determine species presence. Notes and GPS points were taken for each observation. Snow depth in the study area was approximately 0.45m up to 0.75m in snow drifts. There was light snow of less than 1 cm during the survey and in the proceeding 24 hours. Approximately 3-7 cm of fresh snow cover fell in the 48 hours prior to the survey.

Due to property access restrictions, the full study area was not surveyed. Where property access was granted, areas were extensively surveyed on foot and with the aid of snowshoes. A road side survey of the study area was also completed where possible. As part of the Winter Wildlife survey, particular effort was applied to locating and identifying raptors, mammal tracks, stick nests, raptor wintering areas, and deer congregation areas. The path traveled during the winter wildlife survey, including roadside driving route is shown on *Figure 4*. All wildlife observations are presented in *Appendix 12*.

2.4.7 Migratory Birds

An assessment for candidate migratory bird habitat and migratory shorebird habitat was completed within the study area, using criteria and guidance from the SWH EcoRegion Criterion Schedule 6E (2015). An assessment of the habitat in the study area that was identified as candidate shorebird migratory staging and stopover was completed on August 5, 2015 and a migratory bird survey of all accessible lands was conducted on October 8, 2015 to determine if the area had significant numbers of migratory species. Detailed survey dates and weather information are provided in *Appendix 3*. Migratory Bird Survey and Shorebird Habitat Assessment Results are provided in *Appendix 13* and *Appendix 14*, respectively.

2.4.8 Incidental Wildlife Observations

Incidental observations of insects, mammals and reptiles were recorded during all field visits, in addition to incidental observations of birds, turtle and amphibians made outside of the formal field surveys for these groups of fauna. Detailed survey dates and weather information are provided in *Appendix 3*. A complete list of all incidental wildlife is provided in *Appendix 15*.

2.5 Significant Wildlife Habitat

With guidance from the *Significant Wildlife Habitat Technical Guide* (2000) and the SWH EcoRegion Criterion Schedule 6E (2015), the study area and adjacent lands were considered for the presence of Significant Wildlife Habitat (e.g. specialized habitats for wildlife, habitat for species of conservation concern). Detailed survey dates and weather information are provided in *Appendix 3*. An assessment of the study area for all SWH is provided in *Appendix 16*.

2.6 SAR Habitat Assessment

A thorough review of all background documents was conducted to compile a master list of all Species at Risk, and species with conservation designation that may occur in the study area. A review of the site, along with habitat requirements for each species was conducted; the site was then evaluated for potential habitat using Ecological Land Classification, guidance from MNR documents, and on-site knowledge acquired through field surveys. Detailed survey dates and weather information are provided in *Appendix 3*. An assessment of the study area of candidate habitat for SAR is provided in *Appendix 17*.

2.7 Aquatic Habitat Assessment

On October 19th, 2015 an Aquatic Habitat Assessment was completed by Ryan Hamelin, Terrestrial and Wetland Ecologist, for all sections of watercourses in the study area, as well as sections of the watercourse directly up-stream and down-stream of the study area. Detailed survey dates and weather information are provided in *Appendix 3*. The Aquatic Habitat Assessment was completed in order to determine the quality of habitat to fish, barriers to fish

movement, and general aquatic habitat characteristics. For the assessment, the watercourse was separated into 14 segments and each characterized with respect to the following criteria:

- Mean channel width;
- mean channel depth;
- mean water depth;
- percent stream shading;
- buffer width;
- substrate;
- flow pattern;
- channel morphology;
- instream cover;
- bank characteristics;
- presence of specific site features.

In addition to the field Aquatic Habitat Assessment, data provided by the MNR and CVC such as fish collection records, CVC water temperature data and thermal fish community classification information was used to characterize each segment of watercourse. Locations of specific fish collection records from the MNR and CVC were used to identify fish species known to be present in each watercourse segment.

Summer water temperatures of the tributaries and open water communities within the study area were collected during 2013 and 2014 by CVC. These data were provided to Aboud & Associates to assist in the assessment of the temperature regimes of aquatic habitat. The data provided did not have complete coverage of the study area and could not be used to determine the temperature regime of all watercourse segments. Where data allowed, water temperature regime was calculated using the definitions provided in *A Guide to Understanding Freshwater Fish Habitat in Ontario* (DFO 2008): where cold water systems are generally below 19°C during summer maximum temperatures, cool water systems are characterized by maximum summer water temperatures between 19 - 25°C and warm water systems are characterized by maximum summer water temperatures above 25°C.

An alternative to the above water temperature regime classification is fish community classification based on thermal preference. A fish community classification was completed for the Erin Service and Settlement Master Plan- Phase 1: Environmental Component Report (ESSMP 2011) by CVC that included all watercourses in the study area. This fish community classification was used in the aquatic habitat assessment to classify stream segments as cold, cool or warm water systems.

Areas of potential trout spawning habitat, barriers to fish passage, fish community classification and other relevant information is presented on *Figure 5. Survey Results* are provided in *Appendix 18*.

2.8 Landscape Evaluation

A landscape level evaluation was completed for the study area and surrounding lands to identify ecologically significant features that extend beyond the boundaries of the study area, and that may be impacted by changes within the study area. The following background resources were reviewed in completing the Landscape Evaluation:

- Erin Service and Settlement Master Plan – Phase 1: Environmental Component (2011);
- The Credit River Fisheries Management Plan (2002);
- Natural Heritage Information Center (NHIC);
- West Credit River Wetland Evaluation Score Card;
- The Ecosystems of Ontario, Part 1 : Ecozones and Ecoregions (MNR 2009);
- Aerial photo interpretation.

DRAFT

3.0 Existing Conditions

Information that characterizes the existing conditions of the study area came from several sources, including but not limited to, background review of existing documents, public information sources, past field studies by others, and extensive field reconnaissance.

3.1 Background Review

3.1.1 Natural Heritage Information Centre - Species at Risk

Preliminary investigation through the Natural Heritage Information Centre (NHIC) identified two provincial Species at Risk (SAR) under the ESA and two species considered rare in the province (S1-S3) recorded within approximately 1km of the study area. These species and their habitat requirements are summarized in *Table 1*.

Table 1. NHIC Species at Risk Records

Scientific Name	Common Name	(COSEWIC) Status ¹	(SARO) Status ²	Last Observed (NHIC)	S-Rank ³	Habitat Requirements
<i>Dolichonyx oryzivorus</i>	Bobolink	Threatened	Threatened	June 2, 2001	S4B	Nest in grassland habitats, including hayfields and meadows with a mixture of grasses and broad-leaved forbs with a high litter cover. Area Sensitive, with increased density in grasslands greater than 10ha (Renfrew et. al. 2015)
<i>Sturnella magna</i>	Eastern Meadowlark	Threatened	Threatened	June 2, 2001	S4B	Nest in grassland habitats, including hayfields, pasture, savannahs, and other open areas. Preferential habitat includes areas with good grass and thatch (litter) cover (Jaster et. al. 2012).
<i>Carex careyana</i>	Carey's Sedge	Not listed	Not listed	June 14, 1977	S2	Grows in dry to moist rich deciduous upland forests (NatureServe 2015)
<i>Sceptridium rugulosum</i>	Rugulose Grapefern	Not listed	Not listed	Nov. 15, 1977	S2	Grows in sandy to silty soil in open fields, young successional forests or at the edge of forests (Wagner and Wagner 1982).

¹ COSEWIC – Committee on the status of endangered wildlife in Canada

² SARO – Species at Risk Act Ontario

³ S-Rank – Denotes the conservation status of a species at the provincial level

S2: Imperiled

S4: Apparently Secure—Uncommon but not rare

S#B- Breeding status rank

3.1.2 Ontario Breeding Bird Atlas

A list of birds determined to be breeding (Possible, Probable or Confirmed) in the 10km x 10km square containing the study area during the 2001-2005 Ontario Breeding Bird Atlas was compiled. This list includes 107 species; eight are considered Species at Risk under the ESA. Potential breeding habitat was identified in the study area for three of these species (Eastern Wood-pewee (*Contopus virens*), Barn Swallow (*Hirundo rustica*), and Canada Warbler (*Cardellina Canadensis*)). Nine of the species identified in the square are considered Species of Conservation Concern Status by CVC (Tier 1) and 51 are considered significant species in Wellington County (Dougan & Associates 2009). The findings of this review are presented in *Appendix 19*. Species with conservation designation identified in the background review and their habitat requirements are presented in *Appendix 17*.

3.1.3 Ontario Reptile and Amphibian Atlas

Review of the Ontario Reptile and Amphibian Atlas identified seven species that are known to occur within the 10km x 10km square containing the study area. This list includes one species at risk under the ESA; Common Snapping Turtle (*Chelydra serpentina*) is listed as special concern provincially and federally. Confirmed nesting or overwintering habitat was identified on the subject parcel for this species.

One of the species known to occur in the square is considered a Species of Conservation Concern by CVC (Tier 1) and one is considered a significant species in Wellington County (Dougan & Associates, 2009). The findings of this review are presented in *Appendix 19*. Species with conservation designation identified in the background review and their habitat requirements are presented in *Appendix 17*.

3.1.4 Atlas of the Mammals of Ontario

Review of the Atlas of the Mammals of Ontario (1994) identified twenty-five species that are known to occur within approximately 10km of the study area. This list includes one species at risk under the ESA; Little Brown Myotis (*Myotis lucifugus*) is listed as endangered provincially and federally. Potential habitat was identified in the study area for this species.

One of the species known to occur in the square is considered a CVC Species of Conservation Concern (Tier 1) and one is considered a significant species in Wellington County (Dougan & Associates, 2009). The findings of this review are presented in *Appendix 19*. Species with conservation designation identified in the background review and their habitat requirements are discussed in *Appendix 17*.

3.1.6 Credit Valley Conservation

3.1.6.1 Botanical Surveys

CVC provided a list of plant species identified from within the study area. The majority of the observations are from the southern portion of the study area, with only a few observations from the north side of Station Street Road. The data was collected from 2008 to 2014; specific sampling methods were not provided. A total of 320 plant species or distinct sub-species were included in the list. Georeferenced location data was provided for some observations. None of the species identified by CVC are listed as provincial or federal species at risk. Eight of the species are considered rare in Wellington County (Dougan & Associates, 2009) with 70 species are considered Species of Conservation Interest (Tier 2) by CVC.

All but one of the native plants identified are ranked as secure in Ontario (S5) or apparently secure (S4) and globally very common (G5) or common (G4) (NHIC, 2015). The one exception is *Fontinalis sullivantii*, a moss species which is classified as an S1 (Extremely rare in Ontario); location and population detail of the species was not provided. All plant species identified by CVC are included in *Appendix 6*.

Since all observations of plant species identified by CVC are from within the Study and there is a large overlap in identified species between the CVC list and the EA field studies, a further analysis using the combined CVC data and field data collected by AA is provided in section 3.2.2.

3.1.6.2 Fish Surveys

CVC provided a georeferenced list of fish species identified within the study area. Fish species data was compiled between 1954 and 2013; specific sampling methods were not provided. Numbers of individual fish observed are provided for some sampling data.

The list contains 16 species, none of which are considered provincial or federal SAR. Three of the species; Slimy Sculpin, Banded Killifish and Brook Trout are listed as CVC Species of Interest (Tier 2). All species observed are included in *Appendix 20*.

3.1.6.3 Breeding Bird Surveys

A list of birds determined to be breeding (Possible, Probable or Confirmed) in the study area during the 2009 Breeding Bird Surveys, completed by CVC was compiled. This list includes 51 species; none are listed as SAR. None of the species determined to be breeding in the square are considered Species of Conservation Concern by Credit Valley Conservation (Tier 1), eight CVC Species of Interest (Tier 2), were observed, and 12 are considered significant species in Wellington County (Dougan & Associates, 2009). The findings of this review are presented in *Appendix 19*.

3.1.6.4 Incidental Wildlife Observations

A list of all fauna observations made in the area of the Hillsburgh Pond was compiled by CVC and provided as a background source for the study area. All observations occurred between 2003 and 2014 and are provided in *Appendix 19*. This list includes 19 species observed in the study area outside of formal surveys; none of these species are listed as SAR, one CVC Species of Conservation Concern Tier 1 species, Great Egret (*Ardea alba*), was observed, six Tier 2 species were observed, and five are considered significant in Wellington County (Dougan & Associates, 2009).

3.1.6.5 Spring and Fall Migration Surveys

CVC completed spring and fall migration surveys during 2012, 55 species were observed in the study area during investigations. None of the species observed are considered Species of Conservation Concern. Two CVC Tier 1 species were observed, Canvasback (*Aythya valisineria*) and Trumpeter Swan (*Cygnus buccinator*). A further 13 Tier 2 species were observed and are listed in *Appendix 19*.

3.1.6.6 Significant Wildlife Habitat Survey - Waterfowl Staging - Aquatic

CVC completed 13 waterfowl staging surveys in 2011, a total of 45 species were observed during the surveys, and included 13 of the SWH listed species. During the surveys, three days met the criteria of having greater than 100 individuals observed; 7 or more days of 100 individual listed species are required to meet the criteria for SWH. All species observed are included in *Appendix 19*.

3.1.7 Ministry of Natural Resources and Forestry

3.1.7.1 Little Brown Myotis Maternity Exit Surveys

The Peterborough district MNRF have been conducting exit surveys and banding Little Brown Myotis (bats) at a property adjacent to the study area since 2012 (pers. Comm. Lesley Hale, 2015). Little Brown Myotis is listed as Endangered provincially and federally, as such, they are afforded general habitat protection. Over the course of the surveys, the maternity population has increased. The MNRF identified that the Hillsburgh Pond may provide important foraging habitat for this maternity colony of Little Brown Myotis.

3.1.7.2 Fish Records

Fish data collection records from within the study area were provided by the Guelph District MNRF. The data were collected through a combination of electrofishing, drift nets, minnow traps and incidental observations during 2013 and 2014. 10 species were identified, all of which were also identified by CVC. All species observed are included in *Appendix 20*. No provincial or federal SAR was identified.

3.2 Vegetation

3.2.1 Ecological Land Classification

A three season ELC evaluation was completed in 2015 by Aboud & Associates. 31 ELC communities were identified and mapped in the study area. The community polygons identified during the ELC surveys are summarized in *Table 2*. Digitized Field forms are provided in *Appendix 5* with detailed ELC descriptions provided in *Appendix 4*. Comparison with the NHIC Rare Plant Communities did not identify any provincially rare plant communities (S1 – S3) within the study area. ELC communities are shown on *Figure 3*.

Table 2. Ecological Land Classification		
ELC Code ¹	Vegetation Type	Map ID
Mixed Meadow (MEM)		
MEMM3	Dry - Fresh Mixed Meadow Ecosite	12
Coniferous Forest (FOC)		
FOCM2-2	Dry-Fresh White Cedar Coniferous Forest	5
FOCM6	Naturalized Coniferous Plantation	27, 6
Mixed Forest (FOM)		
FOMM7-2	Fresh - Moist White Cedar - Hardwood Mixed Forest	23
Deciduous Forest (FOD)		
FODM5-8	Dry-Fresh Sugar Maple - White Ash Deciduous Forest	4
FODM6	Fresh - Moist Sugar Maple Deciduous Forest Ecosite	16
FODM7-7	Fresh - Moist Manitoba Maple Lowland Deciduous Forest	30
FODM8-1	Fresh - Moist Poplar Deciduous Forest	25, 15
Coniferous Forest (FOD)		
SWCM1-2	White Cedar - Conifer Mineral Coniferous Swamp	2, 17, 21
Mixed Swamp (SWM)		
SWM01-1	White Cedar - Hardwood Organic Mixed Swamp	10
SWM03-3	White Birch - Conifer Organic Mixed Swamp	3
Deciduous Swamp (SWD)		
SWDM2-1	Black Ash Mineral Deciduous Swamp	26
SWDM4-5	Poplar Mineral Deciduous Swamp	24, 29
Thicket Swamp (SWT)		
SWT02-3	Meadow Willow Organic Deciduous Thicket Swamp	28
SWT02-6	Mixed Willow Organic Thicket Swamp Type	22
SWT03-5	Red-osier Organic Deciduous Swamp	9
Treed Fen (FET)		
FETC1-2	Tamarack - White Cedar Treed Fen	14
Meadow Marsh (MAM)		
MAMM1-1	Cattail Graminoid Mineral Meadow Marsh Type	31
MAMO1-2	Cattail Graminoid Organic Mineral Meadow Marsh	1

Table 2. Ecological Land Classification		
ELC Code ¹	Vegetation Type	Map ID
Shallow Marsh (MAS)		
MAS01-1	Cattail Organic Shallow Marsh Type	8
Submerged Shallow Aquatic (SAS)		
SAS_1	Submerged Shallow Aquatic Ecosite	7
Mixed Shallow Aquatic		
SAM_1-8	Water Lily - Bullhead lily Mixed Shallow Aquatic	11
SAM_1-8	Water Lily - Bullhead lily Mixed Shallow Aquatic	19
SAM_1-8	Water Lily - Bullhead lily Mixed Shallow Aquatic	18
Open Aquatic (OAO)		
OAW	Open Aquatic	20
Cultural (CU)		
CS	Cultural Savannah	13
Constructed (CV)		
CVR_1	Residential	Res
CVI_1	Transportation	Road

¹ ELC Codes generally follows the ELC Second Approximation (Lee 2008)

3.2.2 Botanical Inventory

A detailed field inventory of accessible properties within the study area was completed and 299 species or distinct sub-species of vascular plants, from 75 families, were identified. All identified plant species are provided in *Appendix 6*. A further 7 species were identified only to the level of genus and have not been designated as native or non-native or included in the overall species count.

The provided CVC plant data was collected from within the study area, much of it from properties where access was restricted for this study, the combined data provides a more complete inventory of the entire study area. Including the CVC data, a further 95 species or distinct sub-species from 11 additional plant families were identified, for a total of 394 species or sub-species from 87 families within the Study Area; of those, 284 species (72%) are native and 110 species (28%) are exotic.

3.2.1.1 Species at Risk, Regional and Local Significance

All but one of the native vascular plants observed in the study area, or identified in CVC data are ranked as secure in Ontario (S5) or apparently secure (S4) and globally, very common (G5) or common (G4) (NHIC 2015). A moss species (*Fontinalis sullivantii*), one of the species identified by CVC, is classified as an S1 (Extremely rare in Ontario), location and population detail of the species was not provided. One distinct sub-species, Tuberous White Water-lily (*Nymphaea odorata ssp. Tuberosa*) was identified by CVC along with the more common White Water-lily (*Nymphaea odorata ssp. Odorata*). Tuberous White Water-lily is provincially unranked, but considered native by CVC (2015).

Ten identified species are considered significant in Wellington County (Dougan & Associates et. al. 2009). 77 of the identified species are classified as Species of Interest (Tier 2) in the CVC Species of Conservation of Concern Project; no Tier 1 species were identified.

Six of the species observed in the study area, or identified in CVC data had a Co-efficient of conservatism of 9 or 10. These species include: Marsh Horsetail (*Equisetum Palustre*) (CC 10); Three-seed Sedge (*Carex trisperma*) (9); Hooded Ladies'-tresses (*Spiranthes romanzoffiana*) (9); Bog Goldenrod (*Solidago uliginosa*) (9); Kalm's Lobelia (*Lobelia kalmia*) (9); Green Keeled Cottongrass (*Eriophorum viridicarinatum*) (9).

3.3 Provincially Significant Wetlands

3.3.1 Boundary Review

The mapped wetland boundary of the West Credit River Wetland Complex was accessed through Land Information Ontario (LIO). The accuracy of the boundary within the study area was reviewed through field survey and ortho-photograph interpretation to determine any discrepancies and update the current boundary.

The boundaries review determined that the provided wetland boundary was generally accurate with only a few minor deviations from the actual boundary in the field. The wetland boundary as provided by LIO showed a total of 44.74ha of wetland within the study area. The boundary verification identified 0.09ha of additional wetland and 1.07ha of area was incorrectly identified as wetland. *Figure 6* shows the wetland boundary as provided by LIO. Inaccuracies in the wetland boundary have not been field verified or confirmed by the CVC or MNRF.

3.3.2 Wetland Characteristics

The LIO wetland file identifies the wetland complex within the study area as containing Swamp, Marsh and Open Water wetland types. This is consistent with the ELC survey which identified; Coniferous Swamp, Mixed Swamp, Deciduous Swamp, Thicket Swamp, Treed Fen, Meadow Marsh, Shallow Marsh, Submerged Shallow Aquatic, Mixed Shallow Aquatic, and Open Aquatic communities within the study area.

Review of the Wetland Evaluation Data and Scoring Record identified that the wetland complex scored the maximum points for flood attenuation, indicating that the wetland is an important feature in reducing the risk of flooding. Flood risk was identified by CVC as an important criterion to consider when identifying the preferred alternatives (Pers. Comm., T. Slaught, 2014). The wetland also scored the maximum number of points for erosion control, a criteria also identified by CVC as important when considering preferred options (Pers. Comm., T. Slaught, 2014).

3.4 Wildlife

3.4.1 Amphibians (Anurans)

The results of the Anuran Point Count Surveys are summarized in *Table 3*, and results are discussed below. The Point Count Locations are illustrated on *Figure 4*, and Call Level Code descriptions, along with the complete survey results, are provided in *Appendix 7*.

Table 3. Summary of Amphibian Observations (2015)

SPECIES	AMPHIBIAN HABITAT						
	1 (C1, G, F)	2 (D)	3 (C2)	4 (E)	5 (B1)	6 (B2)	7 (A)
Gray Treefrog	25	4	3		2		1
Spring Peeper	8	Chorus	12	1	19	4	3
Green Frog	8	2	2				6
Northern Leopard Frog			2				
Wood Frog		3	1		10		
Significant Habitat*	Y	Y	Y	N	Y	N	N

*Significance: Y-Indicates Amphibian Habitat meets the criteria listed under the Ecoregion 6E SWH Criteria guide (2015). N-Indicates Amphibian Habitat did not meet the criteria listed under the Ecoregion 6E SWH Criteria guide (2015).

Amphibian Habitat 1

Three species of frog were detected calling from within Amphibian Habitat 1. This site targeted the Hillsburgh Pond from three locations (point count locations C1, G and F), at distances of at least 250m apart. One species, Gray Treefrog (*Hyla versicolor*), had greater than 20 individuals. All frog species were heard calling from the edges of the pond, particularly the south and north shoreline, which includes abundant aquatic vegetation. The Hillsburgh Pond meets the criteria for Significant Wildlife Habitat-Amphibian breeding (woodland), as there were greater than 2 species observed and greater than 20 individuals detected.

Amphibian Habitat 2

Four Species of frog were detected calling from within Amphibian Habitat 2. This site targeted the swamp thicket habitat (point count location D), located in the western portion of the study area, north of the Elora-Cataract Trail. One species, Spring Peeper (*Pseudacris crucifer*), was estimated to have greater than 20 individuals. All frog species were heard calling from within a shallow thicket swamp community. The swamp thicket meets the criteria for Significant Wildlife Habitat-Amphibian breeding (woodland), as there were greater than 2 species observed and greater than 20 individuals detected.

Amphibian Habitat 3

Five Species of frog were detected calling from within Amphibian Habitat 3. This site targeted the shallow pond (point count location C2), located in the western portion of the study area, south of the Elora-Cataract Trail. None of the species observed had greater than 20 individuals. The shallow pond meets the criteria for Significant Wildlife Habitat-Amphibian breeding (woodland), as there were greater than 2 listed species observed and 20 individuals detected.

Amphibian Habitat 4

One species of frog was detected calling from within Amphibian Habitat 4. This site targeted the east side of the Hillsburgh Dam (point count location E). One species, Spring Peeper (one individual), was heard calling from the edges of the feature. The east side of the dam does not meet the criteria for Significant Wildlife habitat-Amphibian breeding (woodland).

Amphibian Habitat 5

Three species of frog were detected calling from within Amphibian Habitat 5. This site targeted the Cattail marsh, on the east side of the study area (point count location B1), north of the Elora-Cataract Trail. The cattail marsh meets the criteria for Significant Wildlife habitat-Amphibian

breeding (woodland), as there were greater than 2 species observed and greater than 20 individuals detected.

Amphibian Habitat 6

One species of frog was detected calling from within Amphibian Habitat 6. This site targeted the Eastern White Cedar swamp, east of the Hillsburgh Dam (point count location B2), and south of the Elora-Cataract Trail. One species, spring peeper (four individuals), was heard calling from the edges of the feature. The Spruce swamp does not meet the criteria for Significant Wildlife habitat-Amphibian breeding (woodland).

Amphibian Habitat 7

Three species of frogs were detected calling from within Amphibian Habitat 7. This site targeted the Rudd Pond and the wetland to the north (point count location A). All frog species were heard calling from the North West edge of the pond, where there is abundant aquatic vegetation. The pond does not meet the criteria for Significant Wildlife habitat-Amphibian breeding (woodland), as there were less than 20 individuals detected.

3.4.1.1 Amphibian SAR, Regional and Local Significance

No amphibian species observed are considered federal or provincial species at risk.

All species detected calling within the study area are ranked S5 (Very Common) in Ontario (NHIC, 2015).

One species, Wood Frog (*Lithobates sylvaticus*), is ranked as CVC Species of Interest (Tier 2), all other species observed are ranked as Tier 3; Species of Urban Interest.

3.4.2 Breeding Birds

The results of the Breeding Bird Survey (BBS) are presented in *Appendix 8*. Locations of significant observations are provided in *Figure 7*, and are approximate. They are designed to give a general indication of the area in which the species may be nesting. During BBS visits, a total of 47 species were detected, of which five were assigned 'confirmed' breeding evidence, sixteen were assigned 'Probable', twenty-one were assigned 'possible' and four showed no sign of breeding evidence observed. All but one species, Eastern Meadowlark (*Sturnella magna*), were detected within the study area. During Marsh Breeding Bird surveys, no target marsh bird species were detected, a list of secondary species and aerial foragers observed is provided in *Appendix 9*.

Due to the contiguity with natural lands to the south and north, it is important to note that, despite high levels of breeding evidence, a given species may not have been breeding specifically in the area in which it was observed. This is particularly true where species were only detected during one of the two Breeding Bird Surveys. These species may have been foraging in these areas or, may have been wandering during post-breeding dispersal. Therefore, the following 21 species are those that can be presumed to have been breeding in, or within 30m of, the study area, and exhibited confirmed or probable breeding evidence: Mallard (*Anas platyrhynchos*), Belted Kingfisher (*Megaceryle alcyon*), Downy Woodpecker (*Picoides pubescens*), Northern Flicker (*Colaptes auratus*), Eastern Wood-pewee, Great Crested Flycatcher (*Myiarchus crinitus*), Eastern Kingbird (*Tyrannus tyrannus*), Blue Jay (*Cyanocitta cristata*), American Crow (*Corvus brachyrhynchos*), Black-capped Chickadee (*Poecile atricapillus*), House Wren (*Troglodytes aedon*), American Robin (*Turdus migratorius*), Warbling Vireo (*Vireo gilvus*), Red-eyed Vireo (*Vireo olivaceus*), Yellow Warbler (*Setophaga petechial*),

Common Yellowthroat (*Geothlypis trichas*), Song Sparrow (*Melospiza melodia*), Swamp Sparrow (*Melospiza georgiana*), Red-winged Blackbird (*Agelaius phoeniceus*), Common Grackle (*Quiscalus quiscula*), and Baltimore Oriole (*Icterus galbula*).

Most of the species presumed to be breeding in the study area, are considered common and abundant species (S-Rank 4-5, CVC Tier 3-5).

3.4.2.1 Breeding Bird Species at Risk

Two species observed are considered species at risk under the ESA. Eastern Meadowlark is listed as Threatened provincially and federally and Eastern Wood-pewee is listed as Special Concern provincially and federally, locations of observations are shown on *Figure 7*.

Eastern Meadowlark is an area sensitive, grassland species, often nesting in hay fields and pastures, as well as occasionally occurring in other types of grassed areas such as golf courses, and airfields. The grassland habitat requires a moderate thatch cover, low shrub and tree density, and moderate or limited forbs cover. Large tracts of grassland are typically preferred over smaller patches (McCracken et. al. 2013). A single male Eastern Meadowlark was observed singing from adjacent lands outside the study area during one breeding bird survey (*Figure 7*), in habitat that may be sufficient for establishing a territory. No habitat of sufficient size, or matching criteria was observed in the study area.

Eastern wood-pewee are associated with mid-age mixed and deciduous forest stands, often dominated by Maple (*Acer*), Elm (*Ulmus*) or Oak (*Quercus*), and include areas with clear-cuts, openings or forest edges. Eastern Wood-pewee also prefers forest stands with little to no understory vegetation (COSEWIC 2012). Eastern Wood-pewee was observed singing during both breeding bird surveys in the deciduous forest communities in the south-eastern portion of the study area (*Figure 7*).

3.4.2.2 Breeding Bird Regional and Local Significance

All species detected in the study area are ranked as either S5 (Very Common) or S4 (Common) or in Ontario. The rank qualifier 'B' denotes the status of a migratory species during the breeding season.

Five species ranked Tier 1 or Tier 2 were observed in the study area, two of which showed probable breeding evidence and are described in *Table 4*.

The County of Wellington has identified a number of species considered significant (Dougan & Associates, 2009). 24 regionally significant species were observed in the study area, 11 of which showed probable or confirmed breeding evidence, locations of species observed and their status are described in *Table 4*.

Table 4. Regionally or Locally Significant Breeding Bird Species

COMMON NAME	SCIENTIFIC NAME	WELLINGTON COUNTY ¹	CVC TIER ²	LOCATION(S) IN STUDY AREA
Eastern Wood-pewee	<i>Contopus virens</i>	✓	1	Observed in the deciduous forest communities located in the south-eastern portion of the study area (Figure 7).
American Crow	<i>Corvus brachyrhynchos</i>	✓	2	Observed at most point counts throughout the study area and is ranked Tier 2.
Mallard	<i>Anas platyrhynchos</i>	✓	3	Observed in the shallow pond community, south of the Elora-Cataract Trail.
Belted Kingfisher	<i>Megaceryle alcyon</i>	✓	3	A pair was observed foraging over the Rudd Pond in the eastern portion of the study area south of the trail.
Downy Woodpecker	<i>Picoides pubescens</i>	✓	3	Observed drumming in the vicinity of the cattail marsh on the edge of the Hillsburgh Pond.
Northern Flicker	<i>Colaptes auratus</i>	✓	3	Observed in numerous locations in the study area drumming and calling.
House Wren	<i>Troglodytes aedon</i>	✓	3	Observed along the trail in the eastern portion of the study area.
Red-eyed Vireo	<i>Vireo olivaceus</i>	✓	3	Observed in both the deciduous forest communities and the mixed swamp community in the eastern portion of the study area.
Common Yellowthroat	<i>Geothlypis trichas</i>	✓	3	Observed along the trail in western portion of the study area, adjacent to the shallow marsh and swamp thicket communities.
Song Sparrow	<i>Melospiza melodia</i>	✓	3	Observed singing throughout the study area in most habitats.
Red-winged Blackbird	<i>Agelaius phoeniceus</i>	✓	3	Observed exhibiting agitated behavior in the area of the cattail marshes at the Hillsburgh pond, and the Ainsworth pond, south of the dam.

¹ Wellington County Significant Plants of Wellington County (Dougan & Associates 2009)

² CVC Species of Conservation Concern Project (CVC 2010b)

3.4.2.2 Breeding Bird Regional Priority Species

The Ontario Landbird Conservation Plan (OLCP): Lower Great Lakes/St. Lawrence Plain, North American Bird Conservation Region 13 (Ontario Partners in Flight, 2008) has identified a number of species that are considered conservation priorities for the region. Six priority species were observed in the study area, including Belted Kingfisher, Northern Flicker, Eastern Wood-pewee, Rose-breasted Grosbeak, Eastern Meadowlark, and Baltimore Oriole. The OLCP does not provide legislative protection of species or their habitat, but rather identifies species that should be conservation priorities on a regional level that were not designated Species at Risk at the time of writing.

3.4.3 Snakes

During snake transect surveys; one individual Eastern Garter Snake (*Hamnophis sirtalis sirtalis*) was detected sunning along the edge of the Elora-Cataract Trail, east of the Dam. During the three rounds of surveys, no other snakes were detected. Two areas of candidate hibernacula habitat were identified in the study area (Appendix 10). These areas included numerous piled stones and rubble in the Naturalized Conifer Plantation in the far eastern portion of the study area (Image 1 and 2). However no snakes were observed in the general area of the candidate hibernacula habitat during transect surveys.



Image 1. Candidate Snake Hibernacula A.

Image 2. Candidate Snake Hibernacula B.

3.4.3.1 Snake SAR, Regional and Local Significance

No snake species observed is considered federal or provincial species at risk. Eastern garter snake is ranked S5 (Very Common) in Ontario (NHIC 2015).

3.4.4 Salamanders

During spring surveys a thorough search of the study area for evidence of habitat that may be suitable for salamanders was conducted. No salamander breeding habitat was observed in the study area, where access was provided. Air photo interpretation and ELC surveys revealed no candidate salamander breeding habitat in study area. As a result, no further detailed studies were conducted.

3.4.5 Turtles

The results of the turtle basking surveys are presented in *Appendix 11* and summarized in *Table 5*. Locations of significant observations are provided in *Figure 7*. They are designed to give a general indication of the area in which the species was observed. These locations are also the areas where turtles are likely to overwinter and/or use as summer habitat. During turtle surveys, two species were observed, Common Snapping Turtle and Midland Painted Turtle (*Chrysemys picta marginata*); one unknown species was also observed, an unconfirmed Red-eared Slider (*Trachemys scripta elegans*). An Assessment of Significance is provided in *Table 5*.

One turtle nest, identified as Common Snapping Turtle, was observed in a man-made wood chip berm, along the eastern edge of the Rudd Pond, confirming breeding in this area.

Table 5. Turtle Habitat Results

Species	Turtle Habitat 1	Turtle Habitat 2	Turtle Habitat 3	Turtle Habitat 4	Turtle Habitat 5
<i>Common Snapping Turtle</i>	9		6		
<i>Midland Painted Turtle</i>	104	1	445	7	1
<i>Unknown Turtle Species</i>				1	
Grand Total*	113	1	451	8	1

* Total number of turtles observed summed over all survey dates conducted for each habitat.

Turtle Habitat 1

Turtle Habitat 1, the Rudd Pond is located in the south-eastern portion of the study area, and was surveyed five times between April and June, 2015. Over the course of 5 surveys a total of 113 turtles were observed, consisting of Midland Painted Turtle and Common Snapping Turtle. Within Turtle Habitat 1, two individual and three individual Common Snapping Turtles were observed during April and May surveys, respectively. The Ecoregion 6E SWH criteria (MNR 2015b), states that: any area with at least five overwintering (observed between March and May) Midland Painted Turtles or one Common Snapping Turtle is considered SWH. Turtle Habitat 1 meets the criteria for SWH.

Turtle Habitat 2

Turtle Habitat 2, the Ainsworth Pond is located below the Hillsburgh Dam, east of Station Street. The pond was surveyed five times between April and June, 2015. One turtle was observed over the course of all surveys, a Midland Painted Turtle. Therefore this pond is not considered SWH for Turtle Overwintering Habitat.

Turtle Habitat 3

Turtle Habitat 3, the Hillsburgh Pond is located above the Hillsburgh Dam and northwest of station Street. The pond was surveyed five times between April and June, 2015. Over the course of surveys a total of 451 turtles of Midland Painted Turtle and Common Snapping Turtle were observed. Common Snapping Turtle was observed during April and May surveys. The Ecoregion 6E SWH criteria (MNR 2015b) states that: any area with at least five overwintering (observed between March and May) Midland Painted Turtles or one Common Snapping Turtle is considered SWH. Turtle Habitat 3 meets the criteria for SWH.

Turtle Habitat 4

Turtle Habitat 4, is a small unnamed pond located south of the Hillsburgh Pond. The pond was surveyed five times between April and June, 2015. A total of 7 turtles were observed in this feature over the course of surveys. As only four Midland Painted Turtle observations were made during the March-Early May window, this pond is not considered SWH.

Turtle Habitat 5

Turtle Habitat 5, the large meadow marsh is located south of the Elora-Cataract Trail in the western portion of the study area. This pond was surveyed five times between April and June, 2015. Only one turtle was observed over the course of all surveys, a Midland Painted Turtle. Therefore this pond is not considered SWH for Turtle Overwintering Habitat.

3.4.5.1 Turtle SAR, Regional and Local Significance

One turtle species observed is considered a species at risk under the ESA; Common Snapping Turtle is listed as special concern provincially (SARO) and federally (SARA).

Common Snapping Turtle inhabit slow-moving waters with soft, muck bottom and dense aquatic vegetation. Ponds, sloughs and shallow bays are all often used as summering and overwintering habitat (COSEWIC 2008). Snapping Turtles were observed during turtle basking surveys at two of the candidate locations, a further observation in ELC Polygon 9 (SWTO 3-5) identified a Snapping Turtle as an incidental observation. All three locations likely provide either overwintering, or summering habitat or both for Common Snapping Turtle and are indicated on Figure 7.

Midland Painted Turtle is ranked S5 (Very Common), and Common Snapping Turtle is ranked S3 (vulnerable) in Ontario (NHIC, 2015).

3.4.6 Winter Wildlife

A total of 18 species were identified during the Winter Wildlife Survey, including nine bird species and nine mammal species. Suspected Red Fox (*Vulpes vulpes*) tracks were identified in a wooded area and crossed a walking trail, these tracks entered the woods and no human footprints were seen in the same location. However, it is possible these tracks were Domestic Dog (*Canis lupus familiaris*) rather than a Red Fox.

Figure 4 illustrates the Winter Wildlife Survey route. All species identified during the survey are listed in Appendix 12.

3.4.6.1 SAR, Regional and Local Significance

A single immature Bald Eagle (*Haliaeetus leucocephalus*) was observed flying over the study area. Bald Eagles are listed as Special Concern under the ESA, and have an S-Rank of S2N, S4B and are a Tier 1 Species of Conservation Concern (CVC 2010b). All other identified species during the winter wildlife survey are considered provincially Secure (S5) or Apparently Secure (S4).

3.4.7 Migratory Birds

The results of the Migratory Bird Survey are presented in Appendix 13. Locations of significant observations are provided on Figure 7. During the migratory bird survey, a total of 19 species were detected, of these, nine species are common winter residents in Ontario. The remaining 10 are likely migrants, and were observed in numbers inconsistent with significant migratory habitat under the SWH Ecoregion 6E criteria. Most of the species observed in the study area, are also considered common, and/or abundant species, and tolerant to human disturbance.

3.4.7.1 Migratory Bird SAR, Regional and Local Significance

None of the species observed are considered federal or provincial species at risk.

One species, Great Egret, observed foraging in the Hillsburgh Pond is ranked S2B (imperiled) in Ontario (NHIC, 2015). Great Egret is considered a rare breeder in Ontario, but has no official status. All other species observed are ranked S4 (apparently secure) or S5 (secure).

Six species ranked Tier 1 or Tier 2 were observed in the study area. Great Egret, described above, is ranked Tier 1. Wood Duck, American Black Duck, American Crow, Golden-crowned Kinglet (*Regulus satrapa*), and Ruby-crowned Kinglet (*Regulus calendula*) are ranked as CVC Species of Conservation Concerns Status Tier 2.

3.4.8 Incidental Wildlife Observations

All Incidental wildlife observations made outside of the above formal field surveys are presented in Appendix 15. All observations were of single individuals unless otherwise stated. Species with conservation designation are described in Table 6, and identified on Figure 7.

Table 6. Incidental Species with Conservation Designation Observations

Common Name	Scientific Name	Taxa	Date -Observation	Significance
Common Snapping Turtle	<i>Chelydra serpentina</i>	Reptile	April 15 - Adult observed sunning on log in T1 (Rudd Pond) May 14 - Observed in meadow marsh stream between T1 and T2 May 28 - Observed on bank of T2 (Ainsworth Pond)	<ul style="list-style-type: none"> Species of Special Concern, provincially and federally CVC Tier 1 S-rank S3
Great Egret	<i>Ardea alba</i>	Bird	May 28 - Observed foraging in Hillsburgh Pond	<ul style="list-style-type: none"> S-Rank S2 CVC Tier 1
Trumpeter Swan	<i>Cygnus buccinator</i>	Bird	April 29 & May 28 - Pair observed foraging in Hillsburgh pond	<ul style="list-style-type: none"> CVC Tier 1

3.4.9 Species Listed under the Endangered Species Act

Observations, habitat requirements, breeding evidence and a habitat assessment of six species at risk, Bald Eagle, Bobolink, Eastern Wood-pewee, Eastern Meadowlark, Little Brown Myotis and Common Snapping Turtle, observed in the study area, are discussed below. No federal or provincially listed plant or fish species were identified within the study area through background research, provided data, or field observations.

3.4.9.1 Bald Eagle

Bald Eagle is listed as special concern provincially (ESA 2007) and federally (Species at Risk Public Registry 2014), general habitat protection is not afforded to special concern species. However, species listed as Special Concern and their habitat is protected under the PPS (2014), through the protection of Significant Wildlife Habitat. This species prefers deciduous and mixed-deciduous forest habitat close to water bodies including lakes and rivers; nests in super canopy trees including Pine (Armstrong 2014). The individual was only observed during the winter wildlife visit, and would not be breeding at that time.

3.4.9.2 Bobolink

Bobolink (*Dolichonyx oryzivorus*) is listed as Threatened provincially (ESA 2007) and federally (Species at Risk Public Registry 2014). Bobolink and their general habitat are afforded protection under the ESA. The species typically nests in open grasslands and hay fields. Bobolink are an area-sensitive species, preferring grassland habitat >10ha in area. The individual was observed incidentally in the forb meadow, adjacent the MEMM3 community; one lone male was flushed from the ground, no singing or signs of agitation or nesting were observed. This species is unlikely to be breeding in the study area. This species is unlikely to be breeding in the study area as a result of low quality, size and availability of preferred habitat.

3.4.9.3 Eastern Wood-pewee

Eastern Wood-pewee is listed as Special Concern provincially (ESA 2007) and federally (Species at Risk Public Registry 2014); general habitat protection is not afforded to Special Concern species under the ESA. However, species listed as Special Concern and their habitat is protected under the PPS (2014), through the protection of Significant Wildlife Habitat. The species typically nests in forest clearings and edges of deciduous and mixed forests with an open understory (MNR 2014b). Nests are built on top of the horizontal limbs of mature

deciduous trees (COSEWIC 2012). Eastern Wood-pewee was observed singing in the Sugar Maple Deciduous Forest, during breeding bird surveys. This species is assumed to be breeding within the deciduous forest of the study area.

3.4.9.4 Eastern Meadowlark

Eastern Meadowlark is listed as threatened provincially (ESA 2007) and federally (COSEWIC 2011a). Eastern Meadowlark and their general habitat are afforded protection under the ESA. The species typically nests in open grasslands and hay fields. Eastern meadowlark are an area-sensitive species, preferring grassland habitat >10ha in area. Eastern meadowlark was observed singing from an adjacent agricultural field, south of the study area. This species is unlikely to be breeding in the study area as a result of low quality, size and availability of preferred habitat.

3.4.9.5 Little Brown Myotis

Little Brown Myotis is listed as Endangered provincially (ESA 2007) and federally (Species at Risk Public Registry 2014). Little Brown Myotis and their general habitat are afforded protection under the ESA. Maternal Roosts of little brown Myotis are usually associated with buildings (attics, barns etc.) and occasionally found in large diameter trees (DBH at least 25-44 cm) (COSEWIC 2013). Little Brown Myotis were observed flying towards the Hillsburgh Pond during attendance at an MNRF survey of a known maternal roost adjacent to the study area (*Figure 7*). Surveys conducted by MNRF (2012-2015) indicate an active maternity colony in a building adjacent to the Hillsburgh EA study area. The MNRF indicated that active foraging likely occurs over the Hillsburgh and Ainsworth ponds. Maternity habitat for this species, in trees, is unlikely to occur in the study area. Further studies may be required to confirm presence or absence of Maternity habitat for this species in the study area.

3.4.9.6 Common Snapping Turtle

Common Snapping Turtle is listed as special concern provincially (ESA 2007) and federally (Species at Risk Public Registry 2014), general habitat protection is not afforded to special concern species. However, species listed as Special Concern and their habitat is protected under the PPS (2014), through the protection of Significant Wildlife Habitat. Snapping Turtle is generally found in shallow waters with soft mud bottoms and leaf litter (COSEWIC 2008a). Nesting occurs on gravelly or sandy areas along streams, roadsides or embankments. Observations of Snapping Turtle were made throughout the study area in ponds, wetlands and creeks. This species is confirmed as overwintering, nesting and breeding in the study area.

3.5 Significant Wildlife Habitat

With guidance from the *Significant Wildlife Habitat Technical Guide* (2000) and the SWH EcoRegion Criterion Schedule 6E (2015), four types of Significant Wildlife Habitat (SWH) were confirmed as present within the study area (*Appendix 16*). Confirmed habitat and its location and assessment are presented in *Table 7*. Studies to confirm Bat Maternity Habitat have not been completed in the study area. Following the MNRF Guelph District Protocol (2014), all Candidate Bat Maternity Habitat will be assumed significant. In the event that site activities will affect significant habitat, it is recommended that detailed studies of Candidate Bat Habitat which may be affected, occur pre-construction.

Table 7. Confirmed Significant Wildlife Habitat

SIGNIFICANT WILDLIFE HABITAT TYPE	RATIONALE	LOCATION (FIGURE 7)
Waterfowl Stopover and Staging (Aquatic)	<ul style="list-style-type: none"> Large shallow, open water feature, with abundant aquatic vegetation and soft muck bottom. Surveys completed by CVC did not meet the criteria outlined in the SWH guide. Aggregate of 100 or more listed bird species for 7 days observed, through a combination of field observations and background resources (e-bird 2015). 	SWH 1
Turtle Wintering Area	<ul style="list-style-type: none"> Two large shallow ponds in the study area met the criteria for turtle overwintering; both have muck bottoms and had observations of greater than 5 Midland Painted Turtles or 1 Snapping Turtle during spring surveys. 	SWH 1, SWH 2
Habitat for Special Concern and Rare Wildlife Species	<ul style="list-style-type: none"> Three of the pond features, and one of the stream/meadow marsh features in the study area had observations of Snapping Turtles, either through surveys, or incidentally. 	SWH 1, SWH 2, SWH 3, SWH4
Habitat for Special Concern and Rare Wildlife Species	<ul style="list-style-type: none"> The deciduous woodland feature had probable breeding evidence of Eastern Wood-pewee during breeding bird surveys. 	SWH5
Bat Maternity Habitat	<ul style="list-style-type: none"> All ELC communities meeting the criteria for bat habitat, as listed in the MNRF Guelph District guidelines, including FOD, FOM, FOC, SWD, SWM, SWC with trees >25cm DBH. 	SWH 4, SWH6, SWH7, SWH9, SWH10, SWH11, SWH12
Amphibian Breeding Habitat (Woodland)	<ul style="list-style-type: none"> Four areas identified as candidate habitat in the study area met the criteria for significance. Each feature included at least two of the listed species and greater than 20 individuals. 	SWH1, SWH13, SWH14, SWH15

3.6 SAR Habitat Assessment

An assessment of all Species at Risk, and species with conservation designation, that have the potential to occur in the study area based on lists provided by CVC, MNRF and the NHIC was completed, and is provided in *Appendix 17*. Species assessed include all species with Provincial SARO status, Federal SARA status, or an S-rank of S1-S3. Species assessed with the potential to occur in the study area, but that were not observed during field studies are discussed in detail below.

3.6.1 Vegetation

3.6.1.1 American Chestnut

American Chestnut (*Castanea dentata*) is listed as Endangered provincially (ESA 2007) and federally (SARA 2014). They primarily occur in deciduous forest communities with sandy soil. The species was highly impacted by the Chestnut blight in the early 1900's, which killed 99% of individual trees (MNRF 2015a). The study area is outside of the current known species occurrences (MNRF 2015a).

3.6.1.2 American Ginseng

American Ginseng (*Panax quinquefolius*) is listed as Endangered provincially (ESA 2007) and federally (SARA 2014). The species occurs in rich, moist undisturbed deciduous forests (MNRF 2015a). The FODM5-8 community within the study area provides potential habitat, although historic disturbances and small size of the community may limit habitat potential. American

Ginseng was not identified in the community during field surveys or through previous CVC surveys.

3.6.1.3 *Butternut*

Butternut (*Juglans cinerea*) is listed as Endangered provincially (ESA 2007) and federally (Species at Risk Public Registry 2014). Butternuts primarily occur in rich, moist well drained soils, often along streams (MNR 2015a). Habitat for Butternut is present along the streams throughout the study area, specifically communities SWMO3-3, FODM7-7, SWDM2-1 and FODM8-1. Butternut was not identified in these communities during field surveys or through previous CVC surveys.

3.6.1.4 *Hill's Pondweed*

Hill's Pondweed (*Potamogeton hillii*) is listed as special concern provincially (ESA 2007). The species is found in slow-moving, clear cold stream, ponds, lakes and wetlands. The Ponds within the study area provide potential habitat for this species, although water temperatures are likely to high. Hill's Pondweed is known to occur east of the study area, within the Credit River System (MNR 2015a). Hill's Pondweed was not identified in the Study Area during field surveys or through previous CVC surveys. Detailed aquatic plant surveys of the ponds were not completed as part of this study.

3.6.1.5 *Carey's Sedge*

Carey's Sedge (*Carex careyana*) is listed as S2 in Ontario (NHIC). The species grows in dry to moist rich deciduous upland forests (NatureServe 2015). Deciduous forests of FODM5-8 and FOCM6 provide potential habitat. Carey's Sedge was not identified in the study area during field surveys or through previous CVC surveys.

3.6.1.6 *Rugulose Grapefern*

Rugulose Grapefern (*Sceptridium rugulosum*) is listed as an S2 in Ontario (NHIC). The species grows in sandy to silty soil in open fields, young successional forests or at the edge of forests (Wagner and Wagner 1982). The edges of deciduous forests communities, FODM5-8 and FOCM6 provide potential habitat. Rugulose Grapefern was not identified in the study area during field surveys or through previous CVC surveys.

3.6.2 **Wildlife**

3.6.2.1 *Monarch Butterfly*

Monarch Butterfly (*Danaus plexippus*) is listed as special concern provincially (ESA 2007) and federally (Species at Risk Public Registry 2014). They occur primarily where milkweed and wildflowers exist; including abandoned farmland, along roadsides, and other open spaces (COSEWIC 2010). Habitat for Monarch Butterfly occurs on the forb meadow, south-west of, and including the MEMM3 community. Common Milkweed is abundant in this community, and would provide excellent habitat for Monarch Butterflies. Monarch Butterfly was not observed incidentally during any surveys in the study area, or previously observed during CVC surveys.

3.6.2.2 *West Virginia White*

West Virginia White (*Pieris virginianensis*) is listed as special concern provincially (ESA 2007) and federally (Species at Risk Public Registry 2014). This species generally prefers moist, deciduous woodlands. The larvae feed only on the leaves of a few host plants, including the Two-leaved Toothwort (*Cardamine diphylla*) and Cut-leaved Toothwort (*Cardamine concatenata*) (Burke 2013). Habitat (including host plants) occurs in the study area in small

areas of the SWMCM1-2 and SWMO1-1 communities. West Virginia White was not observed incidentally within the potential habitat communities during spring surveys on the host plant, or during studies completed by CVC.

3.6.2.3 Barn Swallow

Barn Swallow is listed as threatened provincially (ESA 2007) and federally (Species at Risk Public Registry 2014). Barn swallow occurs in farmland, along lake/river shorelines, in wooded clearings and in urban populated areas. Nesting may occur inside or outside buildings; under bridges and in road culverts (COSEWIC 2011b). Habitat for barn swallow is present in the study area, including under bridges, and dam structures. Barn Swallow was not observed during breeding bird surveys, incidentally in the study area, or during studies completed by CVC.

3.6.2.4 Canada Warbler

Canada Warbler is listed as special concern provincially (ESA 2007) and threatened federally (Species at Risk Public Registry, 2014). Canada Warbler prefers wet, coniferous, deciduous and mixed forest types, with a dense shrub layer (COSEWIC 2008b). Habitat for Canada Warbler may occur in the wet mixed forest occurring throughout the study area (SWCM3-2, SWCM1-2). Canada warbler was not observed during breeding bird surveys, or incidentally in the study area, or during studies completed by CVC.

3.6.3 Fish

3.6.3.1 Redside Dace

Redside Dace (*Clinostomus elongatus*) is listed as Endangered provincially (ESA 2007) and, is Under Consideration for listing federally (SARA 2014) and listed as Endangered under COSEWIC (2007). Redside Dace inhabit cool to cold water tributaries, with most Ontario populations occurring in streams flowing to the west basin of Lake Ontario (MNR 2015a). The stream segments within the study area provide adequate habitat for the species. Redside Dace are known to occur within the Credit River (NHIC), but have not been identified in the study area. This may be due to exclusion from up-stream reaches by natural and created barriers to fish passage.

3.7 Aquatic Habitat Assessment

CVC temperature data showed that the two stream reaches, upstream of the Hillsburgh pond, are cold water systems, with maximum summer water temperatures generally below 19° C. The open water communities of the Hillsburgh Pond and Ainsworth Pond and directly downstream from these communities are warm water, with maximum summer temperatures above 25°C. The CVC temperature data showed that the tributary sections directly downstream of the Ainsworth and Rudd Ponds are warm water, but transition to cold water further downstream from the ponds. Due to the spatial scale of the data provided by CVC and uncertainty in the exact location of the sampling points, it was not possible to determine the temperature regime for all segments of the watercourse within the study area. The data provides an overall picture indicating that water tends to be cold within the tributaries upstream of the ponds, increases in temperature to warm water systems within the ponds, and gradually cools back down in the tributaries downstream of the ponds. This is consistent with what would be expected based on ground water fed streams, and the known thermal influence of ponds.

CVC has classified the water courses within the study area into thermal fish communities, such as cold water, cool water and warm water systems based on Fish collection Records (ESSMP

2011 – fig 2.6.1). These classifications are established according to the fish species present and their preferred thermal conditions. This classification characterizes all the tributaries within the study area as cold water fish communities, while the three, online, open water ponds are classified as warm water fish communities. The thermal fish community classification is shown on the Aquatic Habitat Assessment map (*Figure 5*).

Based on the aquatic habitat survey, potential trout (Brook and Brown) spawning habitat was identified in seven stream segments (4, 5, 7, 8, 11, 13 and 14). These areas generally correspond with areas of fish spawning activity, as identified in the Erin Servicing and Settlement Master Plan – Phase 1: Environmental Component (ESSMP 2011). If used as spawning grounds, these areas would be sensitive to sedimentation from erosion or upstream activities.

The aquatic assessment identified three full barriers to fish passage within the study area that would prevent all upstream passage of fish from downstream reaches. These full barriers occur at the outflows of the three main online ponds (Hillsburgh, Ainsworth, and Rudd). Additionally, two partial barriers were identified within the study area. One partial barrier is located at the secondary (North) outfall of the Ainsworth Pond, where sandbags and plastic lining are used to reinforce the outfall and reduce erosion. This partial barrier consists of multiple small drops in elevation, with pools in-between. It may be possible for jumping fish to ascend and pass this barrier during high water conditions. The second partial barrier to fish passage is at the south tributary flowing into the Hillsburgh Pond, connecting segment two and segment three. This partial barrier consists of a presumed fish gate that is likely intended to keep common carp out of the upstream reach. At the same general location, a log jams persists that could make passage difficult for larger and non-jumping fish.

Areas of potential trout spawning habitat, barriers to fish passage, thermal fish community classification and other relevant information are presented on *Figure 5*. *Appendix 18* provides a summary of each watercourse segment assessed.



Image 3. Partial Fish Barrier, upstream reach of segment 3.



Image 4. Full Fish Barrier, upstream reach of segment 5.



Image 5. Partial Fish Barrier, north upstream reach of segment 7.



Image 6. Full Fish Barrier, south upstream reach of segment 7.



Image 7. Full Fish Barrier, upstream reach of segment 11.



Image 8. Full Fish Barrier, upstream reach of segment 11.

3.8 Landscape Evaluation

3.8.1 Ecoregion

The study area is located within Ecoregion 6E. This is the second most densely populated ecoregion in Ontario (MNR 2009), containing a number of large urban centers. The climate of the ecoregion is mild and moist with mean annual precipitation between 759 to 1,087 mm. The underlying geology of the ecoregion is dolomite and limestone, with deep glacially deposited surface soils covering the bedrock in most areas.

Forest cover of the ecoregion is approximately 30.1% and composed of a diverse mixture of hardwood forests, lowlands and flood plain forest. Common tree species within the Ecoregion include; Sugar Maple (*Acer saccharum*), American Beech (*Fagus grandifolia*), White Ash (*Fraxinus americana*), Eastern Hemlock (*Tsuga canadensis*), Green Ash (*Fraxinus pennsylvanica*), Silver Maple (*Acer saccharinum*), Red Maple (*Acer rubrum*), Eastern White Cedar (*Thuja occidentalis*), Yellow Birch (*Betula alleghaniensis*), Balsam Fir (*Abies balsamea*), and Black Ash (*Fraxinus nigra*) (MNR 2009).

3.8.2 Surficial Geology and Groundwater

The surface geology of the study area and surrounding landscape are part of the Orangeville Moraine and contain; silt to clay till, silty sand to sandy silt, ice-contact stratified drift, alluvium, glacio-fluvial outwash gravel, glacio-fluvial outwash sand, and organics (Cowan 1976 as cited in ESSMP 2011). The area generally has a high ground water recharge rate, which supplies much of the base flow of the West Credit River through ground water discharge (ESSMP 2011). This discharge of cold ground water into the river systems contributes to the cold water system of the area.

3.8.3 Connectivity and Existing Natural Features

Natural features of the study area such as; Significant Woodland, the West Credit River and the Provincially Significant West Credit River Wetland Complex serve as linkage corridors within the broader landscape. The Hillsburgh Pond and surrounding wetland are part of the Provincially Significant West Credit River Wetland Complex and are connected to the up-stream Alton-Hillsburgh Provincially Significant Wetland Complex by the West Credit River. The study area also provides a direct corridor between large forest complexes to the southeast and the forests to the north and west of the study area. Many of these forests and natural communities are within or adjacent to watercourses of the West Credit River watershed (*Figure 7*).

3.8.4 Aquatic Habitat

Within the study area and downstream of the study area, there are multiple barriers to fish movement. These have been created through the impoundment of water for the creation of ponds (ESSMP 2011). These barriers limit or stop upstream migration of fish species and prevent access to spawning areas. Upstream of the study area, the ESSMP identifies no barriers to fish passage, and air photo interpretation identified two potential barriers to fish passage.

3.8.5 Rare Features

The open water aquatic community of the Hillsburgh Pond is considered to be rare in the landscape, with only 2.8% of the subwatershed consisting of open water aquatic communities (ESSMP 2011). The Hillsburgh Pond also provides waterfowl stopover and staging habitat, which is considered rare in the larger landscape. The Treed Fen Community within the study area (FETC1-2) is considered a rare community within the landscape, with only 0.3% coverage of fens identified in the ESSMP (2011). The fen community within the study area was not previously identified as a fen community by CVC.

DRAFT

4.0 Summary of Natural Heritage Constraints

The following is a summary of the existing natural heritage conditions assessed and identified within the study area of the Hillsburgh Dam Environmental Assessment. A summary of significant features is provided in *Table 8*.

4.1 Summary of Existing Conditions

4.1.1 Vegetation

Within the study area, three season ELC and three season botanical inventories were completed, where property access was permitted.

1. 31 natural or naturalized vegetation communities were identified, characterised and mapped. None of the ELC communities are considered provincially rare. The fen community (FETC1-2) and open water communities (SAS_1, SAM_1-8, and OAW) are considered rare in the Town of Erin (ESSMP 2011).
2. 394 species or distinct sub-species of plants were identified within the study area through field inventory and background sources. 72% of identified species are native to Ontario, with the remaining 28% of identified species exotic to Ontario.
3. No provincial or federal Species at Risk were identified within the study area. One species identified through background resources, A moss species (*Fontinalis sullivantii*), is ranked as an S1 (Extremely rare in Ontario). 10 identified species are considered significant in Wellington County, and 77 species are classified as CVC Species of Conservation Concern Status Tier 2 on the CVC's ranking system.

4.1.2 Wetlands

1. The Provincially Significant West Credit Wetland Complex is a core natural feature within the study area and surrounding landscape.
2. The wetland was evaluated under the Ontario Wetland Evaluation System (OWES) by the MNRF in 1995 and updated in 2005.
3. The mapped wetland boundary was field verified using the OWES 2013, and found to be largely accurate within the study area, with only a few minor deviations from the mapped boundary provided by LIO.
4. Within the study area the wetland consists of Coniferous Swamp, Mixed Swamp, Deciduous Swamp, Thicket Swamp, Treed Fen, Meadow Marsh, Shallow Marsh, Submerged Shallow Aquatic, Mixed Shallow Aquatic, and Open Aquatic communities within the study area.
5. The OWES Scoring Record identified the wetland as an important feature in reducing the risk of flooding and erosion; which are identified by CVC as important criteria when considering preferred EA options.

4.1.3 Wildlife

1. Where access was permitted, surveys for Amphibians, Breeding Birds, Snakes, Turtles, Winter Wildlife, and Bird Migration Monitoring were completed in appropriate habitats in the study area.
 - a. Three rounds of calling amphibian surveys occurred at seven candidate locations during the months of April, May and June 2015.
 - b. Breeding Bird Surveys were conducted twice, once in June and once in early July; a total of 10 point counts and four Marsh Bird playback surveys were completed.
 - c. Three visual and hand search surveys for snakes were completed in candidate areas between April and early June.
 - d. No breeding habitat for Salamanders was identified in the study area; as a result, further surveys were not required.
 - e. Turtle surveys were conducted between April and June, for a total of five rounds, five candidate habitats were investigated.
 - f. Two migratory bird surveys were completed, one targeted shorebirds and one targeted songbirds, occurring in August and October, respectively.
2. Six frog, 70 bird, one snake, two turtle and 11 mammal species were observed in the study area over the course of all field investigations.
3. Five species listed under the ESA were identified during field investigations: Eastern Meadowlark (THR), Bobolink (THR), Eastern Wood-pewee (SC), Bald Eagle (SC) and Common Snapping Turtle (SC).
 - a. Eastern meadowlark was observed outside the study area on adjacent lands to the south. One individual was heard signing during one round of breeding bird surveys.
 - b. Bobolink was observed incidentally and showed no signs of breeding; one lone male was flushed from the forb meadow, east of the Elora-Cataract Trail way, ELC Polygon 12 (MEMM3).
 - c. Eastern Wood-pewee had probable breeding in the study area, occurring in the deciduous forest community in the eastern portion of the study area, ELC Polygon 4 (FODM5-8).
 - d. Bald Eagle was observed during a winter wildlife survey, soaring over the West Credit River in the study area.
 - e. Common Snapping Turtle was confirmed as overwintering in the Hillsburgh and Rudd Ponds, one nest was identified along the berm of the Rudd Pond.
4. One species identified, Trumpeter Swan, is ranked as a CVC Species of Conservation Concern Status Tier 1. Trumpeter Swans were identified on the Hillsburgh Pond during surveys conducted during the spring migration season.
5. One species identified is considered rare in Ontario; Great Egret and is ranked S2B. No evidence of breeding occurred in the study area. Individuals were seen during spring and fall migration.
6. Thirty-nine species of wildlife identified in the study area are considered significant in Wellington County.

4.1.4 Significant Wildlife Habitat (SWH)

1. A review of the study area using a combination of methods presented in the Ecoregion criteria guide, air photo interpretation and field investigations assessed the study area for Significant Wildlife Habitat that may occur in ecoregion 6E.
2. A total of six types of SWH were identified in the study area, and confirmed or assumed significant using the results of all surveys completed in the study area and background resources.
3. Waterfowl Stopover and Staging (Aquatic), Turtle Wintering Area, Habitat for Special Concern and Rare Wildlife Species, and Amphibian Breeding Habitat (Woodland) were identified as candidate, confirmed and delineated in the study area.
4. Candidate Bat Maternity Habitat was identified in the study area, and assumed significant. Further surveys are proposed pre-construction where impacts to candidate habitat may occur.

4.1.5 Species with Conservation Designation Habitat Assessment

1. A review of the study area was completed, using habitat requirements from reference documents, air photo interpretation and field investigations, to assess for habitat that may be suitable for Species with conservation designation (listed under the ESA or an S-rank of S1-S3). This list included all species identified through background review as occurring in Wellington County (MNR 2015c), identified by CVC (2008-2014), or identified through NHIC (2015) that may occur in the study area.
2. Potential habitat for 16 species was identified in the study area. Surveys conducted targeted habitat that may be suitable for these species through the completion of breeding bird surveys, vegetation surveys, snake surveys, and turtle surveys.
3. During all surveys completed in the study area, seven of the wildlife species with candidate habitat were identified as occurring in the study area. Of those, four (Eastern Meadowlark, Eastern Wood-pewee, Common Snapping Turtle and Little Brown Myotis) were completing important life stages in the vicinity of the study area, and three were observed incidentally, outside the breeding season, or showing no signs of breeding evidence (Bobolink, Great Egret and Bald Eagle).

4.1.6 Aquatic Habitat Assessment

1. An Aquatic Habitat Assessment was completed for all open water communities and stream reaches within the study area. The Aquatic Habitat Assessment was based on a field survey completed on October 19th, 2015 and background resources of CVC temperature data, fish community classification, fish species records and the Credit River Fishery Management Plan.
2. Surveys identified areas of potential trout spawning habitat throughout the cold water watercourses of study area, as well as immediately upstream and downstream of the study area.
3. Three full and two partial barriers to fish passage were identified within the study area. These barriers reduce or prevent passage of fish to adjacent habitats.

4. The West Credit River is a natural, cold water system fed primarily by ground water, the thermal influences of the three online ponds within the study area increases the temperature of the watercourse.
5. Cold water fish communities are generally found within the tributary sections, while the online ponds contain primarily warm water fish communities.
6. The invasive Round Goby has been identified within the Upper West Credit River system, including up-stream and downstream of the study area.

DRAFT

4.2 Summary of Significant Features

A summary of existing conditions of natural heritage are provided in Section 4.1. Several existing condition features are significant, including but not limited to, Species at Risk under Ontario’s Endangered Species Act and Significant Wildlife Habitat under the Provincial Policy Statement. In addition to the natural heritage present across the study area, significant features are given elevated levels of protection and management. A summary of significant features is provided in *Table 8*.

Significance /Type	Site Assessment and Observations	Legislation, Policy and Management Considerations	Figure
Species at Risk	<ul style="list-style-type: none"> •Bald Eagle (SC), observed in the study area, no habitat or breeding evidence. •Bobolink (THR), observed in the study area, no habitat or breeding evidence. •Eastern Meadowlark (THR), Breeding Evidence outside study area. •Eastern Wood-pewee (SC), Breeding evidence in the study area. •Little Brown Myotis (END), observed outside study area, Hillsburgh pond may provide important foraging habitat •Common Snapping Turtle (SC), overwintering, nesting and breeding Habitat in the study area. 	<p><i>Endangered Species Act, 2007</i></p> <ul style="list-style-type: none"> •Species listed as Special Concern (SC) are not afforded general habitat protection under the ESA. •Threatened (THR) and Endangered (END) species are afforded General Habitat Protection under the ESA. <p><i>Provincial Policy Statement, 2014</i></p> <ul style="list-style-type: none"> •The habitat of Species listed as Special concern is protected under the PPS as Significant Wildlife Habitat. 	7
Fish Habitat	<ul style="list-style-type: none"> • All watercourse and open water communities provide fish habitat, with known fish communities. •Within the study area 3 full and 2 partial barriers to fish passage exist that may prevent/restrict fish species from reaching appropriate spawning grounds. •All watercourses and bodies of water within the study area are managed as Coldwater fisheries, with a specific focus of Brook Trout and Resident Brown Trout. 	<p><i>Fisheries Act, 2013</i></p> <ul style="list-style-type: none"> •Protects the productivity of recreational, commercial and Aboriginal fisheries. Fish communities and habitat within the study area are afforded protection. <p><i>Credit River Fisheries management Plan.</i></p> <ul style="list-style-type: none"> •Barriers to fish passage are recommended for removal or mitigation within the Credit River Fisheries Management Plan. • Construction must respect the Coldwater fisheries timing window of no in-water work from October 1 – June 30. 	5

Table 8. Summary of Significant Features

Significance /Type	Site Assessment and Observations	Legislation, Policy and Management Considerations	Figure
Significant Wildlife Habitat (SWH)	<ul style="list-style-type: none"> •Waterfowl Stopover and Staging (Aquatic). •Turtle Wintering Area. •Habitat for Special Concern and Rare Wildlife Species. •Bat Maternity Habitat. •Amphibian Breeding Habitat (Woodland). 	<p><i>Provincial Policy Statement, 2014</i></p> <ul style="list-style-type: none"> •Under the PPS, development and site alteration are not permitted in significant wildlife habitat unless it has been demonstrated that there will be no negative impacts on the natural features or their ecological functions. 	7
Rare Species Habitat	<ul style="list-style-type: none"> •Great Egret - S2, CVC Tier 1 •Trumpeter Swan - CVC Tier 1 	<p><i>Provincial Rarity Rank (S-Rank)</i></p> <ul style="list-style-type: none"> •An S-Rank of S2 indicates that the species is considered imperiled in the province, with few known populations. <p><i>CVC's Species of Conservation Concern project (2010b)</i></p> <ul style="list-style-type: none"> •Tier 1 species are those with low abundance, low population density, specialized habitat requirements, and/or a narrow tolerance to disturbance. CVC Tier 1 species should be identified and managed in order to avoid changes to habitat or site alteration. 	7
Rare or Important Landscape Features	<ul style="list-style-type: none"> • The natural lands within the study area create continuous corridors with surrounding natural features such as Provincially Significant Wetlands, fish spawning habitat and Significant Woodlands outside the study area. • The open water communities of the Hillsburgh, Ainsworth and Rudd Ponds are rare communities in the landscape (ESSMP 2011). • The Treed Fen Community is a rare community within the landscape (ESSMP 2011). 	<p><i>Wellington County Official Plan</i></p> <ul style="list-style-type: none"> •Terrestrial, Wetland, Riparian and Aquatic connecting corridors considered "Protection Area 1" and are included in Wellington Counties Greenlands System designation. Activities that diminish or degrade the essential function of Greenlands Systems will be prohibited. <p><i>Erin Service and Settlement Master Plan (2011)</i></p> <ul style="list-style-type: none"> • Rare communities should be considered a high priority for protection in the landscape (ESSMP 2011). 	6

Table 8. Summary of Significant Features

Significance /Type	Site Assessment and Observations	Legislation, Policy and Management Considerations	Figure
Provincially Significant Wetlands	<ul style="list-style-type: none"> The West Credit Wetland Complex comprises approximately 44.6 ha of the study area. 	<p><i>Credit Valley Conservation</i></p> <ul style="list-style-type: none"> Interference or alteration to wetlands or watercourses are generally not permitted <p><i>CVC's Policies are regulated under the Administration of the Development, Interference with Wetlands and Alterations to Shorelines and Watercourses Regulation (Ontario Regulation 160/06).</i></p> <ul style="list-style-type: none"> CVC and MNR may permit development or site alteration where impacts have been addressed through an environmental assessment. 	6

DRAFT

5.0 Assessment of Alternatives, and Impacts to the Natural Environment

The Assessment of Alternatives was completed by evaluating the potential and actual impacts of each proposed alternative on the identified Significant Natural Heritage Features in the study area and surrounding landscape. The current state of the dam and natural environment were considered to be neutral, and positive or negative impacts were assessed relative to this condition.

5.1 Alternatives Considered for the Hillsburgh Dam and Pond

Four alternatives (A, B, C and D), two of which (C and D) have two options, were considered in the evaluation of impacts to the Natural Environment. The following alternatives and summary descriptions were provided by Triton Engineering.

1) **Alternative A - Do Nothing**

This is the neutral “null” alternative, against which all other alternatives will be measured. If nothing is done to repair and/or replace the dam and bridge, the dam and bridge will continue to deteriorate and eventually fail.

2) **Alternative B - Rehabilitate Hillsburgh Dam and Reconstruct the Bridge**

Construct a new bridge at the same location or a new location along the dam. Alternative B consists of a larger bridge to contain the Regional Storm event without overtopping the road to comply with the Lakes and Rivers Improvement Act. The dam will be rehabilitated to meet the Ministry of Natural Resources and Forestry current dam safety standards.

3) **Alternative C - Rehabilitate Station Street Bridge**

Option 1: Rehabilitate Station Street Bridge, Decommission Dam. This alternative consists of doing only the work necessary to bring the bridge to meet current safety and construction requirements. The dam will be decommissioned, which will alter the pond to a watercourse.

Option 2: Rehabilitate Station Street Bridge, Decommission Dam, and Construct an Offline Pond. This alternative consists of doing only the work necessary to bring the bridge to meet current safety and construction requirements. The dam will be decommissioned, which will alter the pond to a watercourse. An off-line pond will be constructed inside the footprint of the existing Hillsburgh Pond.

4) **Alternative D - Reconstruct Station Street Bridge**

Option 1: Reconstruct Station Street Bridge, Decommission Dam. Construct a new bridge at the same location or a new location along the dam. This alternative consists of decommissioning the dam, which will alter the pond to a watercourse. The bridge will be constructed under the MTO Highway Drainage Design Standards.

Option 2: Reconstruct Station Street Bridge, Decommission Dam, and Construct an Offline Pond. Construct a new bridge at the same location or a new location along the dam. This alternative consists of decommissioning the dam, which will alter the pond

to a watercourse. The bridge will be constructed under the MTO Highway Drainage Design Standards. An off-line pond will be constructed inside the footprint of the existing Hillsburgh Pond.

5.2 Evaluation of Impacts

In order to evaluate the EA Alternatives provided, each alternative was assessed with respect to potential impacts to the significant features of the natural environment identified in the study area (Table 9). These are:

- Impacts to Habitat of Species at Risk;
- Impacts to Fish Habitat, (including thermal regime and fish passage);
- Impacts to Significant Wildlife Habitat;
- Impacts to Rare Species Habitat;
- Impacts to Rare or Important Landscape Features, and;
- Impacts to Provincially Significant Wetlands.

DRAFT

Table 9: Comparison of Alternatives

CRITERIA	ALTERNATIVE A "Do Nothing"	ALTERNATIVE B Rehabilitate Hillsburgh Dam and Reconstruct Station Street Bridge	ALTERNATIVE C Rehabilitate Station Street Bridge and Option 1 or Option 2		ALTERNATIVE D Reconstruct Station Street Bridge and Option 1 or Option 2	
			OPTION 1 Decommission Dam	OPTION 2 Construct Offline Pond	OPTION 1 Decommission Dam	OPTION 2 Construct Offline Pond
NATURAL ENVIRONMENT						
Species at Risk (SAR)	No impacts are anticipated under current state. Uncontrolled dam failure could cause significant negative impacts to Species at Risk.	Yes- Minor impacts to SAR Habitat are expected during construction. If appropriate mitigation measures are put in place, no long term impacts are anticipated following construction and restoration.	Yes-Impacts to SAR Habitat are expected during construction; and long term impacts to the habitat through permanent removal of overwintering habitat for Common Snapping Turtle, and permanent removal of foraging habitat for Little Brown Myotis.	Yes- Minor impacts to SAR Habitat are expected during construction. If appropriate design and mitigation measures are put in place, no long term impacts are anticipated following construction and restoration.	Yes-Impacts to SAR Habitat are expected during construction; and long term impacts to the habitat through permanent removal of overwintering habitat for Common Snapping Turtle, and permanent removal of foraging habitat for Little Brown Myotis.	Yes- Minor impacts to SAR Habitat are expected during construction. If appropriate design and mitigation measures are put in place, no long term impacts are anticipated following construction and restoration.
Fish Habitat	On-going negative impacts to the thermal regime of the watercourse and barrier to fish movement. Uncontrolled dam failure could cause significant negative impacts to Fish and Fish Habitat	Yes- Impacts to fish and fish habitat are expected during construction. If appropriate mitigation measures are put in place, no long term impacts are anticipated following construction and restoration. On-going negative impacts to the thermal regime of the watercourse and barrier to fish movement.	Yes- Positive impacts to the managed Cold Water Fishery are anticipated from removing the dam and re-establishing the watercourse. Warm water fish species, which are not managed, would be negatively impacted by loss of habitat.	Yes- Positive impacts to the managed Cold Water Fishery are anticipated from removing the dam. The off-line pond may negatively impact the thermal regime if warm water is allowed to enter the watercourse.	Yes- Positive impacts to the managed Cold Water Fishery are anticipated from removing the dam and re-establishing the watercourse. Warm water fish species, which are not managed, would be negatively impacted by loss of habitat.	Yes- Positive impacts to the managed Cold Water Fishery are anticipated from removing the dam. The off-line pond may negatively impact the thermal regime if warm water is allowed to enter the watercourse.
Significant Wildlife Habitat (SWH)	No impacts are anticipated under current state. Uncontrolled dam failure could cause significant negative impacts to SWH.	Yes- Minor impacts to SWH are expected during construction. If appropriate mitigation measures are put in place, no long term impacts are anticipated following construction and restoration.	Yes-Impacts are expected during construction and long term negative impacts on the features and functions of the following SWH: Waterfowl Stopover and Staging, Turtle overwintering, and Habitat for Special Concern Species and Rare Wildlife Species.	Yes- Minor impacts to SWH are expected during construction. If appropriate mitigation measures are put in place, no long term impacts are anticipated following construction and restoration.	Yes-Impacts are expected during construction and long term negative impacts on the features and functions of the following SWH: Waterfowl Stopover and Staging, Turtle overwintering, and Habitat for Special Concern Species and Rare Wildlife Species.	Yes- Minor impacts to Significant Wildlife Habitat are expected during construction. If appropriate mitigation measures are put in place, no long term impacts are anticipated following construction and restoration.
Rare Species Habitat	No impacts are anticipated under current state. Uncontrolled dam failure could cause significant negative impacts to Rare Species.	Yes- Minor impacts to Rare Species habitat are expected during construction. If appropriate mitigation measures are put in place, no long term impacts are anticipated following construction and restoration.	Yes-Impacts to Rare species are expected during construction, and long term impacts include permanent changes to potential foraging/stopover habitat for Great Egret and Trumpeter Swan.	Yes- Minor impacts to Rare Species habitat are expected during construction. If appropriate mitigation measures are put in place, no long term impacts are anticipated following construction and restoration.	Yes-Impacts to Rare species are expected during construction, and long term impacts include permanent changes to potential foraging/stopover habitat for Great Egret and Trumpeter Swan.	Yes- Minor impacts to Rare Species habitat are expected during construction. If appropriate mitigation measures are put in place, no long term impacts are anticipated following construction and restoration.
Rare or Important Landscape Features	No impacts are anticipated under current state. Uncontrolled dam failure could cause significant negative impacts to Landscape Features.	No impacts are anticipated.	Yes- Impacts to landscape features are expected through the removal of the Hillsburgh Pond open water community, which is a rare community in the Town of Erin. Possible negative impact to the Treed Fen Community if hydrological changes (e.g. lower water table) are associated with the decommissioning of the dam.	Open water community will be maintained through construction of off-line pond. Possible negative impact to the Treed Fen Community if hydrological changes are associated with the decommissioning of the dam.	Yes- Impacts to landscape features are expected through the removal of the Hillsburgh Pond open water community, which is a rare community in the Town of Erin. Possible negative impact to the Treed Fen Community if hydrological changes (e.g. lower water table) are associated with the decommissioning of the dam.	Open water community will be maintained through construction of off-line pond. Possible negative impact to the Treed Fen Community if hydrological changes are associated with the decommissioning of the dam.
Provincially Significant Wetlands (PSW)	No impacts are anticipated under current state. Uncontrolled dam failure could cause significant negative impacts to the PSW	Potential changes to hydrology could impact the upstream and downstream extent and quality of wetland.	Potential changes to hydrology could impact the upstream and downstream extent and quality of wetland.	Potential changes to hydrology could impact the upstream and downstream extent and quality of wetland.	Potential changes to hydrology could impact the upstream and downstream extent and quality of wetland.	Potential changes to hydrology could impact the upstream and downstream extent and quality of wetland.

Legend	Positive	Neutral	Negative (Short-term, minor or potential)	Negative (Long-term, extensive or definitive)
--------	----------	---------	---	---

5.3 Summary of Impacts to the Natural Environment

5.3.1 Alternative A - Do Nothing

The Do Nothing Alternative will result in no immediate additional negative impacts to the existing natural features under the current dam configuration. However, the presence of the dam is known to be causing negative impacts to the natural environment, including altering the watercourse and reducing the quality of fish habitat.

The Do Nothing Alternative will likely cause significant and unknown impacts to the natural environment, in the case of an uncontrolled dam failure.

5.3.2 Alternative B - Rehabilitate Hillsburgh Dam and Reconstruct Station Street Bridge

Rehabilitation of the Station Street Dam and reconstruction of the bridge is not anticipated to result in new long-term negative impacts to the natural Environment, relative to the current state. This alternative will continue to negatively impact the watercourse.

Short-term impacts to the natural environment are expected during construction and rehabilitation of the dam and bridge:

Overwintering Turtles – Draining of the Hillsburgh pond for construction when turtles are overwintering may cause stress or death to turtles, including the Common Snapping Turtle, a species of Special Concern.

Foraging habitat for Little Brown Myotis (SAR) – Draining of the pond for construction, during the maternal season for Little Brown Myotis could reduce feeding opportunities for the colony adjacent the pond during a critical life stage.

Amphibian Breeding Significant Wildlife Habitat – Draining the pond during the amphibian breeding season will lower water levels and reduce the success rate for breeding, and survival of eggs and tadpoles, in wetland areas adjacent the pond.

Waterfowl Stopover and Staging Significant Wildlife Habitat – Draining the pond during the stopover and staging season will affect species of waterfowl reliant on the pond to provide stopover feeding and roosting opportunities during migration.

Fish Habitat – Without mitigation through construction design, rebuilding the dam in a similar configuration to the current state will maintain the negative impacts to fish in the watercourse, through a continued barrier to fish passage, and negative thermal influences.

Provincially Significant Wetland – Changes to hydrology may impact the upstream and downstream extent and quality of the wetland. Since Alternative B will rehabilitate the dam and rebuild the bridge, the impacts to the wetland are anticipated to be short term.

Landscape Features – Alternative B will retain the open water community of the Hillsburgh pond, which is rare in the Town of Erin. Hydrological changes could also result in impacts to the rare treed fen community downstream of the pond.

In general, the anticipated impacts associated with Alternative B are short term, while construction is ongoing. Mitigation measures can be applied to reduce or eliminate long-term negative impacts to the habitats and minimize any short-term impacts.

5.3.3 Alternative C - Rehabilitate Station Street Bridge and Decommission the Hillsburgh Dam

Option 1 – Without an Off-line Pond

Rehabilitation of the Station Street Bridge and decommissioning the Dam will result in both negative and positive impacts to the natural environment, relative to the current state. This alternative will positively impact the watercourse by returning the system to a naturalized stream environment.

Short-term impacts to the natural environment associated with construction are expected during rehabilitation of the bridge. Removal of the dam will cause long-term changes and impacts to the natural environment:

Overwintering Turtles – Draining of the Hillsburgh pond when turtles are overwintering may cause stress or death to turtles, including the Common Snapping Turtle, a species of Special Concern and Midland Painted Turtle. The loss of the Hillsburgh pond to a naturalized watercourse will permanently remove this Significant Wildlife Habitat and cause negative impacts to the resident turtle populations.

Foraging habitat for Little Brown Myotis (SAR) – Draining of the Hillsburgh pond may reduce or eliminate feeding opportunities for the adjacent Little Brown Myotis colony during critical life stages, reducing overall success rates of the colony in the long-term.

Amphibian Breeding Significant Wildlife Habitat – Draining the Hillsburgh pond will reduce the success rate for breeding amphibians in the short and possibly long term, with the potential to affect the hydrology of the wetlands both upstream and downstream of the pond, and reducing or eliminating a number of Significant Amphibian Breeding Areas.

Waterfowl Stopover and Staging Significant Wildlife Habitat – Draining the Hillsburgh pond will affect species of waterfowl reliant on the pond to provide stopover feeding and roosting opportunities during migration. It is considered a rare landscape feature in the area, and provides an important function for the successful migration of waterfowl species.

Rare Species Habitat - Draining of the Hillsburgh pond may reduce feeding and staging opportunities for Great Egret and Trumpeter Swan.

Fish Habitat – Removal of the Hillsburgh pond and establishment of a naturalized watercourse will have positive impacts on the managed cold water fish species, including Brook and Brown Trout, by removing a barrier to fish passage and reduced thermal impacts to the watercourse. Warm water fish species, which are not managed, will be negatively impacted through the loss of habitat with the removal of the Hillsburgh Pond. However, Alternative C – Option 1 provides an overall positive benefit for the more desirable, managed cold water fish species.

Provincially Significant Wetland - Changes to hydrology could impact the upstream and downstream extent and quality of the wetland. Since Alternative C - Option 1 will remove the dam, it is likely this will result in a lower water table and reduce the upstream extent and quality of the Provincially Significant Wetland; detailed hydrological changes are unknown at this time.

Landscape Features – Alternative C - Option 1 will result in the loss of an open water community, which is described as rare in the Town of Erin Service and Settlement Master Plan, through the draining of the Hillsburgh Pond and reestablishment of a watercourse. Hydrological changes could result in impacts to the rare treed fen community downstream of the pond.

Option 2 – With an Off-line Pond

The rehabilitation of the Station Street Bridge, decommissioning the Dam and re-constructing a new offline pond will result in possible short-term impacts to the natural environment during construction and rehabilitation of the dam and bridge. Minor long-term impacts and changes in the natural features are expected. However, the ecological function of Significant Features will be preserved. There are potential impacts to the natural environment:

Overwintering Turtles – Draining of the Hillsburgh pond when turtles are overwintering may cause stress or death to turtles, including the Common Snapping Turtles, a species of Special Concern and Midland Painted Turtle. The loss of the Hillsburgh pond and return to a naturalized watercourse will permanently affect this Significant Wildlife Habitat and cause negative impacts to the resident turtle populations. Through a created offline pond, the Overwintering Turtle Significant Wildlife Habitat would be maintained, resulting in no long term impacts to turtle populations.

Foraging habitat for Little Brown Myotis (SAR) – Drainage of the pond for construction, during the maternal season for Little Brown Myotis may reduce feeding opportunities for the colony adjacent to the pond during a critical life stage, in the short-term. It is expected that the offline pond would continue to provide foraging habitat.

Amphibian Breeding Significant Wildlife Habitat – Draining the pond during the amphibian breeding season would reduce the success rate for breeding, and affect the success of eggs and tadpoles in wetland areas adjacent the pond that may be affected by low water levels, in the short term. It is expected that the offline pond, would continue to provide amphibian breeding SWH.

Waterfowl Stopover and Staging Significant Wildlife Habitat – Draining the pond during the stopover and staging season may affect species of waterfowl reliant on the pond to provide stopover feeding and roosting opportunities during migration, in the short term. It is expected that the offline pond, would continue to provide Waterfowl Stopover and Staging SWH.

Fish Habitat – Removal of the Hillsburgh pond and establishment of a naturalized watercourse will have positive impacts of the managed cold water fish species, including Brook and Brown Trout by removing a barrier to fish passage and reduced thermal impacts to the watercourse. Depending on the design, resident warm water fish species, which are not managed for, will be able to persist within the offline pond.

Provincially Significant Wetland – Changes to hydrology could impact the upstream and downstream extent and quality of the wetland. Since Alternative C - Option 2 will completely remove the dam, it is likely this could result in a lower water table and could reduce the upstream extent of the Provincially Significant Wetland; detailed hydrological changes are unknown at this time.

Landscape Features – Alternative C - Option 2 will retain an open water community, which is rare in the Town of Erin, through the creation of an offline pond within the footprint of the current Hillsburgh Pond. Hydrological changes could result in impacts to the rare treed fen community downstream of the pond.

5.3.4 Alternative D - Reconstruct Station Street Bridge, Decommission Dam

Option 1 – Without an Offline Pond

Reconstruction of the Station Street Bridge and decommission the Dam will result in both negative and positive impacts to the natural environment relative to the current state. Short-term impacts to the natural environment are possible during construction and rehabilitation of the bridge; and the removal of the dam will cause long-term impacts to the natural environment:

Overwintering Turtles – Draining of the Hillsburgh pond when turtles are overwintering may cause stress or death to turtles, including the Common Snapping Turtles, a species of Special Concern and Midland Painted Turtles. The loss of the Hillsburgh pond to a naturalized watercourse will permanently remove this Significant Wildlife Habitat and cause negative impacts to the resident turtle populations.

Foraging habitat for Little Brown Myotis (SAR) – Draining of the Hillsburgh pond may reduce or eliminate feeding opportunities for the adjacent Little Brown Myotis colony during critical life stages, reducing overall success rates of the colony in the long term.

Amphibian Breeding Significant Wildlife Habitat – Draining the Hillsburgh pond is likely to reduce the success rate for breeding amphibians in the short and possibly long term, with the potential to affect the hydrology of the wetlands both upstream and downstream of the pond, reducing or eliminating a number of Significant Amphibian Breeding Areas.

Waterfowl Stopover and Staging Significant Wildlife Habitat – Draining the Hillsburgh pond is likely to affect species of waterfowl reliant on the pond to provide stopover feeding and roosting opportunities during migration. It is considered a rare landscape feature in the area, and provides an important function for the successful migration of waterfowl species.

Rare Species Habitat - Draining of the Hillsburgh pond may reduce feeding and staging opportunities for Great Egret and Trumpeter Swan.

Fish Habitat – Removal of the Hillsburgh pond and establishment of a naturalized watercourse will have positive impacts to the managed cold water fish species, including Brook and Brown Trout, by removing a barrier to fish passage and reduced thermal impacts to the watercourse. Warm water fish species, which are not managed for, will be negatively impacted through the loss of habitat with the removal of the Hillsburgh Pond.

However, Alternative D – Option 1 provides an overall positive benefit for the more desirable, managed cold water fish species.

Provincially Significant Wetland - Changes to hydrology could impact the upstream and downstream extent and quality of the wetland. Since Alternative D - Option 1 will completely remove the dam, it is likely this will result in a lower water table and could reduce the upstream extent and quality of the Provincially Significant Wetland, although detailed hydrological changes are unknown at this time.

Landscape Features – Alternative D - Option 1 will result in the loss of an open water community, which is rare in the Town of Erin, through the draining of the Hillsburgh Pond and reestablishment of a watercourse. Hydrological changes could result in impacts to the rare treed fen community downstream of the pond.

Option 2 – With an Offline Pond

The re-construction of the Station Street Bridge, decommissioning the Dam and construction of a new offline pond, will result in both long-term negative and positive impacts to the natural environment relative to the current state. This alternative will positively impact the watercourse through establishing a naturalized river system. Expected impacts to the natural environment include:

Overwintering Turtles – Draining of the Hillsburgh pond when turtles are overwintering may cause stress or death to turtles, including the Common Snapping Turtles, a species of Special Concern and Midland Painted Turtles. The loss of the Hillsburgh pond and return to a naturalized watercourse will permanently affect this Significant Wildlife Habitat and cause negative impacts to the resident turtle populations. Through the creation of an off-line pond, the Overwintering Turtle Significant Wildlife Habitat could be maintained, resulting in no long term impacts to turtle populations.

Foraging habitat for Little Brown Myotis (SAR) – Drainage of the pond for construction, during the maternal season for Little Brown Myotis may reduce feeding opportunities for the colony adjacent to the pond during a critical life stage, in the short-term. It is expected that the offline pond would continue to provide foraging habitat.

Amphibian Breeding Significant Wildlife Habitat – Draining the pond during the amphibian breeding season may reduce the success rate for breeding, and affect the success of eggs and tadpoles in wetland areas adjacent the pond that may be affected by low water levels, in the short term. It is expected that the offline pond, would continue to provide amphibian breeding SWH.

Waterfowl Stopover and Staging Significant Wildlife Habitat – Draining the pond during the stopover and staging season will affect species of waterfowl reliant on the pond to provide stopover feeding and roosting opportunities during migration, in the short term. It is expected that the offline pond will be of similar size and composition of existing, and would continue to provide Waterfowl Stopover and Staging SWH.

Fish Habitat – Removal of the Hillsburgh pond and establishment of a naturalized watercourse will have positive impacts of the managed cold water fish species, including Brook and Brown Trout, by removing a barrier to fish passage and reduced thermal

impacts to the watercourse. Depending on the design, resident warm water fish species, which are not managed for, will be able to persist within the offline pond.

Provincially Significant Wetland – Changes to hydrology could impact the upstream and downstream extent and quality of the wetland. Since Alternative D - Option 2 will completely remove the dam, it is anticipated that this may result in a lower water table and could reduce the upstream extent of the Provincially Significant Wetland, although detailed hydrological changes are unknown at this time.

Landscape Features – Alternative D - Option 2 will retain an open water community, which is rare in the Town of Erin, through the creation of an off-line pond within the footprint of the current Hillsburgh Pond. Hydrological changes could also result in impacts to the rare treed fen community downstream of the pond.

DRAFT

6.0 Conclusion

The Natural Environment Report was completed as part of the Schedule B Municipal Class Environmental Assessment. The EA is being completed in order to determine the best option to ensure the long term safety of the Hillsburgh Dam, while considering the natural environment, transportation, socio-economic impacts and construction costs. The Natural Environment Report has identified significant species, features and ecological functions within the study area, which were considered while developing and ranking EA options.

The Assessment of Alternatives and Impacts to the Natural Environment identified potential and actual impacts of each proposed EA Alternative with respect to the identified existing natural heritage features in the study area and surrounding landscape. The analysis concluded that either Alternative C - Option 2 or Alternative D - Option 2, which includes the construction of an off-line Pond, are the preferred alternatives from a natural heritage perspective. These alternatives have the least negative impacts to the natural heritage features and provide positive benefits to the Natural Environment in the long term.

Should it be determined that the preferred alternatives listed above are not feasible, Alternative B is the next preferred alternate. If Alternative B is selected, the design should include measures to improve fish habitat long term through the installation of a fish bypass and bottom draw system.

Alternative C - Option 1 and Alternative D - Option 1 are the least preferred alternatives. These options would result in long term negative impacts to the existing natural environment through the complete loss of the open water community of the Hillsburgh Pond. This would have substantial negative impacts to SAR habitat, Significant Wildlife Habitat, Rare Species habitat, and to the overall ecological value of the study area.

Mitigation measures should be developed for any selected Alternative to minimize impacts to protected natural heritage features during construction and retain and enhance the overall ecological integrity of the area.

Prepared By:

ABOUD & ASSOCIATES INC.

Ryan Hamelin, M.Sc.
Terrestrial and Wetland Ecologist

Cheryl-Anne Ross, B.Sc.
Wildlife Ecologist

Reviewed By:

Steven Aboud, B.Sc. Principal.
ISA Certified Arborist No. ON-0323A
ISA Tree Risk Assessment Qualified
Butternut Health Assessor No. 497
Senior Ecologist

DRAFT

5.0 References

- Armstrong, Ted (E.R.). 2014. Management Plan for the Bald Eagle (*Haliaeetus leucocephalus*) in Ontario. Ontario Management Plan Series. Prepared for the Ontario Ministry of Natural Resources and Forestry, Peterborough, Ontario. vii + 53 pp.
- Bird Studies Canada. 2001. Ontario Breeding Bird Atlas: Guide for Participants. Bird Studies Canada in cooperation with Environmental Canada (Canadian Wildlife Service), Federation of Ontario Field Naturalists, Ontario Field Ornithologists, Ontario Ministry of Natural Resources.
- Bird Studies Canada. 2009. Marsh Monitoring Program Participant's Handbook: Getting Started. 2009 Edition. 13 pages. Published by Bird Studies Canada in cooperation with Environment Canada and the U.S. Environmental Protection Agency. February 2009.
- Burke, Peter S. 2013. Management Plan for the West Virginia White (*Pieris virginianensis*) in Ontario. Ontario Management Plan Series. Prepared for the Ontario Ministry of Natural Resources, Peterborough, Ontario. v + 44 pp.
- Cadman, M.D., D.A. Sutherland, G.G. Beck, D. Lepage, and A.R. Couturier. 2007. The Atlas of the Breeding Birds Ontario 2001-2005. Bird Studies Canada, Environment Canada, Ontario Field Ornithologists, Ontario Ministry of Natural Resources, and Ontario Nature, Toronto, xxii + 706pp.
- Colin Jones, Ross Layberry, and Alan Macnaughton. Ontario Butterfly Atlas Online. 2015. Toronto Entomologists' Association: April 2015. Available at: http://www.ontarioinsects.org/atlas_online.htm
- COSEWIC 2007. COSEWIC assessment and update status report on the redbide dace *Clinostomus Elongates* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vii + 59 pp. Available at: www.sararegistry.gc.ca/status/status_e.cfm
- COSEWIC. 2008a. COSEWIC assessment and status report on the Snapping Turtle *Chelydra serpentina* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vii + 47 pp. Available at: (www.sararegistry.gc.ca/status/status_e.cfm).
- COSEWIC. 2008b. COSEWIC assessment and status report on the Canada Warbler *Wilsonia Canadensis* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vi + 35 pp. Available at: www.sararegistry.gc.ca/status/status_e.cfm
- COSEWIC. 2010. COSEWIC assessment and status report on the Monarch *Danaus plexippus* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vii + 43 pp. Available at: www.sararegistry.gc.ca/status/status_e.cfm
- COSEWIC. 2011a. COSEWIC assessment and status report on the Eastern Meadowlark *Sturnella magna* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. x + 40 pp. Available at: www.sararegistry.gc.ca/status/status_e.cfm

- COSEWIC. 2011b. COSEWIC assessment and status report on the Barn Swallow *Hirundo rustica* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. ix + 37 pp. Available at: www.sararegistry.gc.ca/status/status_e.cfm
- COSEWIC. 2012. COSEWIC assessment and status report on the Eastern Wood-pewee *Contopus virens* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. x + 39 pp. Available at: www.registrelep-sararegistry.gc.ca/default_e.cfm
- COSEWIC. 2013. COSEWIC assessment and status report on the Little Brown Myotis *Myotis lucifugus*, Northern Myotis *Myotis septentrionalis* and Tri-colored Bat *Perimyotis subflavus* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. xxiv + 93 pp. (www.registrelep-sararegistry.gc.ca/default_e.cfm).
- County of Wellington. 2013. Wellington County Official Plan. May 6, 1999 (Last Revision March 24, 2015).
- CRFMP. 2002. Credit River Fisheries Management Plan. Credit Valley Conservation and Ontario Ministry of Natural Resources. Queen's Printer for Ontario. 2002.
- CVC. 2010a. Credit Valley Conservation Authority - Watershed Planning and Regulation Policies. April 2010.
- CVC. 2010b. Credit Valley Conservation Authority, Species of Conservation Concern Project. 2010.
- DFO 2008. A Guide to Understanding Freshwater Fish Habitat in Ontario. Fisheries and Ocean Canada. March 2008
- Dobbyn, John. 1994. Atlas of the Mammals of Ontario. Federation of Ontario Naturalists, viii + 120pp.
- Dougan & Associates, with Snel & Cecile Environmental Research. 2009. List of Significant Wildlife in Wellington County; In City of Guelph Natural Heritage Strategy - Phase 2: Terrestrial Inventory & Natural Heritage System (VOL. 2 – APPENDICES). Final Report March 2009.
- Endangered Species Act, 2007, S.O. 2007, c. 6
- ESSMP. 2011. Erin Service and Settlement Masterplan. Phase 1 – Environmental Component – Existing Conditions Report. May 2011. Town of Erin. Credit Valley Conservation.
- Fisheries Act, 1985. R.S.C., 1985, c. F-14
- GRCA. 2013. Grand River Conservation Authority. *Grand River Information Network*. Available at: <http://www.grandriver.ca/index/document.cfm?sec=63&sub1=0&sub2=0>
- Lee, H.T., W.D. Bakowsky, J.L. Riley, J. Bowles, M. Puddister, P. Uhlig and S. McMurry. 1998. Ecological Land Classification for Southern Ontario: First Approximation and its Application. Ontario Ministry of Natural Resources, Southcentral Science Section. Science Development and Transfer Branch. SCSS Field Guide FG-02

- Lee H.T. 2008. Southern Ontario Ecological Land Classification – Vegetation Type List Ontario Ministry of Natural Resources.
- McCracken, J.D., R.A. Reid, R.B. Renfrew, B. Frej, J.V. Jalava, A. Cowie, and A.R. Couturier. 2013. Recovery Strategy for the Bobolink (*Dolichonyx oryzivorus*) and Eastern Meadowlark (*Sturnella magna*) in Ontario. Ontario Recovery Strategy Series. Prepared for the Ontario Ministry of Natural Resources, Peterborough, Ontario. viii + 88 pp.
- MNRF. 2000. Ontario Ministry of Natural Resources. *Significant Wildlife Habitat Technical Guide*. October 2000.
- MNRF. 2009. The Ecosystems of Ontario, Part 1: Ecozones and Ecoregions. Science and Information Branch. Inventory, Monitoring and Assessment Section. Technical Report SIB TER IMA TR-01. Ministry of Natural Resources and Forestry. 2009
- MNRF. 2010. Ontario Ministry of Natural Resources. *Natural Heritage Reference Manual for Natural Heritage Policies of the Provincial Policy Statement, 2005*. Second Edition. Toronto: Queen's Printer for Ontario.
- MNRF. 2012. Blanding's Turtle Survey Protocol – MNRF Guelph District. April 2012.
- MNRF. 2013. Milksnake Survey Protocol – MNRF Guelph District. June 2013.
- MNRF. 2014. Bat and Bat Habitat Surveys of Treed Habitats – MNRF Guelph District, August 2014.
- MNRF. 2015a. Ontario Ministry of Natural Resources. Ontario Species at Risk website. Available at: <http://www.mnr.gov.on.ca/en/Business/Species/index.html>
- MNRF. 2015b. Significant Wildlife Habitat Ecoregion 6E Criteria Guide. Ministry of Natural Resources and Forestry. Regional Operations Division, Peterborough Ontario, January, 2015.
- MNRF. 2015c. Wellington County Upper Tier Species at Risk. Ministry of Natural Resources and Forestry. Provided by Guelph District MNRF, February, 2015.
- Mortarello, S., Mike Barry, M., Gann G., Zahina, J., Channon, S., Hilsenbeck, C., Scofield, D., Wilder, G., and Wilhelm, G. 2012. Coefficients of Conservatism Values and the Floristic Quality Index for the Vascular Plants of South Florida. *Southeastern Naturalist*. Volume 11(Monograph 3):1–62.
- NatureServe. 2015. *Carex careyana* – Torr. Ex Dewey. NatureServe Explore. An Online Encyclopedia of Life. Available at: <http://explorer.natureserve.org/servlet/NatureServe?searchName=CAREX+CAREYANA>
- NHIC. 2015. Natural Heritage Information Centre. *NHIC online database and mapping*. Ontario Ministry of Natural Resources. Available at: <https://www.ontario.ca/environment-and-energy/natural-heritage-information-centre>

- NRVIS. 2010. Natural Resource Values Information System. Ministry of Natural Recourses. Accessed through GRCA Online Mapping. Available at: <http://www.grandriver.ca/index/document.cfm?Sec=17&Sub1=80>
- OMMHA. 2014. Ontario Ministry of Municipal Affairs and Housing. *Ontario Provincial Policy Statement*. April 30, 2014.
- Ontario Nature. 2015a. *Ontario Reptile and Amphibian Atlas*. Available at: http://www.ontarionature.org/protect/species/herpetofaunal_atlas.php
- Ontario Nature. 2015b. Ontario Reptile and Amphibian Atlas: a citizen science project to map the distribution of Ontario's reptiles and amphibians. Ontario Nature, Ontario. Available: <http://www.ontarionature.org/atlas>; Accessed April 29, 2015].
- Ontario Partners in Flight. 2008. *Ontario Landbird Conservation Plan: Lower Great Lakes/St. Lawrence Plain, North American Bird Conservation Region 13*. Ontario Ministry of Natural Resources, Bird Studies Canada, Environment Canada. Draft Version 2.0
- OWES. 2013. Ontario Ministry of Natural Resources. *Ontario Wetland Evaluation System: Southern Manual*. 3rd Edition.
- Renfrew, Rosalind, Allan M. Strong, Noah G. Perlut, Stephen G. Martin and Thomas A. Gavin. 2015. Bobolink (*Dolichonyx oryzivorus*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Available at: <http://bna.birds.cornell.edu/bna/species/176>
- SARA 2014. Federal Species at Risk Registry. Government of Canada. Available at http://www.registrelep-sararegistry.gc.ca/species/speciesDetails_e.cfm?sid=110
- Town of Erin. 2012a. Corporation of the Town of Erin. Staff Report. Station Street Dam – Hillsburgh. May, 2012.
- Town of Erin. 2012b. The Official Plan of the Town of Erin. Approved by Wellington County Council December 14, 2004. Contains Modification & Applications to May 2012.
- VASCAN. 2015. Database of Vascular Plants of Canada (VASCAN). Université de Montréal Biodiversity Centre. Version 36. Published September 24, 2015.
- Wagner, W. H. and F. S. Wagner. 1982. *Botrychium rugulosum* (Ophioglossaceae), a newly recognized species of evergreen grapefern in the Great Lakes Area of North America. Contributions from the University of Michigan Herbarium.

Agency and Personal Correspondence

Clark, Chris. E.I.T. Triton Engineering Services Limited. Email Correspondence, phone and in-person.

Hale, Lesley. Project Areas Species at Risk Biologist. Ministry of Natural Resources and Forestry, Policy Division. Re: Little Brown Bat Maternity Colony, Banding and Monitoring Project. Dated June 30, 2015. Email Correspondence and in-person.

Slaght, Tyler. Regulations Officer. Credit Valley Conservation. Email Correspondence and in-person.

Buck, Graham. Management Biologist. Ministry of Natural Resources and Forestry, Guelph District. Email Correspondence.


Thompson, Melinda. Management Biologist. Ministry of Natural Resources and Forestry, Guelph District. Email Correspondence.

Timmerman, Art. Management Biologist. Ministry of Natural Resources and Forestry, Guelph District. Email Correspondence

Whalen, Rose. Lands and Waters Technical Specialist. Ministry of Natural Resources and Forestry, Guelph District. Email Correspondence and in-person.

CVC. 2015. Re: Credit Valley Conservation Species of Conservation Concern Project. Dated April 30, 2015.

DRAFT



LEGEND

-  STUDY AREA
-  WATERCOURSES
-  CONSERVATION AUTHORITY BOUNDARY

- Information Sources:
1. Conservation Authority Boundaries: Provided by Land Information Ontario (LIO). Accessed 2015.
 2. Watercourse: Provided by Land Information Ontario (LIO). Accessed 2015.
 3. Roads: Provided by Land Information Ontario (LIO). Accessed 2015.
 4. Ortho Image: Provided by First Base Solutions Web Mapping Service 2015.



Title: SITE LOCATION
 Project: HILLSBURGH DAM MUNICIPAL CLASS EA

Date: Dec, 2015
 Project: AA12-137A
 Scale: 1 : 100000




ABOUT & ASSOCIATES INC.
 Consulting Arborists • Ecologists • Landscape Designers
 591 Woolwich Street • Guelph • Ontario • Canada • N1H 3Y5
 T: 519.822.6839 • F: 519.822.4052 • info@aboutdng.com • www.aboutdng.com

Figure No: **1**



LEGEND

-  WATERCOURSE
-  ELORA CATARACT TRAILWAY
-  STUDY AREA
-  CVC BOTANICAL STUDY AREA
-  OPEN WATER COMMUNITIES
-  PROPERTY ACCESS
-  HILLSBURGH DAM

- Information Sources:
1. Property Access: Based on Erin Parcels. Provided Wellington County under data sharing agreement. Accessed 2015. Modified by AA based on Property Access provided.
 2. CVC Botany Study Area: Based on Peel Region Natural Areas Inventory Accessed 2015.
 3. Open Water Communities: Provided by Wellington County under data sharing agreement. Accessed 2015.
 4. Watercourse: Provided by Land Information Ontario (LIO) Accessed 2015.
 5. Roads: Provided by Wellington County under data sharing agreement. Accessed 2015.
 6. Ortho Image: Wellington County 2010. Provided by First Base Solutions Web Mapping Service 2015.




Title: PROPERTY ACCESS & CVC BOTANICAL SURVEY AREA
 Project: HILLSBURGH DAM MUNICIPAL CLASS EA

Date: Dec, 2015
 Project: AA12-137A
 Scale: 1 : 8000



Figure No: 2



ECOLOGICAL LAND CLASSIFICATION		
POLYGON	ELC CODE	COMMUNITY
1	MAMO1-2	Cattail Graminoid Organic Mineral Meadow Marsh
2	SWCM1-2	White Cedar - Conifer Mineral Coniferous Swamp
3	SWMO3-3	White Birch - Conifer Organic Mixed Swamp
4	FODM5-8	Dry-Fresh Sugar Maple - White Ash Deciduous Forest
5	FOCM2-2	Dry-Fresh White Cedar Coniferous Forest
6	FOCM6	Naturalized Coniferous Plantation
7	SAS_1	Submerged Shallow Aquatic Ecosite
8	MASO1-1	Cattail Organic Shallow Marsh Type
9	SWTO3-5	Red-osier Organic Deciduous Swamp
10	SWMO1-1	White Cedar - Hardwood Organic Mixed Swamp
11	SAM_1-8	Water Lily - Bullhead lily Mixed Shallow Aquatic
12	MEMM3	Dry - Fresh Mixed Meadow Ecosite
13	CS	Cultural Savannah
14	FETC1-2	Tamarack - White Cedar Treed Fen
15	FODM8-1	Fresh - Moist Poplar Deciduous Forest
16	FODM6	Fresh - Moist Sugar Maple Deciduous Forest Ecosite
17	SWCM1-2	White Cedar - Conifer Mineral Coniferous Swamp
18	SAM_1-8	Water Lily - Bullhead lily Mixed Shallow Aquatic
19	SAM_1-8	Water Lily - Bullhead lily Mixed Shallow Aquatic
20	OAW	Open Aquatic
21	SWCM1-2	White Cedar - Conifer Mineral Coniferous Swamp
22	SWTO2-6	Mixed Willow Organic Thicket Swamp Type
23	FOMM7-2	Fresh - Moist White Cedar - Hardwood Mixed Forest
24	SWDM4-5	Poplar Mineral Deciduous Swamp
25	FODM8-1	Fresh - Moist Poplar Deciduous Forest
26	SWDM2-1	Black Ash Mineral Deciduous Swamp
27	FOCM6	Naturalized Coniferous Plantation
28	SWTO2-3	Meadow Willow Organic Deciduous Thicket Swamp
29	SWDM4-5	Poplar Mineral Deciduous Swamp
30	FODM7-7	Fresh - Moist Manitoba Maple Lowland Deciduous Forest
31	MAMM1-1	Cattail Graminoid Mineral Meadow Marsh Type
32	RES	Low Density Residential

LEGEND

- ECOLOGICAL LAND CLASSIFICATION COMMUNITIES
- WATERCOURSE (West Credit River)
- ELORA CATARACT TRAILWAY

Information Sources:
 1. Ecological Land Classification Communities Assessed and Mapped by Aboud & Associates Inc. 2015
 2. Roads. Provided by Wellington County under data sharing agreement. Accessed 2015.
 3. Watercourse. Provided by Land Information Ontario (LIO) Accessed 2015.
 4. Ortho Image. Wellington County 2010. Provided by First Base Solutions Web Mapping Service 2015.



Title: **ECOLOGICAL LAND CLASSIFICATION**
 Project: **HILLSBURGH DAM MUNICIPAL CLASS EA**

Date: Dec, 2015
 Project: AA12-137A
 Scale: 1 : 8000



ABOUT & ASSOCIATES INC.
 Consulting Arborists • Ecologists • Landscape Designers
 591 Woolwich Street • Guelph • Ontario • Canada • N1H 3Y5
 T: 519.822.6839 • F: 519.822.4052 • info@abouding.com • www.abouding.com



Legend

- STUDY AREA
- WINTER WILDLIFE SURVEY TRANSECT
- BIRD MIGRATION TRANSECT (1-5)
- BREEDING BIRD POINT COUNT (1-10)
- AMPHIBIAN SURVEY (A-H)
- CANDIDATE WILDLIFE HABITAT
- SNAKES (S1-S4)
- TURTLES (T1-T5)
- MARSH BIRDS (MB1-MB4)

Information Sources:
 1. Roads. Provided by Wellington County under data sharing agreement. Accessed 2015.
 2. Ortho Image. Wellington County 2010. Provided by First Base Solutions Web Mapping Service 2015.



Title: WILDLIFE HABITAT TARGETS & SURVEY LOCATIONS


Project: HILLSBURGH DAM MUNICIPAL CLASS EA

Date: Dec, 2015
 Project: AA12-137A
 Scale: 1 : 8000



ABOUT & ASSOCIATES INC.
 Consulting Arborists • Ecologists • Landscape Designers
 591 Woolwich Street, Guelph, Ontario, Canada, N1H 3Y5
 T: 519.822.6839 • F: 519.822.4052 • info@aboutinc.com • www.aboutinc.com

Figure No:



LEGEND

- | | |
|---|--|
|  STUDY AREA |  COLD WATER |
|  ELORA CATARACT TRAILWAY |  COOL WATER |
| FISH PASSAGE |  WARM WATER |
|  FULL BARRIER |  AQUATIC HABITAT SEGMENTS |
|  PARTIAL BARRIER | |


Information Sources:
 1. Fish Passage Barriers Assessed and Mapped by Aboud & Associates Inc. 2015
 2. Trout Spawning Habitat Assessed and Mapped by Aboud & Associates Inc. 2015
 3. Fish Communities: Assessed and Mapped by CVP as shown in Erin Service and Settlement Master Plan, 2011.
 4. Watercourse: Provided by Land Information Ontario (LIO) Accessed 2015.
 5. Roads: Provided by Wellington County under data sharing agreement. Accessed 2015.
 6. Ortho Image: Wellington County 2010. Provided by First Base Solutions Web Mapping Service 2015.



Title: AQUATIC HABITAT ASSESSMENT
 Project: HILLSBURGH DAM MUNICIPAL CLASS EA

Date: Dec, 2015
 Project: AA12-137A
 Scale: 1 : 8000





LEGEND

STUDY AREA	OPEN WATER COMMUNITIES
WATERCOURSES	WETLANDS (PSW)
ELORA CATARACT TRAILWAY	West Credit River Wetland Complex
WOODED AREAS	Alton Hillsburgh Wetland Complex
HILLSBURGH DAM	Speed Lutteral Swan Creek Wetland Complex

Information Sources:

1. Wooded Areas: Provided by Land Information Ontario (LIO) Accessed 2015.
2. Wetland: Provided by Land Information Ontario (LIO) Accessed 2015.
3. Open Water Communities: Provided by Wellington County under data sharing agreement. Accessed 2015.
4. Watercourse: Provided by Land Information Ontario (LIO) Accessed 2015.
5. Roads: Provided by Wellington County under data sharing agreement. Accessed 2015.
6. Ortho Image: Wellington County 2010. Provided by First Base Solutions Web Mapping Service 2015.

TOWN OF ERIN
PROGRESS IN UNITY

TRITON
ENGINEERING
SERVICES
LIMITED
Consulting Engineers

Title: **LANDSCAPE EVALUATION**

Project: **HILLSBURGH DAM MUNICIPAL CLASS EA**

Date: Dec, 2015

Project: AA12-137A

Scale: 1 : 20000

ABOUT & ASSOCIATES INC.
Consulting Arborists • Ecologists • Landscape Designers
591 Woolwich Street • Guelph • Ontario • Canada • N1H 3Y5
T: 519.822.6839 • F: 519.822.4052 • info@aboutdng.com • www.aboutdng.com

Figure No: **6**

SIGNIFICANT SPECIES OBSERVATION

- ★ Little Brown Myotis (END)
- ★ Boboblink (THR)
- ★ Eastern Meadowlark (THR)
- ★ Bald Eagle (SC)
- ★ Eastern Wood-Pewee (SC)
- ★ Snapping Turtle (SC)
- ★ Great Egret (CVC Tier 1)
- ★ Trumpeter Swan (CVC Tier 1)

SIGNIFICANT WILDLIFE HABITAT (SWH)

SWH	SWH Types
1	Waterfowl Stopover, Turtle Overwintering, Special Concern Species, Amphibian Breeding
2	Turtle Overwintering, Special Concern Species
3	Special Concern Species
4	Special Concern Species, Bat Maternity Habitat
5	Special Concern Species
6	Bat Maternity Habitat
7	Bat Maternity Habitat
8	Bat Maternity Habitat
9	Bat Maternity Habitat
10	Bat Maternity Habitat
11	Bat Maternity Habitat
12	Bat Maternity Habitat
13	Amphibian Breeding
14	Amphibian Breeding
15	Amphibian Breeding



LEGEND

- STUDY AREA
- SIGNIFICANT WILDLIFE HABITAT (SWH)

Information Sources:
 1. Significant wildlife Habitat and Significant Species Observations Assessed and Mapped by Aboud & Associates Inc. 2015
 2. Roads. Provided by Wellington County under data sharing agreement. Accessed 2015.
 3. Ortho Image. Wellington County 2010. Provided by First Base Solutions Web Mapping Service 2015.
 4. Significant Species Status from SARO (2015) and CVC Species of Special Concern Project (2015).



Title: SIGNIFICANT WILDLIFE OBSERVATIONS & SIGNIFICANT WILDLIFE HABITAT

Project: HILLSBURGH DAM MUNICIPAL CLASS EA

Date: Dec, 2015
 Project: AA12-137A
 Scale: 1 : 8000



ABOUT & ASSOCIATES INC.
 Consulting Arborists • Ecologists • Landscape Designers
 591 Woolwich Street • Guelph • Ontario • Canada • N1H 3Y5
 T: 519.822.6839 • F: 519.822.4052 • info@aboudinc.com • www.aboudinc.com

Figure No:

Appendix 1
Notice of Study Commencement

DRAFT



December 19, 2014

Dear Mr. Ziegler:

**Re: Class Environmental Assessment Study – Schedule B
Proposal for the improved safety of the Hillsburgh Dam and Bridge
Town of Erin**

Further to receipt of the Notice of Commencement, November 27, 2014, CVC staff offer the following preliminary comments:

It is the understanding of CVC staff that the Town of Erin is undertaking a Class Environmental Assessment (EA) for the purpose of improving safety of the Hillsburgh Dam and bridge.

Site Characteristics:

The study area is traversed by the West Credit River and a tributary of the West Credit River, associated hazards (flooding and erosion). In addition, the area contains wetlands and associated adjacent lands. As a result portions of the study area are subject to the Authority's Development, Interference with Wetlands, and Alterations to Shorelines & Watercourses Regulation (Ontario Regulation 160/06). This regulation prohibits altering a watercourse or wetland and prohibits development within the regulated area without the prior written approval of CVC (i.e. a permit).

Permit Approval Requirements:

In accordance with Ontario Regulation 160/06 (our Development, Interference with Wetlands and Alterations to Shorelines and Watercourses Regulation), a permit would be required from the CVC prior to commencement of the works involving development, interference with a wetland and/or alterations to a watercourse or shoreline.

Fish Habitat and Department of Fisheries and Oceans (DFO):

Please note that CVC is no longer administering the *Fisheries Act* on behalf of Fisheries and Oceans Canada (DFO). As a result, it is up to the proponent to ensure that his/her project meets the DFO requirements under the self-assessment process. See the link below for a description of the self assessment process and a list of projects/activities where DFO review is not required: <http://www.dfo-mpo.gc.ca/pnw-pppe/index-eng.html>. Measures to avoid causing harm are noted here: <http://www.dfo-mpo.gc.ca/pnw-pppe/measures-mesures/index-eng.html>

EA Study Objectives:

The EA Study must clearly identify and quantify the environmental constraints and enhancement opportunities within the study area, including the following:

Aquatic Habitat and Valleylands:

The project needs to evaluate alternatives that minimize impacts to the form and function of the West Credit River and Hillsburgh Pond and if possible include opportunities for enhancement. The EA should list and describe the natural features (fish habitat, etc.) and site characteristics (e.g. Rolling topography, high water table, buffering vegetation, etc.) in the study area that may pose constraints to the project.

December 19, 2014

Re: **Proposal for the improved safety of the Hillsburgh Dam and Bridge**
Town of Erin

Stormwater Management:

The project should include quality and quantity control measures to treat stormwater runoff in accordance with Ministry of Environment and CVC guidelines. Typically we request that the proponent provide treatment for all new proposed impervious areas and where possible existing road surfaces.

Hydraulics and Meander Belt

Any alterations to any watercourse crossings may require a hydraulic analysis to ensure that there are no negative up or down stream impacts. In addition, the road improvements or reconstruction at a minimum should maintain existing depth flooding on the road or improve the road such that it is flood free under Regional Storm conditions. In addition, CVC requests that new bridges and where possible replacement bridges span the calculated meander belt of the watercourse.

Subwatershed Study:

The EA should ensure that the subwatershed study environmental targets and objectives are identified and identify proposed measures that implemented these targets and objectives.

Erosion and Sediment Controls

During the detailed design period of this project, all proposed methods to control sedimentation during construction and potential erosion following the completion of the project must be detailed. Furthermore, as means of minimizing impacts to aquatic habitat all works must be completed in the dry.

Restoration

All disturbed areas will need to be stabilized and restored with native/non-invasive seed mixes and woody species.

Given CVC's interest staff would like to be kept informed of future meetings and proceedings through the Environmental Assessment process. Please forward any information or reports when available to ensure that this Authority's policy and program interest are reflected in the planning and design components for this project.

Should you have any further questions please contact the undersigned at (905) 670-1615 extension 406

Regards,



Tyler Slaght
Regulations Officer

cc: **Town of Erin**
Attention: Larry Van Wyck (via email)

Ministry of Natural Resources and Forestry Guelph District
Attention: Rose Whalen (via email)

Ministry of Natural Resources and Forestry Peterborough District
Attention: Doug Ryan (via email)

Ryan Hamelin

From: Buck, Graham (MNRF) <Graham.Buck@ontario.ca>
Sent: April-02-15 1:45 PM
To: Ryan Hamelin
Cc: Whalen, Rose (MNRF)
Subject: RE: Hillsburgh SAR

Hi Ryan,

With respect to bats we generally only recommend surveys if there is the potential for impacts to the hibernation or roost habitats. If the impact is only to foraging habitat I do not think the activity will damage or destroy habitat.

With respect to Rusty-patched Bumblebee I agree the likelihood is extremely low, given the number of hours of survey completed for this species with very few individuals seen (3, all at Pinery). I would accept incidental observations during plant surveys. Staff can always take pictures of Bumblebees and submit them to <http://bumblebeewatch.org/> if they are not confident if identification.

The 20102 protocol for Blanding's is still acceptable.

I will follow up with the survey protocols in another email.

Graham

*Graham Buck
Management Biologist
Ministry of Natural Resources and Forestry
1 Stone Road West
Guelph ON
N1G 4Y2
519 826 4505
graham.buck@ontario.ca*

From: Ryan Hamelin [mailto:ryan@abouttng.com]
Sent: March-25-15 3:59 PM
To: Whalen, Rose (MNRF); Buck, Graham (MNRF)
Cc: Chris Clark; Paul Ziegler
Subject: RE: Hillsburgh SAR

Hi Rose and Graham,

As part of the Hillsburgh Dam EA the CVC has requested Targeted Surveys of all SAR with potential habitat within the Study Area.

Based on some winter field work and orthophotography interpretations, I have developed a proposed list of SAR to be surveyed for based on the identified habitat features. Attached is an Excel file with a comprehensive explanation of the proposed species to be surveyed for, and what methods will be used.

Would the MNRF be able to provide the survey protocols for the following species? Do any of the survey protocols require handling permits or specific permission?

- Jefferson Salamander
- Eastern Small Footed Myotis
- Little Brown Myotis
- Northern Myotis
- Butlers Garter Snake
- Eastern Ribbon Snake
- Massasauga Rattlesnake
- Milk Snake
- Rusty-patched Bumble Bee

If possible could you comment on the need or benefit of surveying for the three SAR bat species? It would be expected that bats would use the study area for feeding, but there are no known caves, abandoned mines, cliffs or rock outcrops that could be used as over wintering habitat within the study area. Due to the lack of these Key Habitat features it is felt that targeted bat survey may not be necessary. The habitat is also not ideal for rusty-patched bumble bees and potential EA options would be unlikely to impact their habitat, could you also comment on the need for Rusty-patched Bumble Bee surveys ?

Could you confirm the appropriateness of using the 2012 Blanding's Turtle Survey Protocol to survey for the presence of Snapping Turtles and Spotted Turtles?

If it would be easier to discuss any of this information over the phone please feel free to call me.

Thanks for the assistance,

Ryan Hamelin

Ryan Hamelin, B.S.c (Env). M.Sc. Terrestrial and Wetland Ecologist
ABOUD & ASSOCIATES INC. 591 Woolwich Street . Guelph . Ontario . N1H 3Y5
T:519.822.6839 x 2 . F:519.822.4052 www.aboudtng.com . ryan@aboudtng.com

From: Buck, Graham (MNRF) [<mailto:Graham.Buck@ontario.ca>]

Sent: February-10-15 11:48 AM

To: Ryan Hamelin

Subject: RE: SAR List

I have attached the most recent version of the Wellington County list.

It is always best to use the Guelph District list over the lists available online because the Guelph District list is more comprehensive and up to date.

The online lists should only be used in instances where the district does not maintain a species at risk by municipality list.

*Graham Buck
Management Biologist
Ministry of Natural Resources and Forestry
1 Stone Road West
Guelph ON
N1G 4Y2*

519 826 4505
graham.buck@ontario.ca

From: Ryan Hamelin [<mailto:ryan@aboudtng.com>]
Sent: February-10-15 11:11 AM
To: Buck, Graham (MNRF)
Subject: SAR List

Hi Graham,

I am developing a list of SAR that have the potential to be present at the Hillsburgh Dam Study Area. We have a list of SAR known to occur in Wellington County that was supplied by the MNRF in the summer of 2013. I have also consulted the MNRF online – SAR by Area web-mapper for Wellington County (<http://www.ontario.ca/environment-and-energy/species-risk-area>). I have noticed that the two lists have some inconsistency in species. Could you confirm what SAR list should be used when assessing for potential SAR within Wellington County?

I have attached a copy of the original SAR list supplied by the MNRF, with an additional column indicating what species are also listed on the MNRF website for Wellington County.

Thanks,

Ryan Hamelin

From: Buck, Graham (MNRF) [<mailto:Graham.Buck@ontario.ca>]
Sent: January-19-15 11:18 AM
To: Ryan Hamelin
Subject: RE: SAR / Turtle Survey

Hello Ryan,

I have attached the survey protocol for Blanding's Turtle. It has been finalized and can be used for this species. You may also be able to adapt it for other species.

MNRF Guelph also has draft survey protocols for Stinkpot, Spotted and Wood Turtle but they are not finalized and not applicable to Hillsburgh Dam EA. They are also very species specific and likely not as transferable to other turtles.

*Graham Buck
Management Biologist
Ministry of Natural Resources and Forestry
1 Stone Road West
Guelph ON
N1G 4Y2
519 826 4505
graham.buck@ontario.ca*

From: Ryan Hamelin [<mailto:ryan@aboudtng.com>]
Sent: December-23-14 4:39 PM
To: Buck, Graham (MNRF)
Cc: Whalen, Rose (MNRF)
Subject: SAR / Turtle Survey

Hello Graham Buck,

I hope you are doing well.

We are in the process of developing a work plan for a Municipal Class Environmental Assessment and are interested in Species at Risk and Turtle Surveys. Would you be able to provide any information on typical requirements for Species at Risk surveys and Turtle Surveys as part of Municipal Class Environmental Assessments, Schedule C ? Also, are there specific protocols or techniques that the MNRF endorses for Species at Risk and Turtle Surveys ?

The specific project and site of interest is the Hillsburgh Dam Environmental Assessment in the Township of Erin. Rose Whalen is the primary MNRF contact for the project. The subject property falls within CVC's watershed and they have asked about surveys for 'all' Species at Risk that could possibly be present.

Thanks,

Ryan Hamelin

Ryan Hamelin, B.S.c (Env). M.Sc. Terrestrial and Wetland Ecologist
ABOUD & ASSOCIATES INC. 591 Woolwich Street . Guelph . Ontario . N1H 3Y5
T:519.822.6839 x 2 . F:519.822.4052 www.aboudtng.com . ryan@aboudtng.com

DRAFT

Archived: December-04-15 1:46:29 PM
From: Hale, Lesley (MNRF)
Sent: June-30-15 11:00:38 AM
To: Cheryl-Anne Ross
Subject: RE: Hillsburgh mill Pond Bat study
Importance: Normal

Hi Cheryl

Here is a summary of the data from the last few years for the Hillsburgh church bat roost (please note the 2015 data has not been input yet):

Site Name	Primary Surveyor	Date	Sky Code	Wind Code	Temp	Start Time	End Time	Total Bats	Technique	Other Surveyors	Comments
Hillsburgh Church	Lesley Hale	June 20, 2012	1	1	26	21:34	22:23	113	visual	Paul Faure, Les Misch, John (Ryan) Caldwell, Lucas Greville	Mist nets set; 3 MYLU captured and banded 002453-002455
Hillsburgh Church	Lesley Hale	July 12, 2012	1	1	24	21:20	22:20	174	visual	Heather Riddell, Iga Stasiak, Ryan Caldwell	Mist nets set; 10 MYLU captured and banded; 002457-58, 002494-002498, 002500, 002483-84
Hillsburgh Church	Heather Riddell	July 4, 2013	1	1	22	21:20	22:08	125	visual	Iga Stasiak, Catherine Jong, Ryan Morin	5 MYLU captured in the mist net (2 mist nets set). 17 bats observed at the front; 108 observed at the back. Not enough surveyors for the side walls of the building.
Hillsburgh Church	Heather Riddell	July 23, 2013	3	4	15	21:01	22:00	106	visual	Les Misch, Dan Bourassa, Alejandra	7 MYLU captured, 2 released accidentally. All male bats. 1 mist net set.
Hillsburgh Church	Lesley Hale	June 9, 2014	1	1	18	21:30	22:15	110	visual	Benoit Talbot, Christy Humphrey	25 exited from south side near centre apex; 85 exited from west side from apex triangle; 6 returned after 22:00; 2 nets set along west line of trees; 17 MYLU and 1 EPFU captured and banded
Hillsburgh Church	Lesley Hale	July 10, 2014	2	1	15	21:15	22:23	219	visual	Christy Humphrey, Christina Smyth	200 bats exited from west wall near peak and 19 exited from south wall; 2 nets erected 37 MYLU captured and banded - no recaptures

As I mentioned, please feel free to join us for the next survey on July 20 (July 21 if it rains on 20th) at 8pm at the church. Bring a headlamp and a folding camp chair (it's a long night otherwise!) and be ready for lots of bugs. My cell number is 705-917-0373 in case you need to reach me.

Cheers

Lesley Hale

From: Cheryl-Anne Ross [mailto:Cheryl@aboudtng.com]
Sent: June-19-15 11:53 AM
To: Hale, Lesley (MNRF)
Cc: Ryan Hamelin
Subject: Hillsburgh mill Pond Bat study

Hi Lesley,

We've received your contact information from one of the councillors in the town of Erin (Jeff Duncan), regarding an ongoing study of a little brown bat population at the century Theater, near the station Street Dam. As your voicemail indicated you are out of the office quite a bit this week, I thought I would send an email to follow up on some of the information that we were provided.

About & Associates is currently gathering Existing Information for a municipal class EA for the Station Street Dam, examining alternatives for the failing Dam, working closely with the Local MNRF branch, and the Credit Valley Conservation authority. It's recently come to our attention that the MNRF has been conducting bat assessment work in the vicinity of the station street Dam, and we would really appreciate any further information you could provide to us about the population. Particularly if any known maternity colonies may exist in the forested landscape in the vicinity of the Dam, as well as a discussion about protection of foraging habitat, and if there are any guidelines regarding foraging habitat or regulated habitat that might be in place. I am available until 4:00 pm today, and should be in the office all day Monday, if you would like to give me a call at the number below, my extension is 7.

Thank you,



**TRITON
ENGINEERING
SERVICES
LIMITED**
Consulting Engineers

MEETING MINUTES

DATE: Wednesday, September 24, 2014

TIME: 10:00 a.m.

LOCATION: Credit Valley Conservation Authority
Headquarters – 1255 Old Derry Road,
Mississauga

OUR FILE: A4685E

RE: HILLSBURGH DAM, MUNICIPAL
CLASS ENVIRONMENTAL
ASSESSMENT, TOWN OF ERIN

Attendees:

T. Slaght, J. Wong, J. Clayton; Credit Valley Conservation Authority (CVC)
R. Whalen, D. Ryan; Ministry of Natural Resources and Forestry (MNR)
L. Van Wyck; Town of Erin (Town)
S. Aboud, R. Hamelin; Aboud & Associates Inc. (Aboud)
C. Clark; Triton Engineering Services Limited (TESL)

1. C. Clark reviewed the projects history, from temporary works completed to repair the Dam/Road in 2011/2012 to the present permanent solution involving the completion of a Municipal Class Environmental Assessment (Class EA) to fulfill the requirements of the MNR's Non-Application Emergency Works under the Lakes and Rivers Improvement Act.
2. D. Ryan asked if project Problem Statement has been formed. C. Clark to detail Problem Statement and circulate to project team. To be included as part of the Class EA Notice of Project Commencement, to be released shortly.
3. Aboud presented the proposed project Study Area with respect to the Natural Heritage investigations. MNR and CVC recommended extension to Wellington Road 22 east to Trafalgar Rd. Logic behind Study Area was connectivity and impact to Provincially Significant Wetlands (PSW).
4. Aboud overviewed the required Natural Heritage information that is still outstanding for "desktop investigations".
 - a. All data requests to go through T. Slaght (CVC) and R. Whalen (MNR)
 - b. All Natural Heritage data requests are to come from Aboud
 - c. Aboud to submit revised Study Area to MNR/CVC as part of formal data requests
 - d. R. Whalen to provide mapping of wetland evaluations, if available

5. Aboud will be utilizing the services of Aquafor Beech for fish habitat investigations. Data required for this portion of investigations are as follows; fish community data (presence/absence, biomass), thermal regimes and temp data, spawning survey, benthic macroinverts, geomorphology, invasive species info (Round Goby) and fish habitat assessment.
6. J. Clayton overviewed available fish related data. This includes periodic fish inventories from 1954 to present, fish biomass collection, thermal records, presence of invasive Round Goby, spawning data (2010 - 2014).
 - a. Temperature loggers currently on-site and logging and could be left longer into the fall/winter season if required.
 - b. Groundwater seeps throughout system, but no specific locations identified in study area.
 - c. Area is historically Brook Trout habitat, with population currently upstream and downstream of the pond.
 - d. CVC considers the Banded Killifish and the Slimy Sculpin as important species due to the rarity in the watershed.
7. J. Clayton added that Round Goby control methods may be implemented this fall or next spring within the Hillsburgh Pond and other affected ponds along the watercourse. This would involve lowering the water levels and removing desirable fish species.
 - a. May be an opportunity to inventory fish species at this time
 - b. During previous public contact related to Goby eradication, public was opposed temporary water drawdowns
8. According to MNRF, there is no known presence of Species at Risk (SAR) within the Study Area. This will be confirmed through Aboud's desktop/field investigations.
9. C. Clark reviewed existing hydraulic data completed as part of the temporary works, as well as, the Dam's "High" Hazard Potential Classification (HPC). CVC agreed to share any relevant data/information completed or acquired post temporary dam repair works in order to perform any additional analysis.
 - a. T. Slaght - CVC main criteria when evaluating EA options will be; 1) Flood hazard reduction 2) Sediment/Erosion impact reduction. There must be no negative impacts to flooding or erosion. The options reviewed should seek to improve these conditions, as well as; improve natural heritage features present. Flooding and erosion must be demonstrated as part of the Project File Report while sediment control can be established during the detailed design stage.
10. It was agreed that Geomorphology and Hydrogeology investigations be completed as part of the Class EA to cover all areas for the potential alternative outcomes.
 - a. CVC has 2005 fluvial geomorphology data completed by PEIL for West Credit Watershed, which can be provided.
 - b. Provincial Groundwater Monitoring Network and Source Water Protection data may help to provide background. Local water bottling company may also be a source of groundwater data.

11. C. Clark and L. Van Wyck reminded the group of the potential restrictions affecting the Class EA due to Town's property limitations. The Town owns the Station Street road right-of-way but not the north and south adjacent properties. A number of field investigations will need to be performed at these locations. The north landowner also owns the Hillsburgh Pond's stop-log control structure.
 - a. D. Ryan reminded everyone of the adjacent landowners "riparian interests" to the Dam. This involves holding their concerns/interests at stake. Further, the Town can perform the Class EA to uphold their responsibilities to the Dam. Adjacent landowner has legislative responsibilities if dam were to fail.
 - b. As the north adjacent landowner and the Town are affiliated "dam owners" and the requirements for land access to south pond (Ainsworth Pond), it was suggested and agreed that a personal letter be distributed to these parties to request their involvement in the Class EA process. This could eliminate any property access restrictions.
12. General discussion of how potential options could affect the existing PSW wetland complex. Due to the overall size of the Provincially Significant West Credit Wetland Complex it would be expected that a local reduction in extent around the Station Street Dam site would not affect the PSW status of the complex as a whole. However, specific areas that transition from wetland to upland due to changes in hydrology would no longer be included in the wetland complex and would therefore not have PSW status. However, this is to be determined as part of the Class EA process.
13. Those parties and members present at this meeting will be the main Project Team moving forward.
 - a. Technical reporting and any project status updates to be provided approximately every three months.
 - b. All documents will be reviewed by the Committee before release to the Public.
 - c. Meetings will be scheduled as needed.

Project Next Steps/Actions Items:

1. Natural Heritage Study Area and project Terms of Reference to be completed and circulated to Committee for data requisitions.
2. Draft a letter to send to adjacent landowners requesting their personal involvement in the Class EA process.
3. Problem/Opportunity Statement to be developed and included in the Notice of Project Commencement to be released to public.
4. Next Status Up-date Mid-December 2014

Archived: December-04-15 1:49:37 PM
From: Thompson, Melinda (MNRF)
Sent: December-04-15 12:29:01 PM
To: Cheryl-Anne Ross
Cc: Buck, Graham (MNRF); Whalen, Rose (MNRF)
Subject: RE: hillsburgh dam EA project - Ecoregion Criteria clarification
Importance: Normal

Hi Cheryl

I believe our interpretation is that you need to have an amphibian breeding population of at least 20 individuals (this can be combined across species).

Melinda

MELINDA J. THOMPSON ❁ ❁ ❁ ❁ ❁

MANAGEMENT BIOLOGIST | ONTARIO MINISTRY of NATURAL RESOURCES and FORESTRY | GUELPH DISTRICT OFFICE
1 Stone Road West, Guelph, Ontario, N1G 4Y2 | ☎ 519.826.6543 | ✉ melinda.thompson@ontario.ca
[Learn more about Ontario's Species at Risk](#)

From: Cheryl-Anne Ross [<mailto:Cheryl@aboutdng.com>]
Sent: November 24, 2015 11:57 AM
To: Whalen, Rose (MNRF)
Cc: Buck, Graham (MNRF); Ryan Hamelin
Subject: hillsburgh dam EA project - Ecoregion Criteria clarification

Hi Rose,

We are working on completing the existing conditions report for the Hillsburgh Dam project, and I've run into some confusion in interpreting the defining criteria for confirmed SWH.

For Amphibian Breeding Habitat (Woodland), it states that:

'a population of 2 or more of the listed frog species with at least 20 individuals or 2 or more of the listed frog species with call level codes of 3' are to be considered significant.

Does this indicate that if a station had, for example, 5 species of frogs, but did not have greater than 20 individuals of any two species, that the site would not be considered significant? Despite the fact that the site may have had greater than 20 individual frogs combined across species?

Any help with interpreting this criteria would be greatly appreciated!

Thank you,

Regards,
Cheryl-Anne Ross B.Sc. .

Appendix 2
Terms of Reference and Approval

DRAFT



591 Woolwich Street
Guelph . Ontario
N1H 3Y5

T: 519.822.6839
F: 519.822.4052
info@aboutdng.com
www.aboutdng.com

URBAN FORESTRY
ARBORIST REPORTS
MANAGEMENT PLANS
TREE PRESERVATION PLANS
TREE RISK ASSESSMENT
GIS TREE INVENTORIES
TREE APPRAISALS
MONITORING

ECOLOGICAL RESTORATION
NATURAL SYSTEMS DESIGN
HABITAT RESTORATION
EDGE MANAGEMENT PLANS
RAVINE STEWARDSHIP PLANS
NATURALIZATION PLANS
INTERPRETIVE DESIGN
MONITORING
CONTRACT ADMINISTRATION

ENVIRONMENTAL STUDIES
SUBWATERSHED STUDIES
ENVIRONMENTAL IMPACT
STATEMENTS
ECOLOGICAL LAND
CLASSIFICATION
WETLAND EVALUATION
VEGETATION ASSESSMENT
BOTANICAL INVENTORIES
WILDLIFE SURVEYS
MONITORING

LANDSCAPE ARCHITECTURE
MASTER PLANNING
RESIDENTIAL COMMUNITIES
COMMERCIAL/INDUSTRIAL
HEALTHCARE AND EDUCATION
STREETSCAPES
PARKS AND OPEN SPACES
TRAIL SYSTEMS
GREEN ROOFS
CONTRACT ADMINISTRATION

EXPERT OPINION
OMB TESTIMONY
LEGAL PROCEEDINGS
PEER REVIEW
RESEARCH
EDUCATION

December 8, 2014

Our Project No: AA12-137A
Sent by email: cclark@trotoneng.on.ca

Tyler Slaght
Credit Valley Conservation
1255 Old Derry Road
Mississauga, Ontario
L5N 6R4

Care of:

Chris Clark
Triton Engineering Services Limited
105 Queen Street West, Unit 14
Fergus, Ontario
N1M 1S6

Re: Terms of Reference for Hillsburgh Dam Natural Heritage Existing Conditions Report as part of the Municipal Class Environmental Assessment


Dear Tyler,

This letter outlines the draft Terms of Reference (ToR) of the Hillsburgh Dam Natural Heritage Existing Conditions Report. This report is part of the Municipal Class Environmental Assessment (EA) to address the structural state of the existing earthen berm and dam.

Background and Context

The proposed study area for the project is a total of 78.5 hectares, centered on the Hillsburgh Dam and extending up stream to include the Hillsburgh pond, surrounding wetland and associated tributary sections. The study area also extends downstream from the dam to Wellington Road 22 and includes the associated wetlands and woodlands (see Natural Heritage Study Area Map). The larger landscape level context of the area will also be examined to evaluate the significance of the natural heritage features within the broader region.

The study area is contained entirely within the Town of Erin's municipal boundaries and the Credit Valley Conservation's (CVC) jurisdiction. The majority of the study area contains naturalized environments and hosts a wide variety of flora and fauna. Large sections of the study area contain part of the Provincially Significant West Credit Wetland Complex. The open water community of the Hillsburgh pond, created by the Hillsburgh dam is considered a 'rare community' within the region



591 Woolwich Street
Guelph . Ontario
N1H 3Y5

T: 519.822.6839

F: 519.822.4052

info@aboutdng.com

www.aboudtng.com

URBAN FORESTRY

ARBORIST REPORTS
MANAGEMENT PLANS
TREE PRESERVATION PLANS
TREE RISK ASSESSMENT
GIS TREE INVENTORIES
TREE APPRAISALS
MONITORING

ECOLOGICAL RESTORATION

NATURAL SYSTEMS DESIGN
HABITAT RESTORATION
EDGE MANAGEMENT PLANS
RAVINE STEWARDSHIP PLANS
NATURALIZATION PLANS
INTERPRETIVE DESIGN
MONITORING
CONTRACT ADMINISTRATION

ENVIRONMENTAL STUDIES

SUBWATERSHED STUDIES
ENVIRONMENTAL IMPACT
STATEMENTS
ECOLOGICAL LAND
CLASSIFICATION
WETLAND EVALUATION
VEGETATION ASSESSMENT
BOTANICAL INVENTORIES
WILDLIFE SURVEYS
MONITORING

LANDSCAPE ARCHITECTURE

MASTER PLANNING
RESIDENTIAL COMMUNITIES
COMMERCIAL/INDUSTRIAL
HEALTHCARE AND EDUCATION
STREETSCAPES
PARKS AND OPEN SPACES
TRAIL SYSTEMS
GREEN ROOFS
CONTRACT ADMINISTRATION

EXPERT OPINION

OMB TESTIMONY
LEGAL PROCEEDINGS
PEER REVIEW
RESEARCH
EDUCATION

according to Phase 1 Erin Service and Settlement Master Plan - Environmental Component. The tributaries above and below the dam are classified as cold water tributaries with associated cold water fish communities, whereas the on-line ponds and adjoining sections of tributaries are classified as warm water systems and fish communities. According to the Credit River Fisheries Management Plan, the Hillsburgh Dam is known to have negative fish community impacts through changes to the thermal regimes and imposed barriers to movement (CVC & MNR, 2002).

DRAFT

Proposed Terms of Reference

The ToR, provided below will be based on background natural heritage information (where available) and site visits by Aboud & Associates to collect detailed natural heritage information related to Ecological Land Classification (ELC) communities, flora, fauna, habitat, watercourses and fish. A description of these existing natural heritage features will be detailed. A preliminary assessment will be provided to determine potential impacts and opportunities to natural heritage features from potential design options to address the structural state of the dam.

ToR for the Natural Heritage Existing Conditions Report are listed below.

1. Conduct background screening of relevant documents, material and online mapping sources (e.g. NHIC, CVC, MNR-Guelph District, and Wellington County).
2. Conduct ELC evaluation and prepare ELC community mapping using available background resources, supplemented with 3 season ELC field evaluations and desktop analysis.
3. Complete a 3 season botanical inventory and review of past available inventories to develop a comprehensive list of flora species present. Review and update status of all identified species (SRank; GRank; COSEWIC; COSSARO; Local significance, as listed in Dougan & Associates and Snell & Cecile. 2009).
4. Provincially Significant Wetland:
 - a. Review Wetland Evaluation file to determine presence of potentially significant features.
 - b. Confirm accuracy of current wetland boundaries through desktop analysis and consultation with Ontario Ministry of Natural Resources and Forestry(MNRF).
 - c. (Provisional): Confirm and re-stake wetland boundary of areas that are not current and that may be altered through changes to the dam structure. Work with MNRF to have new boundaries approved.
5. Bird Surveys:
 - a. Complete breeding bird survey of study area, following the protocol of the Breeding Bird Atlas (Bird Studies Canada. 2001). Confirm the presence or absence of Eastern Meadowlark and Bobolink.
 - b. Assess for the presence of the following Significant Wildlife habitat (MNR. 2000):
 - i. Waterfowl Stopover and Staging areas
 - ii. Shorebird migratory stopover area
 - iii. Song bird migratory area
 - iv. Raptor Wintering area

6. Winter Wildlife Survey:

- a. Conduct a survey for signs or sightings of winter wildlife and their associated habitat. Location of observed species will be recorded and mapped.
- b. Assess for the presence of the following Significant Wildlife habitats (MNR. 2000):
 - i. Deer wintering yards
 - ii. Deer Movement Corridors

7. Anuran Survey: Complete three evening anuran (frog and toad) call counts surveys for all potentially suitable habitat locations. Protocols described in the Marsh Monitoring Program will be followed (Marsh Monitoring Program. 2003).

8. Record incidental wildlife observations made during field investigations and combine data with existing wildlife inventories to create a comprehensive wildlife species list. Review and update status of all identified species (SRank; GRank; COSEWIC; COSSARO; Local significance, as listed in Dougan & Associates and Snell & Cecile. 2009).

9. Identify, describe and map wildlife habitat areas and assess for significance using Significant Wildlife Technical Guide (MNR. 2000) and MNR input.

10. Identify specialized habitat or potential SAR habitat for SAR's known to occur in Wellington County. Will be completed using MNR Wellington County SAR and Habitat Requirements Table along with ELC community maps, field investigation and aerial photo interpretation.

11. Fish:

- a. Compile fish community records from MNR and CVC to create a comprehensive fish species list; supplement existing data with additional field sampling if necessary.
- b. Characterise fish habitat within the study area based on thermal regime, vegetation, barriers to movement, depth, pools and riffles, and substrate.

12. Assess the landscape level context of the study area within the broader region, including drainage line, migratory corridors, extended ELC communities, wetlands, and adjacent habitat and wildlife linkages. Specific focus of the landscape level context will be on fish communities and their movement within the West Credit River System.

13. Prepare an interim summary report of existing natural heritage conditions and a preliminary assessment of potential impacts and opportunities to natural heritage features. Detailed project information of species lists, maps, photographs and GIS files will be provided.

Yours truly,

ABOUD & ASSOCIATES INC.



Ryan Hamelin
Terrestrial and Wetland Ecologist

DRAFT

REFERENCES

- Bird Studies Canada. 2001. *Ontario Breeding Bird Atlas Guide for Participants*. Environment Canada, Ministry of Natural Resources, Bird Studies Canada, Federation of Ontario Naturalists, and Ontario Field Ornithologists. 43 pp
- CVC & MNR. 2002. Credit River Fisheries Management Plan. Credit Valley Conservation Authority and Ministry of Natural Resources. 5-36 pp
- Dougan & Associates and Snell & Cecile. 2009. *Guelph Natural Heritage Strategy. Appendix A: Significant Plant List for Wellington County & Appendix B2: List of Significant Wildlife in Wellington County*. Guelph, Ontario.
- Marsh Monitoring Program. 2003 Edition. *Training Kit and Instructions for Surveying Marsh Birds, Amphibians, and Their Habitats*. Published by Bird Studies Canada in cooperation with Environment Canada and the U.S. Environmental Protection Agency. 44 pp
- MNR. 2000. Significant Wildlife Habitat: Technical Guide. Ministry of Natural Resources.

DRAFT

Ryan Hamelin

From: Slaght, Tyler <TSlaght@creditvalleyca.ca>
Sent: December-17-14 2:43 PM
To: 'Chris Clark'; 'rose.whalen@ontario.ca'
Cc: Steven Aboud; Ryan Hamelin; Paul Ziegler
Subject: RE: Hillsburgh Dam and Bridge Class EA - Natural Heritage Component - Project Terms of Reference

Hi Chris,

CVC has reviewed the terms of reference for the natural heritage component and provide the following comments:

1. CVC recommends the limits of the study area on the eastern tributary upstream of the pond be a formal reach break (e.g. road crossing, feature boundary) rather than an arbitrary break.
2. List and describe the natural areas on site, including any natural area designations as defined by CVC, the Town of Erin, Wellington County and/or the Ministry of Natural Resources.
3. Outline relevant federal, provincial, municipal and agency legislation and policies related to the natural area/s and designations that will be applied to options associated with the dam.
4. Please note that while the fish community in the Hillsburgh Pond is characterized by warm water species, the pond is managed as coldwater due to the presence of Brook Trout upstream and downstream of the pond. Mapping in the EA and mitigation measures (e.g. timing windows) should reflect this.
5. Review CVC's available water temperature data (to be provided) and fill data gaps as required. A thermal profile of the pond over as long a period as possible should be considered to assess stratification and the dissolved oxygen profile.
6. Please note that Round Goby, an invasive aquatic species, have been found upstream and downstream of Hillsburgh Dam. The presence and potential spread of this species should be considered in the EA.
7. Breeding bird surveys are to be completed in accordance with the Marsh Monitoring Program (CWS and Bird Studies Canada). That is, two surveys must be conducted at least 10 days apart between late May and July 5th. The surveys must be conducted in either the early morning and/or early evening depending on habitat and potential species present, as per the protocol.
8. In addition to the Significant Wildlife Habitat Technical Guide (MNR 2000), the assessment of Significant Wildlife habitat should follow the MNRF's SWH Ecoregion 6E Criterion Schedule. Based on criteria for Ecoregion 6E, data collected by CVC in 2011 and 2012 indicates that the Hillsburgh Pond is Significant Wildlife Habitat for Waterfowl Stopover and Staging Areas (Aquatic).
9. Surveys for Species at Risk should target all possible Species at Risk based on the presence of suitable habitat, and not just Meadowlark and Bobolink. Based on the habitat features present CVC questions whether surveys for Meadowlark and Bobolink are warranted. Target species may include, but are not limited to: Blanding's Turtle, Snapping Turtle, Least Bittern, Butternut, Chorus Frog, Barn Swallow, and Species-at-Risk bats. MNRF should be contacted for Species at Risk screening.
10. Complete turtle surveys and provide discussion on the suitability of features within the study area for overwintering, nesting and movement habitat.

11. In addition to assessing local rarity based on *Guelph Natural Heritage Strategy* (Dougan & Associates and Snell & Cecile, 2009), GPS the location and describe the distribution of all rare or uncommon species based upon *Vascular Plant Flora of the Region of Peel and the Credit River Watershed* (Kaiser, 2001 and amendments). CVC may request detailed mapping of the species occurrence at a later date.
12. CVC requests an invitation to be present for the staking of the PSW with the MNRF.
13. Identify mitigation measures/restoration opportunities to eliminate and/or minimize negative impacts associated with the preferred option.

Please let me know if you have any questions. Please note I will be out of the office between December 24 returning January 19.

Regards,



Tyler Slaght
 Regulations Officer
 Credit Valley Conservation
tslaght@creditvalleyca.ca | 905.670.1615 ext 406

From: Chris Clark [mailto:cclark@tritoneng.on.ca]
Sent: December 8, 2014 2:40 PM
To: Slaght, Tyler; 'rose.whalen@ontario.ca'
Cc: Steven Aboud; Ryan Hamelin (ryan@aboudtng.com); Paul Ziegler
Subject: Hillsburgh Dam and Bridge Class EA - Natural Heritage Component - Project Terms of Reference

Hi Tyler/Rose,

I have attached the Natural Heritage portion of the project's Terms of Reference for CVC and MNR review and comment. The Fluvial Geomorphology and Hydro technical Terms of Reference will follow under separate cover.

Let us know if you have any questions or require clarification on anything.

Thanks,

Chris Clark, M.A.Sc. E.I.T.



Triton Engineering Services Limited
 105 Queen Street West, Unit 14 Fergus, ON N1M 1S6
 Tel - (519) 843-3920 • Fax - (519) 843-1943 • www.tritoneng.on.ca

The information contained in this Credit Valley Conservation e-mail message is directed in confidence solely to the person(s) named above and may be otherwise disclosed or disseminated without the permission of the sender. The message may contain information that is privileged, confidential and exempt from disclosure under the *Access to Information Act* and the *Privacy Act* and by the *Personal Information Protection and Electronic Document Act*. The use of such personal information is prohibited. If you have received this message in error, please notify the sender immediately by e-mail. If you are not the named addressee you should not disseminate, distribute or copy this e-mail. Please notify the sender immediately by e-mail if you have received this e-mail by mistake. If you are not the named addressee you should not disseminate, distribute or copy this e-mail. Thank you.

Ryan Hamelin

From: Ryan Hamelin
Sent: January-06-15 10:34 AM
To: 'Slaght, Tyler'
Cc: Chris Clark; Steven Aboud; Larry Van Wyck
Subject: RE: Hillsburgh Dam and Bridge Class EA - Natural Heritage Component - Project Terms of Reference

Hello Tyler,

Thank you for your comments on our proposed Terms of Reference (ToR) for the Hillsburgh Dam EA.

Throughout this process we want to be as efficient as possible in our project, and to make sure we are not completing any unnecessary work or analyses. Based on that, is there any proposed actions in our initial ToR that the CVC feels would not need to be included as part of the EA process?

Based on your provided comments there are a few points that we would like some clarification on before finalizing our ToR. I have addressed each of your comments below and where applicable requested additional information or clarification on a few of the points (4,5,7,8,9).

CVC Comments and Aboud & Associates Notes:

1. CVC recommends the limits of the study area on the eastern tributary upstream of the pond be a formal reach break (e.g. road crossing, feature boundary) rather than an arbitrary break.
 - This can be accommodated by moving the study boundary downstream approximately 100 m to Covert Lane in Hillsburgh. The further upstream reaches will still be included in the landscape level analysis already proposed. An updated study area map has been provided to show the new limits of the study area.
2. List and describe the natural areas on site, including any natural area designations as defined by CVC, the Town of Erin, Wellington County and/or the Ministry of Natural Resources.
 - This is already accounted for as part of the background screening outlined in Term 1 of the proposed Terms of Reference. We will re-write the Term to more directly address your comment in the final ToR.
3. Outline relevant federal, provincial, municipal and agency legislation and policies related to the natural areas and designations that will be applied to options associated with the dam.
 - This is already accounted for as part of the background screening outlined in Term 1 of the proposed Terms of Reference. We will re-write the Term to more directly address your comment in the final ToR.
4. Please note that while the fish community in the Hillsburgh Pond is characterized by warm water species, the pond is managed as coldwater due to the presence of Brook Trout upstream and downstream of the pond. Mapping in the EA and mitigation measures (e.g. timing windows) should reflect this.
 - This difference between the actual thermal regime and associated fish species vs. how the Hillsburgh Pond is managed will be noted and may have implications around recommended mitigation measures. Are the two ponds directly downstream from the Hillsburgh Pond also managed in the same way (i.e. Cold water)? Besides timing windows, are there other CVC active fish management decisions or actions associated with cold water management?

5. Review CVC's available water temperature data (to be provided) and fill data gaps as required. A thermal profile of the pond over as long a period as possible should be considered to assess stratification and the dissolved oxygen profile.
 - We have already received temperature Data from CVC that has continuous monitoring from June 3rd to November 15th for 2013. From this data we can assess temperature profiles of the distinct tributary reaches and comment on the seasonal fluctuations. Does CVC have Temperature data for additional years? The data we have already received appears to be a complete and accurate temperature profile for the study area and I would not anticipate collecting additional temperature data.
 - Does CVC already have the data on lake stratification and dissolved oxygen profile, or would this be something that has to be collected? If the data has not been collected could you expand on the expectation of the study as well as the relevance to the Dam EA and how it should be used to assess impacts or determine best options?
6. Please note that Round Goby, an invasive aquatic species, have been found upstream and downstream of Hillsburgh Dam. The presence and potential spread of this species should be considered in the EA.
 - We can specifically assess the potential impact of Round Goby movement and habitat as part of our already proposed background fish screening studies to be completed. Since the CVC and MNRF already have records of the Goby upstream and downstream of the Dam I wouldn't expect any additional sampling to be required.
7. Breeding bird surveys are to be completed in accordance with the Marsh Monitoring Program (CWS and Bird Studies Canada). That is, two surveys must be conducted at least 10 days apart between late May and July 5th. The surveys must be conducted in either the early morning and/or early evening depending on habitat and potential species present, as per the protocol.
 - This was part of our initial ToR, but was under a different protocol reference. The actual study methodology is the same between the ToR and CVC comments and will be completed in accordance to the Marsh Monitoring Program.
 - Part of the data already received from the CVC includes a two visit Breeding Bird Survey Completed by Bob Curry in June and July 2009. Can this data be used to fulfil the Breeding Bird Survey Requirements of the EA, or does a new full Breeding Bird Survey need to be completed? If the CVC survey suffices, could the meta data such as study area maps be provided ?
8. In addition to the Significant Wildlife Habitat Technical Guide (MNR 2000), the assessment of Significant Wildlife habitat should follow the MNRF's SWH Ecoregion 6E Criterion Schedule. Based on criteria for Ecoregion 6E, data collected by CVC in 2011 and 2012 indicates that the Hillsburgh Pond is Significant Wildlife Habitat for Waterfowl Stopover and Staging Areas (Aquatic).
 - The SWH Ecoregion 6E Criterion Schedule will be used in conjunction with the Significant Wildlife Habitat Technical Guide.
 - Can the detailed data from the past CVC SWH studies be provided?
9. Surveys for Species at Risk should target all possible Species at Risk based on the presence of suitable habitat, and not just Meadowlark and Bobolink. Based on the habitat features present CVC questions whether surveys for Meadowlark and Bobolink are warranted. Target species may include, but are not limited to: Blanding's Turtle, Snapping Turtle, Least Bittern, Butternut, Chorus Frog, Barn Swallow, and Species-at-Risk bats. MNRF should be contacted for Species at Risk screening.
 - Our initial Terms of Reference proposed a background Species at Risk Habitat Screening using the Wellington MNRF SAR list. Consultation with MNRF has started regarding their requirements for

Species at Risk surveys, techniques and to identify specific target species. The CVC will be provided a list of proposed species at risk to be surveyed for based on consultation with MNRF.

10. Complete turtle surveys and provide discussion on the suitability of features within the study area for overwintering, nesting and movement habitat.
 - Will be added to our terms of reference. Consultation with MNRF has started regarding requirements and methodology for Turtle surveys.
11. In addition to assessing local rarity based on *Guelph Natural Heritage Strategy* (Dougan & Associates and Snell & Cecile, 2009), GPS the location and describe the distribution of all rare or uncommon species based upon *Vascular Plant Flora of the Region of Peel and the Credit River Watershed* (Kaiser, 2001 and amendments). CVC may request detailed mapping of the species occurrence at a later date.
 - This component will be added to our ELC surveys and plant inventories field surveys and mapped.
12. CVC requests an invitation to be present for the staking of the PSW with the MNRF.
 - Wetland boundary delineation was included as a provisional item in the ToR if the existing wetland boundary was found to be inaccurate and needed to be refined. If boundaries are staked and re-delineated for any portions of the study area CVC would be informed and invited to participate in the approval of the new boundary delineation. Based on preliminary interpretation of the current wetland boundary and ortho images it appears the delineated boundary may be a reasonably accurate representation of the actual wetland feature. It is our opinion that the wetland boundary would only need to be re-delineated if found to be inaccurate and unsuitable for identifying preferred EA options.
 - Could you please comment as to CVC's position regarding if portions of the wetland boundary need to be re-delineated as part of the existing features study, or if the 2005 MNRF updated boundary should suffice.
13. Identify mitigation measures/restoration opportunities to eliminate and/or minimize negative impacts associated with the preferred option.
 - As part of the EA, opportunities for mitigation / restoration of the preferred options will be provided.

It should also be noted that the majority of the study area is on private property and access has not yet been granted for large sections of the Natural Heritage study area. Communication with landowners is ongoing and permission to access properties has been requested. However, it is still likely that much of the proposed study area will not be accessible. In these instances, where access to portions of the study area are not granted, alternative study methods such as observation from adjacent lands, orthophotography analysis, and background materials will be used to characterise the existing conditions of the property.

Thank you Tyler for reviewing the ToR and providing detailed comments. Perhaps it would be best to have a phone conversation to clarify the above points and to better understand the CVC's positions. If you could let me know if there is a time we could talk that would be appreciated.

We look forward to hearing back from you.

Ryan Hamelin

Ryan Hamelin, B.S.c (Env). M.Sc. Terrestrial and Wetland Ecologist

Ryan Hamelin

From: Slaght, Tyler <TSlaght@creditvalleyca.ca>
Sent: January-23-15 10:13 AM
To: Ryan Hamelin
Cc: 'Chris Clark'
Subject: RE: Hillsburgh Dam and Bridge Class EA - Natural Heritage Component - Project Terms of Reference

Hi Ryan,

We've put together some responses to your questions, see below. If have you any further questions, perhaps our ecology staff can chat with you, it probably makes more sense for something like this. Let me know which points you still have questions about and I'll arrange to have them call you.

#4. The 2 ponds downstream of the Station Street pond are also managed the same way (i.e. warmwater species present but managed for the coldwater species that are up and downstream). The only other "formal" management action that was in the Credit River Fisheries Management Plan would be requesting a 30m buffer rather than a 15m buffer.

#5. Continuous Temperature logging data for six sites in the summer of 2014 is available. CVC has no data on water temperatures or dissolved oxygen levels in the pond. This would be useful for assessing the existing impacts of the pond (e.g. does it stratify, do anoxic conditions exist) and benefits of some mitigation options (e.g. installation of a bottom draw and determining discharge volumes).

#6. No additional surveys for Round Goby are needed.

#7: Additional surveys are required as Bob Curry's surveys were completed over a smaller study area and did not include the pond. The meta data can be provided.

#8: As above, no concerns with providing the data.

#12: As indicated in the ToR, CVC is of the understanding that staking of the wetland boundary is provisional and dependent upon consultation with MNRF. If MNRF determines that the PSW boundary requires staking, CVC requests an invitation to be present.

For the additional information above we will put that together and send it to you as soon as possible.

Regards,



Tyler Slaght
Regulations Officer
Credit Valley Conservation
tslaght@creditvalleyca.ca | 905.670.1615 ext 406

From: Ryan Hamelin [<mailto:ryan@aboutdng.com>]
Sent: January 21, 2015 11:01 AM
To: Slaght, Tyler

Guelph District
1 Stone Road West
Guelph, Ontario
N1G 4Y2

Telephone: (519) 826-4955
Facsimile: (519) 826-4929

January 26, 2015

Ryan Hamelin
Aboud & Associates Inc.
591 Woolwich Street
Guelph, ON
N1H 3Y5

Dear Mr. Hamelin

Re: Terms of Reference for Hillsburgh Dam, Natural Heritage Existing Conditions Report
As Part of the Municipal Class Environmental Assessment

The Ministry of Natural Resources and Forestry Guelph District Office (MNRFO) had a chance to review the attached terms of reference (ToR) for the Hillsburgh dam regarding the existing natural heritage conditions report as part of the Municipal Class Environmental Assessment and offer the following comments:

- If the order in which the surveys/screening presented in the ToR is in chronological order, it may be beneficial for the survey to identify specialized habitat or potential SAR habitat for SAR to occur immediately after the Ecological Land Classification (ELC) evaluation.
This may inform the types of surveys required for the subject properties.
- Some of the described surveys may be limited due to property access. How does Aboud & Associates plan to resolve this?
- A proposed clarification in the Background and Context section, regarding the comment "whereas the on-line ponds and adjoining sections of tributaries are classified as warm water systems and fish communities". This should be amended to reflect that the stream sections between the Hillsburgh and Ainsworth pond and between Ainsworth and Rudd pond have coldwater fish communities.
- Regarding 5 a. Bird surveys, confirm the presence of other SAR along with Eastern Meadowlark and Bobolink.
- Regarding 6 b. i. Deer wintering yards are referred to as Deer Winter Congregation Areas in this area.

Thank you for giving us the opportunity to review. If you have any questions, please do not hesitate to contact me

Sincerely,



Rose Whalen
T: 519-826-4910



591 Woolwich Street
Guelph . Ontario
N1H 3Y5

T: 519.822.6839
F: 519.822.4052
info@aboutdng.com
www.aboutdng.com

URBAN FORESTRY
ARBORIST REPORTS
MANAGEMENT PLANS
TREE PRESERVATION PLANS
TREE RISK ASSESSMENT
GIS TREE INVENTORIES
TREE APPRAISALS
MONITORING

ECOLOGICAL RESTORATION
NATURAL SYSTEMS DESIGN
HABITAT RESTORATION
EDGE MANAGEMENT PLANS
RAVINE STEWARDSHIP PLANS
NATURALIZATION PLANS
INTERPRETIVE DESIGN
MONITORING
CONTRACT ADMINISTRATION

ENVIRONMENTAL STUDIES
SUBWATERSHED STUDIES
ENVIRONMENTAL IMPACT
STATEMENTS
ECOLOGICAL LAND
CLASSIFICATION
WETLAND EVALUATION
VEGETATION ASSESSMENT
BOTANICAL INVENTORIES
WILDLIFE SURVEYS
MONITORING

LANDSCAPE ARCHITECTURE
MASTER PLANNING
RESIDENTIAL COMMUNITIES
COMMERCIAL/INDUSTRIAL
HEALTHCARE AND EDUCATION
STREETSCAPES
PARKS AND OPEN SPACES
TRAIL SYSTEMS
GREEN ROOFS
CONTRACT ADMINISTRATION

EXPERT OPINION
OMB TESTIMONY
LEGAL PROCEEDINGS
PEER REVIEW
RESEARCH
EDUCATION

April 10, 2015

Our Project No: AA12-137A
Sent by email: cclark@trotoneng.on.ca

Tyler Slaght
Credit Valley Conservation
1255 Old Derry Road
Mississauga, Ontario
L5N 6R4

c/o:

Chris Clark
Triton Engineering Services Limited
105 Queen Street West, Unit 14
Fergus, Ontario
N1M 1S6

Re: Proposed Targeted Species at Risk Survey

Dear Tyler,

As requested, Aboud & Associates Inc. have undertaken a review of all Species at Risk (SAR) which may occur in the project location using the Wellington County MNR species at risk list.

The following targeted surveys for SAR are proposed for the Hillsburgh dam, existing conditions report. Species which were not considered likely in the project location are discussed in brief, following the recommended surveys, and the specifics of their exclusion. The accompanied summary table includes proposed surveys for all SAR within Wellington County, including SAR surveys which follow general survey protocols (e.g. Breeding Bird Protocol).

Jefferson Salamander Surveys

Likelihood of occurrence: Possible, populations located north-east of project location in Orangeville area and east of project location, south of Caledon.

Proposed field work:

1. In 2015, Visual surveys for *Ambystoma* egg masses in candidate ponds identified during initial site visit will be inspected in early April by a qualified wildlife ecologist to determine the presence or absence of any *Ambystoma* species occurring in the project location. Jefferson Salamanders are one of three *Ambystoma* species in Ontario, these survey will help to determine the possible presence of Jefferson Salamander within the Study area. Site visit timings will occur within less than 15 days of approximate salamander movement windows, in order to ensure salamander egg hatches have not yet occurred.

Survey Methods: Visual inspection of any candidate pools will be performed on sunny cloudless days in April, using polarized lenses, with no entry into candidate pools. All egg masses will be identified based on characteristics as frog, toad or salamander, with no effort to determine salamander species in order to avoid disturbance of egg masses and entry into ponds.

2. Should *Ambystoma* egg masses occur in candidate habitat during 2015 visual surveys, application for permits and subsequent field planning for salamander trapping surveys would be expected to occur in late March-early April 2016, after acquiring all permits and training personnel. Survey methods will follow the Jefferson Salamander sampling protocol as provided by the Guelph MNR (2013).

Bat Maternity Roost Surveys

Likelihood of occurrence: probable, all three species are found throughout Ontario.

Target Species: Eastern small-footed Myotis (*Myotis Leibii*), Little Brown Myotis (*Myotis lucifugus*), and Northern Myotis (*Myotis septentrionalis*)

Proposed Desktop work:

1. Identification of all ELC communities (FOD, FOM, FOC, SWD, SWM, SWC) which may be considered candidate bat maternity habitat, following guidelines provided in the bat and bat habitat: guidelines for wind projects (2011), will be treated as confirmed habitat and appropriate mitigation will be applied as outlined below. This proposed methodology is based on communication with Guelph District MNR, which “only recommend surveys if there is potential for impacts to the hibernation or roost habitat.” (pers. comm. Graham. Buck 2015)

Mitigation recommendations- tree removal must occur outside bat maternity season, from September-April, in all habitats considered candidate bat maternity habitat based on ELC results.

Turtle Basking Surveys

Likelihood of occurrence: Blanding's turtle-Possible, populations occur in the vicinity of Guelph and Luther Marsh. Snapping turtle-Probable, populations occur throughout southern Ontario. Spotted turtle-unlikely, populations of spotted turtle are generally found in the vicinity of Georgian Bay and along the Lake Erie shoreline.

Target species: Blanding's turtle (*Emydonidea Blandingii*), snapping turtle (*chelydra serpentine*), spotted turtle (*Clemmys guttata*)

Proposed field work:

A total of 5 Basking surveys in all candidate habitats within the project location will be conducted in 2015 following the MNR Guelph district Blanding's survey protocol (2012). Basking surveys, including overwintering (late march-early April) and summer habitat (late April-June 15), will be conducted in all waterbodies and wetlands.

Methods: All shorelines and potential basking sites in the project location will be surveyed from the sunlit side using high power binoculars or a spotting scope. If shorelines are obstructed by vegetation, surveys will be conducted from canoe or while wearing waders in water as required; provided that access is granted. Between late March and early May, surveys will be conducted between 9am and 5pm. between late May and early June turtles are less reliably found late in the day, as a result surveys will occur between 9am and 12pm. When temperatures fall between 6c and 10c, surveys may only occur on sunny days with no wind between 10am and 5pm, at full sunlight basking sites. When temperatures fall between 10c and 25c, surveys will be conducted between 9am and noon on sunny days.

Snake Visual Encounter and Active Hand Search Surveys

Target Species: Eastern ribbonsnake (*Thamnophis sauritus*), milksnake (*Lampropeltis Triangulum*)

Proposed field work:

Visual encounter and active hand search surveys will occur from late April through late June in all candidate habitats identified during initial ELC screening and site visit. A minimum of 3 surveys, two weeks apart, searching all suitable habitats and flipping any natural or naturalized cover, will occur in all suitable habitat identified in the project location.

Methods: surveys will occur on sunny days when air temperatures are between 8c and 25c, and on overcast day's air temperatures must be above 15c. Surveys will follow pre-determined transects, traversing all areas of suitable habitat for both eastern ribbonsnake and milksnake.

West Virginia White Visual Survey

Likelihood of occurrence: possible, species host plant occurs in the project location.

Proposed field work:

Visual surveys for adults and caterpillars will occur within moist, deciduous woodlands in areas where two-leaved toothwort has been previously identified by the CVC. Surveys will be conducted during spring botanical surveys. Caterpillars feed on the two-leaved toothwort which blooms from April to June. Caterpillars will be looked for carefully on the host plant.

Species that are unlikely to occur in the project location for which targeted surveys exist:

Barn Owl- No habitat is present within the project location, barn owl have not been identified as occurring in the vicinity of the project location. During the second breeding bird Atlas, a single Barn owl was identified in Wellington County with no confirmation on breeding status. This species is unlikely to occur in the project location.

Bobolink and Eastern meadowlark- these grassland bird species are unlikely to occur in the project location, no grassland habitat, pasture or fallow fields were identified through air photo interpretation or initial site visits. Presence/absence will be confirmed through Breeding bird Surveys.

Nightjar survey (Common nighthawk and Whip-poor-will) – habitat for these species was not identified in the project location based on air photo interpretation and initial site visit. As a result, no additional targeted surveys are recommended.

Least bittern – No suitable habitat was identified in the project location. Targeted surveys are not recommended. General Marsh monitoring playback surveys for marsh birds will occur in appropriate habitat in the project location.

Short-eared owl - No suitable habitat was identified in the project location. Targeted surveys are not recommended.

Fish Species at Risk- Black redhorse, Redside Dace and Silver Shiner were not documented in past fish surveys conducted by MNRF or CVC. Ideal habitat is not present. No surveys to be conducted.

Rusty-patched bumble- Not documented in project location. No suitable habitat was identified in the project location. Discussion with Graham Buck at the MNRF indicates that there is no requirement to complete targeted surveys for this species in the project location, if a bee is identified as suspect, photos and UTM will be recorded during botanical surveys.

Mollusc Species at Risk –Rainbow mussel and Wavy-rayed lampmussel have not been identified in the Upper Credit River, Ideal habitat is not present in project location, not detected during previous aquatic sampling.

Butler's gartersnake- Ideal habitat for this species is unlikely to occur in the project location based on air photo interpretation and initial site visit. Butler's gartersnake occur in fragmented populations in Ontario, the nearest population is located in Luther marsh, which is ~30km from the project location. One home range study in Michigan indicated that Butler's gartersnake occupy a very small home range, with a maximum distance of 300m. It is unlikely that Butler's gartersnake would occur in the Project location as a result of the distance to the nearest known population.

Massasauga rattlesnake- Ideal habitat is unlikely to occur in the project location. This species is only known to occur historically in Wellington County; as a result, it is unlikely to occur in the project location.

Yours truly,

ABOUD & ASSOCIATES INC.



Cheryl-Anne Ross, B.Sc., Wildlife Ecologist



Ryan Hamelin, M.Sc, Terrestrial and Wetland Ecologist

cc. P. Ziegler, Triton Engineering Services Ltd
C. Clark, Triton Engineering Services Ltd
R. Whalen, Ministry of Natural Resources and Forestry

WELLINGTON - Upper Tier - MNRF SAR List			
Species At Risk Designations			
ENDANGERED			
THREATENED			
SPECIAL CONCERN			
EXTIRPATED			
AMPHIBIANS			
		ESA Protection	Proposed Survey Action
Jefferson Salamander (<i>Ambystoma jeffersonianum</i>)	Known to Occur	Species Protection and Habitat Regulation	Targeted Survey - Egg Survey
BIRDS			
		ESA Protection	Proposed Survey Action
Acadian Flycatcher (<i>Empidonax virescens</i>)	Suspected to Occur	Species and General Habitat Protection	Breeding Bird Survey
Bald Eagle (<i>Haliaeetus leucocephalus</i>)	Known to Occur	N/A	Breeding Bird Survey
Bank Swallow (<i>Riparia riparia</i>)	Known to Occur	Species and General Habitat Protection June 27, 2014	Breeding Bird Survey
Barn Owl (<i>Tyto alba</i>)	Known to Occur	Species Protection and Habitat Regulation	Breeding Bird Survey - No targeted night survey based on lack of suitable habitat
Barn Swallow (<i>Hirundo rustica</i>)	Known to Occur	Species and General Habitat Protection	Breeding Bird Survey
Black Tern (<i>Chidonias niger</i>)	Known to Occur	N/A	Breeding Bird Survey
Bobolink (<i>Dolichonyx oryzivorus</i>)	Known to Occur	Species and General Habitat Protection	Breeding Bird Survey
Canada Warbler (<i>Cardellina canadensis</i>)	Suspected to Occur	N/A	Breeding Bird Survey
Cerulean Warbler (<i>Setophaga cerulea</i>)	Known to Occur	Species and General Habitat Protection	Breeding Bird Survey
Chimney Swift (<i>Chaetura pelagica</i>)	Known to Occur	Species and General Habitat Protection	Breeding Bird Survey
Common Nighthawk (<i>Chordeiles minor</i>)	Known to Occur	N/A	Breeding Bird Survey- No additional targeted survey based on lack of appropriate habitat.
Eastern Meadowlark (<i>Sturnella Magna</i>)	Known to Occur	Species and General Habitat Protection	Breeding Bird Survey - Three survey days to confirm absence
Eastern Wood-Pewee (<i>Contopus virens</i>)	Known to Occur	N/A	Breeding Bird Survey
Eastern Whip-poor-will (<i>Caprimulgus vociferus</i>)	Known to Occur	Species and General Habitat Protection	Breeding Bird Survey- No additional survey based on lack of appropriate habitat.
Golden-winged Warbler (<i>Vermivora chrysoptera</i>)	Known to Occur	N/A	Breeding Bird Survey
Henslow's Sparrow (<i>Ammodramus henslowii</i>)	Known to Occur	Species and General Habitat Protection	Breeding Bird Survey
Least Bittern (<i>Ixobrychus exilis</i>)	Known to Occur	Species and General Habitat Protection	Breeding Bird Survey- No additional survey based on lack of appropriate habitat.
Loggerhead Shrike (<i>Lanius ludovicianus</i>)	Historically Known to Occur	Species and General Habitat Protection	Breeding Bird Survey

BIRDS		ESA Protection	Proposed Survey Action
Louisiana Waterthrush (<i>Seiurus motacilla</i>)	Suspected to Occur	N/A	Breeding Bird Survey
Northern Bobwhite (<i>Colinus virginianus</i>)	Known to Occur	Species and General Habitat Protection	Breeding Bird Survey
Olive-sided Flycatcher (<i>Contopus cooperi</i>)	Suspected to Occur	N/A	Breeding Bird Survey
Red-Headed Woodpecker (<i>Melanerpes erythrocephalus</i>)	Known to Occur	N/A	Breeding Bird Survey
Short-eared Owl (<i>Asio flammeus</i>)	Known to Occur	N/A	Breeding Bird Survey - No targeted night survey based on lack of habitat
Wood Thrush (<i>Hylocichla mustelina</i>)	Known to Occur	N/A	Breeding Bird Survey
Yellow-breasted Chat (<i>Icteria virens</i>)	Historically Known to Occur	Species and General Habitat Protection	Breeding Bird Survey
FISH		ESA Protection	Proposed Survey Action
Black Redhorse (<i>Moxostoma duquesnei</i>)	Known to Occur	Species and General Habitat Protection	Not identified in past MNRF or CVC sampling. No targeted survey to be conducted.
Redside Dace (<i>Clinostomus elongatus</i>)	Known to Occur	Species Protection and Habitat Regulation	Not identified in past MNRF or CVC sampling. No targeted survey to be conducted.
Silver Shiner (<i>Notropis photogenis</i>)	Known to Occur	Species and General Habitat Protection	Not identified in past MNRF or CVC sampling. No targeted survey to be conducted.
INSECTS		ESA Protection	Proposed Survey Action
Monarch Butterfly (<i>Danaus plexippus</i>)	Known to Occur	N/A	Survey following MNRF survey protocol
Rusty-patched Bumble Bee (<i>Bombus affinis</i>)	Formerly Occurred and May Still Occur	Species and General Habitat Protection	Incidental observation during plant surveys. (pers. Comm. Graham Buck 2015)
West Virginia White (<i>Pieris virginianensis</i>)	Known to Occur	N/A	Survey following MNRF survey protocol
MAMMALS		ESA Protection	Proposed Survey Action
Eastern Small-footed Myotis (<i>Myotis leibii</i>)	Suspected to Occur	Species and General Habitat Protection as of June 27, 2014	Desktop habitat identification, following Guideline for Wind Projects (2011)
Grey Fox (<i>Urocyon cinereoargenteus</i>)	Known to Occur	Species and General Habitat Protection	Winter Wildlife survey and incidental wildlife. No targeted survey based on habitat and past observations.
Little Brown Myotis (<i>Myotis lucifugus</i>)	Known to Occur	Species and General Habitat Protection	Desktop habitat identification, following Guideline for Wind Projects (2011)
Northern Myotis (<i>Myotis septentrionalis</i>)	Known to Occur	Species and General Habitat Protection	Desktop habitat identification, following Guideline for Wind Projects (2011)
MOLLUSCS		ESA Protection	Proposed Survey Action
Rainbow Mussel (<i>Villosa iris</i>)	Known to Occur	Species and General Habitat Protection	No Targeted Survey - Not identified in the Upper Credit River
Wavy-rayed lampmussel (<i>Lampsilis fasciola</i>)	Known to Occur	Species and General Habitat Protection	No Targeted Survey - Not identified in the Upper Credit River

PLANTS		ESA Protection	Proposed Survey Action
American Chestnut (<i>Castanea dentata</i>)	Known to Occur	Species and General Habitat Protection	Plant Inventory
American Ginseng (<i>Panax quinquefolius</i>)	Known to Occur	Species and General Habitat Protection	Plant Inventory
Butternut (<i>Juglans cinerea</i>)	Known to Occur	Species and General Habitat Protection	Plant Inventory
Hill's Pondweed (<i>Potamogeton hillii</i>)	Known to Occur	N/A	Plant Inventory
REPTILES		ESA Protection	Proposed Survey Action
Blanding's Turtle (<i>Emydonidea blandingii</i>)	Known to Occur	Species and General Habitat Protection	Turtle Survey - Blanding's Turtle Protocol
Butler's Gartersnake (<i>Thamnophis butleri</i>)	Known to Occur	Species and General Habitat Protection	No Targeted Survey - Unlikely to occur in project study area based on habitat
Eastern Ribbonsnake (<i>Thamnophis sauritus</i>)	Known to Occur	N/A	Visual Encounter and Active Hand Search Surveys
Massassauga Rattlesnake (<i>Sistrurus catenatus</i>)	Historically Known to Occur	Species and General Habitat Protection	No Targeted Survey - Unlikely to occur in project study area based on habitat
Milksnake (<i>Lampropeltis triangulum</i>)	Known to Occur	N/A	Visual Encounter and Active Hand Search Surveys
Snapping Turtle (<i>Chelydra serpentina</i>)	Known to Occur	N/A	Turtle Survey - Blanding's Turtle Protocol
Spotted Turtle (<i>Clemmys guttata</i>)	Known to Occur	Species and General Habitat Protection	Turtle Survey - Blanding's Turtle Protocol

DRY

Ryan Hamelin

From: Slaght, Tyler <TSlaght@creditvalleyca.ca>
Sent: April-13-15 1:06 PM
To: Ryan Hamelin
Cc: Cheryl-Anne Ross; Chris Clark; Paul Ziegler; Whalen, Rose (MNRF)
Subject: RE: Hillsburgh Dam and Bridge Class EA - Natural Heritage Component - Project Terms of Reference

Hi Ryan,

This looks good to us.

Regards,



Tyler Slaght
Regulations Officer
Credit Valley Conservation
tslaght@creditvalleyca.ca | 905.670.1615 ext 406

From: Ryan Hamelin [mailto:ryan@aboudtng.com]
Sent: April 10, 2015 2:41 PM
To: Slaght, Tyler
Cc: Cheryl-Anne Ross; Chris Clark; Paul Ziegler; Whalen, Rose (MNRF)
Subject: FW: Hillsburgh Dam and Bridge Class EA - Natural Heritage Component - Project Terms of Reference

Hi Tyler,

I hope you are doing well.

In response to Comment 9. of CVC's earlier Terms of Reference review, Aboud & Associates have completed a SAR Targeted Survey proposal for all SAR possibly present within the Hillsburgh Dam Study Area, based on available habitat. A SAR habitat assessment was completed based on winter field observations, background resources and orthophotography interpretation. A proposed list of possible SAR was circulated to MNRF Guelph District for comments and recommendation on survey protocol.

The attached letter details our proposed survey protocol for specie specific targeted surveys or desk top analysis's. The accompanying table outlines the proposed action for all Wellington County SAR. Please let us know if you have any comments or recommendation regarding our proposed SAR Survey approach.

Also, I would like to introduce you to Aboud & Associates newest staff Member, Cheryl-Anne Ross. Cheryl-Anne is our new Wildlife Ecologist and will be leading the wildlife portion of the Hillsburgh Dam EA.

Thanks,

Ryan Hamelin

APPENDIX 4. SURVEY DETAILS

Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

SURVEY	DATE	TIME	OBSERVER(S)	TEMP	WIND	CLOUD COVER	PRECIPITATION	PRECIPITATION - PAST 24h
Winter Wildlife	25/02/2015	10:00-16:35	R. Hamelin, M. Iles	-10	2	20	none	snow-1cm
Anuran	15/04/2015	20:31-22:04	C.A. Ross, R.Hamelin	7	1	10	none	none
Salamander Egg Mass Survey	15/04/2015	13:30-17:45	C.A., Ross, R. Hamelin	12	2	5	none	none
Snake Basking	15/04/2015	13:30-17:45	C.A., Ross, R. Hamelin	12	2	5	none	none
Wildlife Habitat Assessment-spring	15/04/2015	13:30-17:45	C.A., Ross, R. Hamelin	12	2	5	none	none
Snake Basking	29/04/2015	9:00-13:00	C.A. Ross, R. Hamelin	10	2	15	none	none
Turtle Basking	29/04/2015	9:25-11:30	C.A. Ross, R.Hamelin	10	1	10	none	none
Turtle Basking	08/05/2015	9:20-11:00	C.A. Ross	19	1	10	none	none
ELC Spring	13/05/2015	9:30 -16:45	R. Hamelin	8	3		none	rain
Snake Basking	14/05/2015	11:30- 12:30	R. Hamelin	16	2	10	none	none
ELC Spring	14/05/2015	12:30- 17:00	R. Hamelin	17	2		none	none
Turtle Basking	15/05/2015	10:10-11:31	R. Hamelin	11	2	10	none	none
ELC Spring	22/05/2015	9:00-16:45	R. Hamelin	12	3		none	none
Anuran	28/05/2015	21:25-22:55	C.A. Ross, R.Hamelin	19	1	0	none	none
Turtle Basking	28/05/2015	10:27-11:45	C.A. Ross	18	2	0	none	rain
Breeding Bird	11/06/2015	7:00-10:53	C.A. Ross	17	3	10	none	rain
Marsh Breeding Birds	11/06/2015	7:00-11:01	C.A. Ross	17	3	10	none	rain
Turtle Basking	11/06/2015	9:10-10:47	R. Hamelin	17	2	20	none	rain
ELC Spring	11/06/2015	12:30- 16:30	R. Hamelin	22	2		rain	rain
Anuran	24/06/2015	21:37-22:54	C.A. Ross, R.Hamelin	16	1	0	none	none
Breeding Bird	09/07/2015	6:22-10:23	C.A. Ross	14	0	80	none	none
Marsh Breeding Birds	09/07/2015	6:22-10:30	C.A. Ross	14	0	80	none	none
ELC Summer	30/07/2015	9:00-16:45	R. Hamelin	27	2		none	none
Shorebird Habitat Assessment	05/08/2015	9:30-10:15	C.A. Ross	17	3	10	none	none
Shorebird Survey	05/08/2015	9:30-10:15	C.A. Ross	17	3	10	none	none
Wetland Boundary Verification	05/08/2015	9:00-14:45	R. Hamelin	23	2		none	rain
ELC Summer	05/08/2015	10:00- 17:00	R. Hamelin	23	2		none	rain
ELC Summer	10/08/2015	9:00-14:45	R. Hamelin	22	1		rain	none
ELC Fall	24/09/2015	10:00- 17:00	R. Hamelin	24	1		none	none
ELC Fall	25/09/2015	9:30 -16:45	R. Hamelin	22	2		none	none
Songbird Migration	08/10/2015	7:20-9:40	C.A. Ross	5	1	0	none	none
Wildlife Habitat Assessment-fall	08/10/2015	9:40-11:00	C.A. Ross	5	1	0	none	none
Aquatic Habitat Assesment	19/10/2015	9:00-14:00	R. Hamelin	13.5	2	5	rain	none

ELC Code	Map ID	Vegetation Type	Community Description
<i>Mixed Meadow (MEM)</i>			
MEMM3	12	Dry - Fresh Mixed Meadow Ecosite	This community is of cultural influence with evidence of past disturbance and clearing. The community is dominated by a mixture of Fringed Brome (<i>Bromus ciliates</i>), Common Milk Weed (<i>Asclepias syriaca</i>), Tall Goldenrod (<i>Solidago altissima</i>), Canada Goldenrod (<i>Solidago canadensis</i>) and a variety of Aster species. A mixture of small shrub and trees are present along the edge of the community where vegetation clearing has not occurred recently.
<i>Coniferous Forest (FOC)</i>			
FOCM2-2	5	Dry-Fresh White Cedar Coniferous Forest	This community is composed almost entirely of Eastern White Cedar (<i>Thuja occidentalis</i>) with little to no understory or ground cover. The Cedar community is dense with some individuals in poor condition due to overcrowding. Soil is a well-drained mineral soil on a moderate slope.
FOCM6	27	Naturalized Coniferous Plantation	Access was not available for this community. ELC is based on observations from a distance and through air photo interpretation. The community is a Naturalized Coniferous Plantation containing approximately equal amounts of mature Norway Spruce (<i>Acer platanoides</i>), White Spruce (<i>Picea glauca</i>), Eastern White Pine (<i>Eastern White Pine</i>) and White Cedar. Understory and ground cover communities are unknown, as well as soil and moisture properties.
FOCM6	6	Naturalized Coniferous Plantation	<p>This community is of cultural influences, with evidence of past disturbance and plantings. The canopy is comprised of a mixture of primarily planted species with some volunteer establishment of native tree species from surrounding communities. Dominant canopy species include; European Larch (<i>Larix decidua</i>) and White Pine, with White Ash (<i>Fraxinus americana</i>), Red Oak (<i>Quercus rubra</i>), Sugar Maple (<i>Acer saccharum ssp. saccharum</i>), Black Cherry (<i>Prunus serotina</i>), Balsam Fir (<i>Abies balsamea</i>) and White Cedar associates. Sub-canopy and understory is dense in areas, with Choke Cherry (<i>Prunus virginiana</i>), Domestic Apple (<i>Malus pumila</i>), European Buckthorn (<i>Rhamnus cathartica</i>), and Tartarian Honeysuckle (<i>Lonicera tatarica</i>).</p> <p>The Community contains an inclusion of Dry-Fresh Forb Meadow Ecosite (MEFM1) composed primarily of Tall Goldenrod, Canada Goldenrod, New England Aster (<i>Symphotrichum novae-angliae</i>) and White Heath Aster (<i>Symphotrichum ericoides var. ericoides</i>).</p>
<i>Mixed Forest (FOM)</i>			
FOMM7-2	23	Fresh - Moist White Cedar - Hardwood Mixed Forest	<p>This narrow community occurs between the Elora-Cataract Trail way and residential properties. Co-dominant species within the community are Eastern White Cedar and White Birch (<i>Betula papyrifera</i>); other canopy tree species include Balsam Poplar (<i>Populus balsamifera</i>), Trembling Aspen (<i>Populus tremuloides</i>), Domestic Apple, American Elm (<i>Ulmus americana</i>), Black Cherry, Sugar Maple and White Ash. The understory contains a variety of woody species, with Alternate-leaf Dogwood (<i>Cornus alternifolia</i>) and White Ash saplings as the most abundant species. Ground layer varies inversely with the abundance of White Cedar and includes Red Trillium (<i>Trillium erectum</i>), Smooth Yellow Violet (<i>Viola macloskeyi</i>), Labrador Violet (<i>Viola labradorica</i>), Wild Leek (<i>Allium tricoccum var. tricoccum</i>) and Sensitive Fern (<i>Onoclea sensibilis</i>).</p> <p>Canopy composition varies and includes a complex of Fresh - Moist White Cedar Coniferous Forest Type, that is almost entirely White Cedar, with little understory of ground cover.</p>
<i>Deciduous Forest (FOD)</i>			
FODM5-8	4	Dry-Fresh Sugar Maple - White Ash Deciduous Forest	This large upland community is dominated by Sugar Maple, with White Ash as sub-dominant. Associates include; Black Cherry, Red Oak, White Cedar, Balsam Fir, American Elm, Trembling Aspen, Eastern Hop-hornbeam (<i>Ostrya virginiana</i>), Basswood (<i>Tilia americana</i>) and Birch species. The community is mature containing a number of large trees with a DBH of 50 cm or greater. Understory and ground layer cover is diverse, and includes; Yellow Trout-lily (<i>Erythronium americanum</i>), Smooth Yellow Violet (<i>Viola pubescens var. scabriuscul</i>), Blood Root (<i>Sanguinaria canadensis</i>), Red Baneberry (<i>Actaea rubra</i>), and Giant Blue Cohosh (<i>Caulophyllum giganteum</i>), as well as a number of upland grass and sedge species.
FODM6	16	Fresh - Moist Sugar Maple Deciduous Forest Ecosite	This long narrow community borders residential properties and has a high edge to area ratio. Cultural influences include plantings within the community and property maintenance along the edge of the community. Sugar Maple is the dominant canopy species in the community with an abundance of Manitoba Maple (<i>Acer negundo</i>) and Green Ash (<i>Fraxinus pennsylvanica</i>) in the wetter areas. Other canopy species include Black Walnut (<i>Juglans nigra</i>), Balsam Poplar, White Ash and planted White Pine, European Alder (<i>Alnus glutinosa</i>), and Freeman's Maple (<i>Acer x freemanii</i>). The sub-canopy and understory is dense with immature canopy species, and a variety of native and non-native shrub and herbaceous species. Low trees and shrubs are often covered with Riverbank Grape (<i>Vitis riparia</i>) and Wild Mock-cucumber (<i>Echinocystis lobata</i>).
FODM7-7	30	Fresh - Moist Manitoba Maple Lowland Deciduous Forest	This culturally influenced riverine community occurs along the tributary within the town of Hillsburgh, upstream of the main pond. The canopy is dominated by Manitoba Maple and Crack Willow (<i>Salix fragilis</i>), with occurrences of American Elm, Sugar Maple, and Black Cherry, White Willow (<i>Salix alba</i>), Black Walnut and Scots Pine (<i>Pinus sylvestris</i>). The understory is a mix of native and non-native shrub and herbaceous species. Soil moisture varies with proximity from the watercourse, with more wetland characteristics directly adjacent to the tributary.

ELC Code	Map ID	Vegetation Type	Community Description
FODM8-1	25	Fresh - Moist Poplar Deciduous Forest	This upland community is located between the Elora-Cataract Trail way and agricultural lands. The co-dominant tree species in the community are Trembling Aspen and Sugar Maple, the two trees species are generally separated within the community, with the Sugar Maple dominant along the trail edge. Associate canopy tree species include Bass Wood, White Ash, and White Cedar. The understory includes young White Ash along with Alternate-leaf Dogwood, Wild Red Raspberry (<i>Rubus idaeus</i> ssp. <i>strigosus</i>) and Staghorn Sumac (<i>Rhus typhina</i>) along the community edge.
FODM8-1	15	Fresh - Moist Poplar Deciduous Forest	This is a narrow culturally influenced community behind residential properties. The community is dominated by Trembling Aspen and Balsam Poplar with occurrences of Norway Maple, Manitoba Maple, Black Cherry, and White Cedar. Understory species are a mixture of native species and exotic weedy species such as Goutweed (<i>Aegopodium podagraria</i>) and Colt's Foot (<i>Tussilago farfara</i>). Soil is mineral with moisture varying from Fresh to Moist.
<i>Coniferous Swamp (SWC)</i>			
SWCM1-2	17	White Cedar - Conifer Mineral Coniferous Swamp	This community surrounds the Ainsworth Pond and the lands directly south of the Pond. The community is dominated by Eastern White Cedar, with occurrences of a variety of other deciduous and coniferous species as minor canopy components. The understory is a diverse variety of mostly native plants, which includes Downy Serviceberry (<i>Amelanchier arborea</i>) and Alderleaf buckthorn (<i>Rhamnus alnifolia</i>), with Dwarf Scouring Rush (<i>Equisetum scirpoides</i>) among the ground layer species. Surface water and organic soil is present throughout much of the community with some areas of dry mineral soil. The community is complexed with Fresh - Moist White Cedar Coniferous Forest Types around the upland edge of much of the community.
SWCM1-2	21	White Cedar - Conifer Mineral Coniferous Swamp	This large community is comprised predominantly of White Cedar - Conifer Mineral Coniferous Swamp, with a complex of Fresh - Moist White Cedar - Hardwood Mixed forest Type (FOMM7-2) in the drier, more upland areas. Soil within the community varies from organic in areas to primarily mineral. The community is mature and includes a variety of canopy tree species. Canopy cover is greater than 60% throughout most of the community, with areas of reduced canopy providing a diversity of light conditions and ground cover. The dominant canopy species is Eastern White Cedar and Balsam Fir as the subdominant species. Associates include White Spruce, Black Spruce, American Larch (<i>Larix laricina</i>), Eastern Hemlock (<i>Tsuga Canadensis</i>), White Birch, Yellow Birch (<i>Betula alleghaniensis</i>), Red Maple (<i>Acer rubrum</i>), Trembling Aspen and Green Ash, among other less frequent species. Sub-canopy has lower coverage of between 10% and 25% and is comprised of immature canopy species along with occurrences of Red-osier Dogwood (<i>Cornus sericea</i>), Alternate-leaf Dogwood and Tartarian Honeysuckle. The understory and ground layer includes a variety of mostly native forbs, graminoids and ferns. Understory cover varied with light exposure and soil moisture.
SWCM3-2	2	White Cedar - Conifer Organic Coniferous Swamp	This large community extends from Wellington Road 22 to the berm of the Rubb pond. The community is dominated by White Cedar, with White Spruce, Black Spruce (<i>Picea mariana</i>), Balsam Fir, American Larch, White Ash, Green Ash, American Elm, Wild Black Cherry, Trembling Aspen, Balsam Poplar, White Birch and Yellow Birch. Canopy cover is greater than 60% throughout most of the community with small patches of opening. Understory is composed of immature canopy species, as well as Alternate-leaf Dogwood, Red-osier Dogwood, Choke Cherry, February Daphne (<i>Daphne mezereum</i>) and Tartarian Honeysuckle. Understory and ground cover is a mixture of graminoids and forbs that varied greatly in cover and composition depending on moisture and canopy cover. The community contains an inclusion of Cattail Organic Meadow Marsh and is complexed with a Fresh - Moist White Cedar - Hardwood Mixed Forest Type (FOMM7-2), which is of similar species composition, but with dry mineral soils.
<i>Mixed Swamp (SWM)</i>			
SWMO1-1	10	White Cedar - Hardwood Organic Mixed Swamp	This community is primarily White Cedar - Hardwood Organic Mixed Swamp, complexed with Fresh - Moist White Cedar - Hardwood Mixed Forest Type (FOMM7-2), which is of similar species composition, but dry and with mineral soil. Dominant canopy species within the community are White Cedar and Balsam Fir with a variety of deciduous species throughout the community. Understory and groundcover varied inversely with canopy cover, species composition varies based on moisture. The community is primarily organic soil with a complex of mineral soils.
SWMO3-3	3	White Birch - Conifer Organic Mixed Swamp	This riverine community is similar in species composition to the adjacent White Cedar - Conifer Organic Coniferous Swamp community, with a lower component of conifers and high proportion of White Birch, Yellow Birch and Poplar species. The community is located along the tributary and is bordered by a driveway that allows greater light penetration to the understory and ground layer. Soil composition is mixed, with organic soil in the lower areas and mineral soil on the slopes. Canopy cover is greater than 60%, with sub canopy, understory and ground layer cover between 25% and 60%.
<i>Deciduous Swamp (SWD)</i>			
SWDM2-1	26	Black Ash Mineral Deciduous Swamp	This community is dominated by Black Ash (<i>Fraxinus nigra</i>), with American Elm, and Trembling Aspen associates; and Sugar Maple occurs in the more upland areas along the trail way. The community has a sparse understory of Common Elderberry (<i>Sambucus Canadensis</i>), Inserted Virginia Creeper (<i>Parthenocissus inserta</i>), Smooth Gooseberry (<i>Ribes hirtellum</i>) and Wild Black Currant (<i>Ribes americanum</i>). Herbaceous ground cover is sparse with Sensitive Fern, Downy Yellow Violet (<i>Viola pubescens</i> var. <i>pubescens</i>) and Garlic Mustard (<i>Alliaria petiolate</i>) as the most common species, along with various grasses and sedges.

ELC Code	Map ID	Vegetation Type	Community Description
SWDM4-5	24	Poplar Mineral Deciduous Swamp	This community is dominated by Trembling Aspen, with White Birch as the sub-dominant canopy species. Other tree species include; White Cedar, Black Spruce and Green Ash, with Black Cherry and Sugar Maple in the more upland areas. The understory contains immature canopy species, and a mix of Willow and Dogwood species.
SWDM4-5	29	Poplar Mineral Deciduous Swamp	This is a culturally influenced community with a residential property occupying a large portion of the community. Trembling Aspen is the dominant species in the community, with Green Ash, Manitoba Maple, White Birch, and White Cedar as associate species. Sugar Maple occurs in the more upland locations. Planted Norway Maple and White Willow are also present. Understory species consist of Inserted Virginia Creeper, Alternate-leaf Dogwood and Riverbank Grape. Pale Jewel Weed (<i>Impatiens pallida</i>) is present in the understory, but may be from anthropogenic origins based on its location along the disturbed edge of a residential property.
<i>Thicket Swamp (SWT)</i>			
SWTO2-3	28	Meadow Willow Organic Deciduous Thicket Swamp	This shrub thicket community contained a mixture of willow species, with Meadow Willow as the most dominant, other shrub species present include; Choke Cherry, Red-osier Dogwood, Wild Red Raspberry, Dwarf Raspberry, Wild Black Currant. Open areas will little low shrub cover contained a variety of herbaceous species, including Common Woolly Bulrush (<i>Scirpus cyperinus</i>), Dark-green Bulrush (<i>Scirpus atrovirens</i>), Spotted Joe-pye Weed (<i>Eutrochium maculatum</i> var. <i>maculatum</i>) and Spotted Jewel-weed (<i>Impatiens capensis</i>). Tree Canopy cover is sparse with Trembling Aspen as the most abundant canopy species.
SWTO2-6	22	Mixed Willow Organic Thicket Swamp Type	This is a diverse community comprised primarily of Mixed Willow Organic Thicket Swamp Type with complexes of Mixed Shallow Water (SAM), Mixed Mineral Meadow Marsh. The community canopy cover is between 10% - 25%, containing White Spruce, Black Spruce, White Cedar, Balsam Poplar, Green Ash and planted White Willow. The understory shrubs are dominated by an assortment of willow species, with abundant Red-osier Dogwood. There is a dense understory and ground layer of herbaceous species, including; Cattail, Marsh Marigold (<i>Caltha palustris</i>), Common Woolly Bulrush, Dark Green Bulrush, Lake Bank Sedge (<i>Carex lacustris</i>), Water Horsetail (<i>Equisetum fluviatile</i>), Broadleaf Arrowhead (<i>Sagittaria latifolia</i>) and various species of Asters and Goldenrods.
SWTO3-5	9	Red-osier Organic Deciduous Swamp	This riverine thicket swamp community is species rich with a variety of native wetland trees, shrubs, and herbaceous species. Red-osier Dogwood is the dominant species within the community, with Narrow-leaved Meadow-sweet (<i>Spiraea alba</i>), Willow Species and Reed Canary Grass (<i>Phalaris arundinacea</i>) as sub-dominant. Greenfruit Bur-reed (<i>Sparganium emersum</i>), Marsh Fern (<i>Thelypteris palustris</i>) and Lake Bank Sedge are among the herbaceous species in the community.
<i>Treed Fen (FET)</i>			
FETC1-2	14	Tamarack - White Cedar Treed Fen	This community is composed primarily of Tamarack - White Cedar Treed Fen vegetation type, with complexes of Tamarack Organic Coniferous Swamp Type (SWOC2-2), Mixed Willow Organic Deciduous Thicket Swamp Type (SWTO2-6); and Cattail Graminoid Organic Meadow Marsh Type (MAMO1-2). Canopy cover within the community varies from less than 10 % to greater than 60% coverage. The community includes a high number of regionally rare species including; Loesel's Twayblade (<i>Liparis loeselii</i>), Hooded Ladies'-tresses (<i>Spiranthes romanzoffiana</i>), Marsh Blue Violet (<i>Viola cucullata</i>), Tussock Sedge (<i>Carex stricta</i>), Kalm's Lobelia (<i>Lobelia kalmii</i>), Marsh Bellflower (<i>Campanula aparinoides</i>), Linear-leaved Willow-herb (<i>Epilobium leptophyllum</i>) and Common St. John's-wort (<i>Hypericum punctatum</i>). Water is present at or just below the surface throughout most of the community. Soil ranged from saturated calcareous mineral soil to greater than 40 cm deep organic soil.
<i>Meadow Marsh (MAM)</i>			
MAMM1-1	31	Cattail Graminoid Mineral Meadow Marsh Type	This community is dominated by Narrow-leaved Cattail (<i>Typha angustifolia</i>) with associates of Redtop (<i>Agrostis gigantea</i>), Reed Canary Grass, Orchard Grass (<i>Dactylis glomerata</i>), Spotted Jewel-weed and Purple-stemmed Aster (<i>Symphotrichum puniceum</i>). Tree and shrub species on the edge of the community include Manitoba Maple, American Elm, Red-osier Dogwood, and Wild Red Raspberry.
MAMO1-2	1	Cattail Graminoid Organic Mineral Meadow Marsh	This small Community is at the corner of Trafalgar Road and Wellington Road 22. The community has evidence of past cultural influence and clearing, as indicated by the absence of mature trees. The community is dominated by Cattails with an abundance of European Reed (<i>Phragmites australis</i> ssp. <i>australis</i>) and Reed Canary Grass. The edges of the community contain a mixture of Cattail Graminoid Organic Mineral Meadow Marsh and the adjacent White Cedar – Conifer Organic Coniferous Swamp community. A low tree and shrub layer of between 2m - 10m, composed of primarily of White Cedar, Trembling Aspen and Tamarack is starting to grow throughout the community.
<i>Shallow Marsh (MAS)</i>			
MASO1-1	8	Cattail Organic Shallow Marsh Type	This riverine community borders the Submerged Shallow Aquatic Ecosite and the associated tributary. The community is dominated by Cattails, with a groundcover of sphagnum moss. Other associated species include; Red-osier Dogwood, Black Currant, Reed Canary Grass, Lake Bank Sedge, and Tussock Sedge. Occasional occurrences of American Larch and White Cedar are present along the upland edge of the community.

ELC Code	Map ID	Vegetation Type	Community Description
<i>Submerged Shallow Aquatic (SAS)</i>			
SAS_1	7	Submerged Shallow Aquatic Ecosite	This aquatic community is of unknown maximum depth, but appears to be less than 2m throughout most of the community. The aquatic community is anthropogenic in origin through the building of a berm to dam the West Credit River. Submergent aquatic species are visible below the surface, with White Water-lily (<i>Nymphaea odorata</i> ssp. <i>odorata</i>) as the associated floating species. Cattail and Dark Green Bulrush are present along the community edge. Occasional standing snags and deadfall are present throughout the community.
<i>Mixed Shallow Aquatic</i>			
SAM_1-8	11	Water Lily - Bullhead lily Mixed Shallow Aquatic	This aquatic community is of unknown maximum depth, but appears to be less than 2m throughout most of the community. The aquatic community is of anthropogenic origin, through creation of a dug offline pond, feed by the adjacent watercourse. Submergent aquatic species are visible below the surface, with Yellow Cowily (<i>Nuphar variegata</i>) as the dominant floating species. Cattail, Dark Green Bulrush, Spotted Joe-pye Weed, Broadleaf Arrowhead are present along the community edge along with shrub species, such as willow and Red-osier Dogwood.
SAM_1-8	19	Water Lily - Bullhead lily Mixed Shallow Aquatic	This open water community comprises the shallow portions of the Hillsburgh Pond. The aquatic community is of unknown maximum depth, but appeared to be less than 2m throughout most of the community. The aquatic community is of anthropogenic origin, occurring as a result of the Hillsburgh dam at Station Street. Submergent aquatic species are visible below the surface, with White Water-lily as the dominant floating species. Cattail, and Dark Green Bulrush, Spotted Joe-pye Weed, Broadleaf Arrowhead, Jewel Weed, Bulb-bearing Water-hemlock (<i>Cicuta bulbifera</i>), Blueflag Iris (<i>Iris versicolor</i>) and Yellow Iris (<i>Iris pseudacorus</i>) are present along the community edge, along with shrub species, such as Willow species and Red-osier Dogwood. Abundant to occasional standing snags and deadfall are present throughout the community.
SAM_1-8	18	Water Lily - Bullhead lily Mixed Shallow Aquatic	This aquatic community is of unknown maximum depth, but appeared to be less than 2m throughout most of the community. The aquatic community is of anthropogenic origin, resulting from the dam. Submergent aquatic species are visible below the surface, with White Water-lily as the dominant floating species. Cattail, Dark Green Bulrush, Spotted Joe-pye Weed, Broadleaf Arrowhead and Bulb-bearing Water-hemlock are present along the community edge. Abundant to occasional standing snags and deadfall are present throughout the community.
<i>Open Aquatic (OAO)</i>			
OAW	20	Open Aquatic	This is an open water community with depth greater than 2m. No floating or emergent aquatic species are present, but submerged aquatic species are suspected.
<i>Cultural (CU)</i>			
CS	13	Cultural Savannah	This community runs along the Elora-Cataract Trail way, south of the Ainsworth Pond and is highly culturally influenced. The community has a mixture of predominantly non-native tree, shrub and herbaceous species. Canopy cover varies along the length of the community from less than 10% cover to approximately 60% cover. The most abundant canopy and sub-canopy species include; Manitoba Maple, Scots Pine, White Willow, Colorado Blue Spruce (<i>Picea pungens</i>), Norway Maple, Staghorn Sumac and Common Lilac (<i>Syringa vulgaris</i>). Understory and ground cover is primarily non-native weedy species with few native herbaceous species, such as Canada Anemone (<i>Anemone canadensis</i>).
<i>Constructed (CV)</i>			
CVR_1	Res	Residential	Residential properties, including building, driveways and yards.
CVI_1	Road	Transportation	Roadway

Appendix 5
Ecological Land Classification Data Sheets

DRAFT

ELC Community Description & Classification

Polygon: P1

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 25 2015

Polygon Description P1		Community Series: MA		Ecosite: MAM		Vegetation Type: MAM01-2 Cattail Graminoid Organic Mineral Meadow Marsh			
System Terrestrial <input type="checkbox"/> Wetland Aquatic		Topographic Feature Lacustrine Riverine <input type="checkbox"/> Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff				Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid <input type="checkbox"/> Forb Lichen Bryophyte Deciduous Coniferous Mixed			
Cover Open Shrub <input type="checkbox"/> Treed		History <input type="checkbox"/> Natural Cultural		Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural <input type="checkbox"/> Swamp Fen Bog Marsh Open Water Shallow Water					
Stand Description:					Soil Analysis:				
Community Age Pioneer Young <input type="checkbox"/> Mid-Aged Mature Old Growth				Basal Area (m²/ha)		Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect <input type="checkbox"/> Poor Very Poor			
Standing Snags <input type="checkbox"/> Rare Occasional Abundant Dominant					Soil Moisture Regime Dry Fresh <input type="checkbox"/> Moist Wet				
Deadfall Logs Rare <input type="checkbox"/> Occasional Abundant Dominant					Effective Soil Texture Organic				
Health Low <input type="checkbox"/> Medium High		Sensitivity Low Medium High		Botanical Quality Low <input type="checkbox"/> Medium High		Depth to Mottles / Gley Sample: M - cm / G - cm			
Slope <input type="checkbox"/> none gentle moderate steep (simple or complex)					Depth to Groundwater metres at surface <input type="checkbox"/> less than 1m more than 1 m		Depth to Bedrock metres at surface less than 1m more than 1 m		

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	1	Larix laricina > Picea mariana = Thuja occidentalis
2 Subcanopy	3	2	Larix laricina = Thuja occidentalis = Populus tremuloides
3 Understorey	4	4	Typha angustifolia >> Thuja occidentalis = Phalaris arundinacea
4 Ground Layer	6	2	Juncus tenuis = Thelypteris palustris = Carex sp.

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4 = >60%

Size Class Analysis ³	A	O	R	NA
	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant

Evidence of Disturbance: Mowed grass around edges

Wildlife / Habitat Observations:
NA

Comments:
Road side observation due to private property restrictions.

Community Name			Code	% Coverage
Inclusion		Complex		
Inclusion		Complex		

ELC Community Description & Classification

Polygon: P2

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 25, 2015

Polygon Description P2		Community Series: SW		Ecosite: SWC		Vegetation Type: SWCM1-2 White Cedar – Conifer Mineral Coniferous Swamp			
System Terrestrial <input type="checkbox"/> Wetland Aquatic		Topographic Feature Lacustrine <input type="checkbox"/> Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff				Dominant Plant Form Plankton Submerged Floating-lvd. <input type="checkbox"/> Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed			
Cover Open Shrub <input type="checkbox"/> Treed		History <input type="checkbox"/> Natural Cultural		Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural <input type="checkbox"/> Swamp Fen Bog Marsh Open Water Shallow Water					
Stand Description:					Soil Analysis:				
Community Age Pioneer Young Mid-Aged <input type="checkbox"/> Mature Old Growth				Basal Area (m²/ha)		Soil Drainage Very Rapid Rapid Well Moderately Well <input type="checkbox"/> Imperfect Poor Very Poor			
Standing Snags Rare Occasional <input type="checkbox"/> Abundant Dominant				Soil Moisture Regime Dry Fresh <input type="checkbox"/> Moist Wet					
Deadfall Logs Rare Occasional <input type="checkbox"/> Abundant Dominant				Effective Soil Texture Organic					
Health Low <input type="checkbox"/> Medium High		Sensitivity Low <input type="checkbox"/> Medium High		Botanical Quality Low Medium High		Depth to Mottles / Gley Sample: M - cm / G - cm			
Slope none <input type="checkbox"/> gentle moderate steep (simple or complex)					Depth to Groundwater metres at surface <input type="checkbox"/> less than 1m more than 1 m		Depth to Bedrock metres at surface less than 1m more than 1 m		

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	4	Thuja occidentalis = Abies balsamea > Larix laricina = Picea glauca
2 Subcanopy	3	3	Cornus alternifolia > Prunus virginiana
3 Understorey	4	3	Typha angustifolia = Impatiens capensis
4 Ground Layer	6	3	Equisetum arvense = Caltha palustris > Fern sp. = Carex sp.

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³		O	A	D	O
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant		< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:
Wildlife / Habitat Observations:
Comments: Assessed from property edge due to private property restrictions.

				Community Name	Code	% Coverage
Inclusion	X	Complex		Cattail Organic Meadow Marsh	MAM01-2	10 %
Inclusion		Complex	X	Fresh – Moist White Cedar – Hardwood Mixed Forest Type	FOMM7-2	20 %

ELC Community Description & Classification

Polygon: P2

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 25, 2015

Representative Photographs of Vegetation Community:



DRAFT

ELC Community Description & Classification

Polygon: P3

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 25, 2015

Polygon Description P3		Community Series: SW		Ecosite: SWM		Vegetation Type: SWM03-3 White Birch – Conifer Organic Mixed Swamp			
System Terrestrial <input type="checkbox"/> <input checked="" type="checkbox"/> Wetland Aquatic		Topographic Feature Lacustrine <input type="checkbox"/> <input checked="" type="checkbox"/> Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff				Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous <input checked="" type="checkbox"/> Mixed			
Cover Open Shrub <input type="checkbox"/> <input checked="" type="checkbox"/> Treed		History <input checked="" type="checkbox"/> Natural Cultural		Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural <input checked="" type="checkbox"/> Swamp Fen Bog Marsh Open Water Shallow Water					
Stand Description:					Soil Analysis:				
Community Age Pioneer Young Mid-Aged <input checked="" type="checkbox"/> Mature Old Growth				Basal Area (m²/ha)		Soil Drainage Very Rapid Rapid Well Moderately Well <input checked="" type="checkbox"/> Imperfect Poor Very Poor			
Standing Snags Rare Occasional <input checked="" type="checkbox"/> Abundant Dominant					Soil Moisture Regime Dry Fresh <input checked="" type="checkbox"/> Moist Wet				
Deadfall Logs Rare Occasional <input checked="" type="checkbox"/> Abundant Dominant					Effective Soil Texture Organic				
Health Low <input checked="" type="checkbox"/> Medium High		Sensitivity Low <input checked="" type="checkbox"/> Medium High		Botanical Quality Low Medium High		Depth to Mottles / Gley Sample: M - cm / G - cm			
Slope none gentle moderate steep (simple or complex)					Depth to Groundwater metres at surface less than 1m more than 1 m		Depth to Bedrock metres at surface less than 1m more than 1 m		

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	4	Betula papyrifera > Thuja occidentalis > Betula alleghaniensis
2 Subcanopy	3	3	Cornus alternifolia
3 Understorey	4	3	Impatiens capensis > Ranunculus abortivus
4 Ground Layer	6	3	Caltha palustris

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4 = >60%

Size Class Analysis ³	Ø	A	D	O
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:
Wildlife / Habitat Observations:
Comments: Assessed from property edge due to private property restrictions.

		Community Name	Code	% Coverage
Inclusion	Complex			
Inclusion	Complex			

ELC Community Description & Classification

Polygon: P3

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 25, 2015

Representative Photographs of Vegetation Community:



DR

ELC Community Description & Classification

Polygon: P4

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 25, 2015

Polygon Description P4	Community Series: FO	Ecosite: FOD	Vegetation Type: FODM5-8 Dry-Fresh Sugar Maple – White Ash Deciduous Forest	
System Terrestrial Wetland Aquatic	Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff		Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed	
Cover Open Shrub Treed	History Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water		

Stand Description:		Soil Analysis:	
Community Age Pioneer Young Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor
Standing Snags Rare Occasional Abundant Dominant		Soil Moisture Regime Dry Fresh Moist Wet	
Deadfall Logs Rare Occasional Abundant Dominant		Effective Soil Texture	
Health Low Medium High	Sensitivity Low Medium High	Botanical Quality Low Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm
Slope none gentle moderate steep (simple or complex)		Depth to Groundwater metres at surface less than 1m more than 1m	Depth to Bedrock metres at surface less than 1m more than 1m

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	1	3	Sugar Maple > Black Cherry = White Ash = White Pine
2 Subcanopy	3	4	Sugar Maple > White Ash > Black Cherry
3 Understorey	4	3	Sugar Maple > Alt Lvs Dogwood > Black Cherry = Chock Cherry
4 Ground Layer	6	3	Yellow Trout Lily > Coltsfoot > Smooth Yellow Violet

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³ ³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	O	A	D	R
	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance: Trails, garden waist
Wildlife / Habitat Observations:
Comments:

		Community Name	Code	% Coverage
Inclusion	Complex			
Inclusion	Complex			

ELC Community Description & Classification

Polygon: P4

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 25, 2015

Representative Photographs of Vegetation Community:



DRAFT

ELC Community Description & Classification

Polygon: P5

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 25, 2015

Polygon Description P5	Community Series: FO	Ecosite: FOC	Vegetation Type: FOCM2-2 Dry-Fresh White Cedar Coniferous Forest	
System Terrestrial Wetland Aquatic	Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff		Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed	
Cover Open Shrub Treed	History Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water		

Stand Description:		Soil Analysis:	
Community Age Pioneer Young Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor
Standing Snags Rare Occasional Abundant Dominant		Soil Moisture Regime Dry Fresh Moist Wet	
Deadfall Logs Rare Occasional Abundant Dominant		Effective Soil Texture	
Health Low Medium High	Sensitivity Low Medium High	Botanical Quality Low Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm
Slope none gentle moderate steep (simple or complex)		Depth to Groundwater metres at surface less than 1m more than 1 m	Depth to Bedrock metres at surface less than 1m more than 1 m

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	4	Thuja occidentalis
2 Subcanopy	3	2	Thuja occidentalis
3 Understorey			
4 Ground Layer			

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³	A	A	O	NA
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:

Wildlife / Habitat Observations:

Comments:

		Community Name	Code	% Coverage
Inclusion	Complex			
Inclusion	Complex			

ELC Community Description & Classification

Polygon: P6

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 25, 2015

Polygon Description P6	Community Series: FO	Ecosite: FOC	Vegetation Type: FOCM6 Naturalized Coniferous Plantation	
System Terrestrial Wetland Aquatic	Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff		Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed	
Cover Open Shrub Treed	History Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water		

Stand Description:		Soil Analysis:	
Community Age Pioneer Young Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor
Standing Snags Rare Occasional Abundant Dominant		Soil Moisture Regime Dry Fresh Moist Wet	
Deadfall Logs Rare Occasional Abundant Dominant		Effective Soil Texture	
Health Low Medium High	Sensitivity Low Medium High	Botanical Quality Low Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm
Slope none gentle moderate steep (simple) or complex		Depth to Groundwater metres at surface less than 1m more than 1m	Depth to Bedrock metres at surface less than 1m more than 1m

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	1	3	Larix decidua > Pinus strobus = Fraxinus americana
2 Subcanopy	3	4	Pinus strobus > Malus pumila > Thuja occidentalis
3 Understorey	4	3	Rhamnus cathartica > Thuja occidentalis > Prunus virginiana = Lonicera tartarian = Solidago altissima
4 Ground Layer	6	1	Fragaria vesca > Tussilago farfara > Epipactis helleborine

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4 = >60%

Size Class Analysis ³ ³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	O	A	D	R
	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance: Non-native planting, trail, clearing.
Wildlife / Habitat Observations:
Comments:

Community Name			Code	% Coverage	
Inclusion	X	Complex	Goldenrod Forb Meadow Type	MEFM1-1	15
Inclusion		Complex			

ELC Community Description & Classification

Polygon: P6


Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 25, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P7

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 24, 2015

Polygon Description P7	Community Series: SA	Ecosite: SAS_1 Submerged Shallow Aquatic Ecosite	Vegetation Type:
System Terrestrial Wetland <u>Aquatic</u>	Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff		Dominant Plant Form Plankton <u>Submerged</u> Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed
Cover <u>Open</u> Shrub Treed	History Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh <u>Open Water</u> Shallow Water	
Stand Description:		Soil Analysis:	
Community Age Pioneer Young <u>Mid-Aged</u> Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor
Standing Snags Rare <u>Occasional</u> Abundant Dominant		Soil Moisture Regime Dry Fresh Moist Wet	
Deadfall Logs Rare <u>Occasional</u> Abundant Dominant		Effective Soil Texture	
Health Low Medium High	Sensitivity Low Medium High	Botanical Quality Low Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm
Slope <u>none</u> gentle moderate steep (simple or complex)		Depth to Groundwater metres <u>at surface</u> less than 1m more than 1m	Depth to Bedrock metres at surface less than 1m more than 1m

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy			
2 Subcanopy			
3 Understorey			
4 Ground Layer		2	Nymphaea odorata ssp. odorata > Typha latifolia > Potamogeton

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m ² Cover Codes: 0 = none, 1 = 0% - 10%, 2 = 10% - 25%, 3 = 25% - 60%, 4 = >60%

Size Class Analysis ³	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant				

Evidence of Disturbance: Earthen Dam creating pond. No fish passage into community from downstream

Wildlife / Habitat Observations: Large Mouth Bass, Mallard Duck, King Fisher, Canada Goose, Erget

Comments: Greater than 2 meters at south end, becoming shallow in the middle and towards the north. Mud flats at the north end. Little to no vegetation

Community Name			Code	% Coverage
Inclusion	Complex			
Inclusion	Complex			

ELC Community Description & Classification

Polygon: P7

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P8

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 24, 2015

Polygon Description P8	Community Series: MA	Ecosite: MAS	Vegetation Type: MAS01-1 Cattail Organic Shallow Marsh Type	
System Terrestrial <input type="checkbox"/> Wetland Aquatic	Topographic Feature Lacustrine <input type="checkbox"/> Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff		Dominant Plant Form Plankton Submerged Floating-lvd. <input type="checkbox"/> Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed	
Cover <input type="checkbox"/> Open Shrub Treed	History Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog <input type="checkbox"/> Marsh Open Water Shallow Water		

Stand Description:		Soil Analysis:	
Community Age Pioneer Young <input type="checkbox"/> Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor <input type="checkbox"/> Very Poor
Standing Snags Rare <input type="checkbox"/> Occasional Abundant Dominant		Soil Moisture Regime Dry Fresh Moist <input type="checkbox"/> Wet	
Deadfall Logs Rare <input type="checkbox"/> Occasional Abundant Dominant		Effective Soil Texture	
Health Low <input type="checkbox"/> Medium High	Sensitivity Low Medium <input type="checkbox"/> High	Botanical Quality Low <input type="checkbox"/> Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm
Slope none gentle moderate steep (simple or complex)		Depth to Groundwater metres <input type="checkbox"/> at surface less than 1m more than 1 m	Depth to Bedrock metres at surface less than 1m <input type="checkbox"/> more than 1 m

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	1	Thuja occidentalis = Betula papyrifera > Larix laricina
2 Subcanopy	3	1	Thuja occidentalis = Betula papyrifera
3 Understorey	4	4	Typha latifolia >> Phalaris arundinacea
4 Ground Layer	6	2	Sphagnum sp.>> Caltha palustris > Carex lacustris

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³ <small>³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant</small>	O	R	NA	NA
	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:

Wildlife / Habitat Observations:

Comments:

			Community Name	Code	% Coverage
Inclusion		Complex			
Inclusion		Complex			

ELC Community Description & Classification

Polygon: P9

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 24, 2015

Polygon Description P9		Community Series: SW		Ecosite: SWT		Vegetation Type: SWT03-5 Red-osier Organic Deciduous Swamp			
System Terrestrial <input type="checkbox"/> Wetland Aquatic		Topographic Feature Lacustrine <input type="checkbox"/> Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff				Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte <input type="checkbox"/> Deciduous Coniferous Mixed			
Cover Open <input type="checkbox"/> Shrub <input type="checkbox"/> Treed		History <input type="checkbox"/> Natural Cultural		Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest <input type="checkbox"/> Thicket Cultural <input type="checkbox"/> Swamp Fen Bog Marsh Open Water Shallow Water					
Stand Description:					Soil Analysis:				
Community Age Pioneer <input type="checkbox"/> Young Mid-Aged Mature Old Growth				Basal Area (m²/ha)		Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect <input type="checkbox"/> Poor Very Poor			
Standing Snags Rare <input type="checkbox"/> Occasional Abundant Dominant					Soil Moisture Regime Dry Fresh Moist <input type="checkbox"/> Wet				
Deadfall Logs Rare <input type="checkbox"/> Occasional Abundant Dominant					Effective Soil Texture Organic				
Health Low <input type="checkbox"/> Medium High		Sensitivity Low <input type="checkbox"/> Medium High		Botanical Quality Low Medium <input type="checkbox"/> High		Depth to Mottles / Gley Sample: M - cm / G - cm			
Slope <input type="checkbox"/> none gentle moderate steep (simple or complex)					Depth to Groundwater metres <input type="checkbox"/> at surface less than 1m more than 1 m		Depth to Bedrock metres at surface less than 1m more than 1 m		

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	3	1	Thuja occidentalis > Acer saccharinum
2 Subcanopy	4	1	Thuja occidentalis = Salix discolor
3 Understory	5	4	Cornus stolonifera >> Salix discolor = Spirea alba
4 Ground Layer	6	3	Phalaris arundinacea > Typha latifolia > Caltha palustris

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³	0	R	NA	NA
	³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH

Evidence of Disturbance:

Wildlife / Habitat Observations:

Comments:

			Community Name	Code	% Coverage
Inclusion		Complex			
Inclusion		Complex			

ELC Community Description & Classification

Polygon: P9

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P10

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 24, 2015

Polygon Description P10	Community Series: SW	Ecosite: SWM	Vegetation Type: SWM01-1 White Cedar – Hardwood Organic Mixed Swamp	
System Terrestrial <input type="checkbox"/> Wetland Aquatic	Topographic Feature Lacustrine <input type="checkbox"/> Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff		Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous <input type="checkbox"/> Coniferous Mixed	
Cover Open Shrub <input type="checkbox"/> Treed	History <input type="checkbox"/> Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural <input type="checkbox"/> Swamp Fen Bog Marsh Open Water Shallow Water		

Stand Description:		Soil Analysis:	
Community Age Pioneer Young <input type="checkbox"/> Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect <input type="checkbox"/> Poor Very Poor
Standing Snags Rare Occasional <input type="checkbox"/> Abundant Dominant		Soil Moisture Regime Dry Fresh Moist <input type="checkbox"/> Wet	
Deadfall Logs Rare Occasional <input type="checkbox"/> Abundant Dominant		Effective Soil Texture	
Health Low <input type="checkbox"/> Medium High	Sensitivity Low <input type="checkbox"/> Medium High	Botanical Quality Low <input type="checkbox"/> Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm
Slope none <input type="checkbox"/> gentle moderate steep (simple or complex)		Depth to Groundwater metres at surface <input type="checkbox"/> less than 1m more than 1 m	Depth to Bedrock metres at surface less than 1m more than 1 m

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	4	White Cedar > Balsam Fir > White Birch
2 Subcanopy	3	3	White Cedar > Balsam Fir > White Birch
3 Understorey	4	3	White Cedar > Red osier dogwood
4 Ground Layer	6	2	Sensitive Fern > Sedge sp. > Sphagnum

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³	A	A	O	NA
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:
Wildlife / Habitat Observations:
Comments:

		Community Name	Code	% Coverage
Inclusion	<input type="checkbox"/>	Complex		
Inclusion	<input type="checkbox"/>	Complex		

ELC Community Description & Classification

Polygon: P10

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 14; Aug 10; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P11

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Polygon Description P11	Community Series: SA	Ecosite: SAM	Vegetation Type: SAM_1-8 Water Lily – Bullhead Lily Mixed Shallow Aquatic		
System Terrestrial Wetland Aquatic	Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff			Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed	
Cover Open Shrub Treed	History Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water			
Stand Description:			Soil Analysis:		
Community Age Pioneer Young Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor		
Standing Snags Rare Occasional Abundant Dominant			Soil Moisture Regime Dry Fresh Moist Wet		
Deadfall Logs Rare Occasional Abundant Dominant			Effective Soil Texture		
Health Low Medium High	Sensitivity Low Medium High	Botanical Quality Low Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm		
Slope none gentle moderate steep (simple or complex)			Depth to Groundwater metres at surface less than 1m more than 1m	Depth to Bedrock metres at surface less than 1m more than 1m	

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy			
2 Subcanopy			
3 Understorey			
4 Ground Layer	6	2	Nuphar variegata > Sagittaria latifolia = Typha latifolia

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=< 0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³	NA	NA	NA	NA
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:

Wildlife / Habitat Observations:

Comments:

Community Name			Code	% Coverage
Inclusion	Complex			
Inclusion	Complex			

ELC Community Description & Classification

Polygon: P11

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P12

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22 ; Aug 10; Sept 24, 2015

Polygon Description P12		Community Series: ME	Ecosite: MEM	Vegetation Type: MEMM3 Dry – Fresh Mixed Meadow	
System Terrestrial Wetland Aquatic		Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff		Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed	
Cover Open Shrub Treed		History Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Meadow Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water		
Stand Description:			Soil Analysis:		
Community Age Pioneer Young Mid-Aged Mature Old Growth		Basal Area (m²/ha)		Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor	
Standing Snags Rare Occasional Abundant Dominant			Soil Moisture Regime Dry Fresh Moist Wet		
Deadfall Logs Rare Occasional Abundant Dominant			Effective Soil Texture		
Health Low Medium High		Sensitivity Low Medium High		Botanical Quality Low Medium High	
Slope none gentle moderate steep (simple or complex)		Depth to Mottles / Gley Sample: M - cm / G - cm		Depth to Groundwater metres at surface less than 1m more than 1m	
				Depth to Bedrock metres at surface less than 1m more than 1m	

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	3	2	Populus balsamifera = Pinus sylvestris
2 Subcanopy	4	2	Populus balsamifera = Pinus sylvestris > Prunus virginiana
3 Understorey	5	4	Phalaris arundinacea = Solidago Canadensis > Bromus inermis = Asclepias syriaca
4 Ground Layer	6	4	Vicia cracca > Linaria vulgaris = Anemone canadensis

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³			
	O	O	Na
	< 10 cm DBH	10 to 24 cm DBH	> 50 cm DBH

³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant

Evidence of Disturbance:

Wildlife / Habitat Observations:

Comments:

		Community Name	Code	% Coverage
Inclusion	Complex			
Inclusion	Complex			

ELC Community Description & Classification

Polygon: P13

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; July 30; Sept 24, 2015

Polygon Description P13	Community Series	Ecosite: CS Cultural Savannah	Vegetation Type
System Terrestrial Wetland Aquatic	Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff	Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed	
Cover Open Shrub Treed	History Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water	

Stand Description:		Soil Analysis:	
Community Age Pioneer Young Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor
Standing Snags Rare Occasional Abundant Dominant		Soil Moisture Regime Dry Fresh Moist Wet	
Deadfall Logs Rare Occasional Abundant Dominant		Effective Soil Texture	
Health Low Medium High	Sensitivity Low Medium High	Botanical Quality Low Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm
Slope none gentle moderate steep (simple or complex)		Depth to Groundwater metres at surface less than 1m more than 1m	Depth to Bedrock metres at surface less than 1m more than 1m

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	2	Acer negundo > Malus pumila > Pinus sylvestris = Populus tremuloides
2 Subcanopy	3	3	Malus pumila > Pinus sylvestris = Syringa vulgaris = Rhus typhina
3 Understorey	4	2	Lonicera tatarica > Solidago Canadensis = Lathyrus latifolius = Vitis Riparia
4 Ground Layer	6	3	Alliaria petiolate > Galium triflorum > Anemone canadensis

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³	O	A	R	NA
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance: Trail, invasive species, cut grass
Wildlife / Habitat Observations:
Comments:

Community Name		Code	% Coverage
Inclusion	Complex		
Inclusion	Complex		

ELC Community Description & Classification

Polygon: P13


Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; July 30; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P14

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Polygon Description P14	Community Series: FE	Ecosite: FET	Vegetation Type: FETC1-2 Tamarack – White Cedar Treed Fen
System Terrestrial <input type="checkbox"/> Wetland Aquatic	Topographic Feature Lacustrine <input type="checkbox"/> Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff	Dominant Plant Form Plankton Submerged Floating-lvd. <input type="checkbox"/> Graminoid Forb Lichen Bryophyte Deciduous Coniferous <input type="checkbox"/> Mixed	
Cover <input type="checkbox"/> Open Shrub Treed	History <input type="checkbox"/> Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp <input type="checkbox"/> Fen Bog Marsh Open Water Shallow Water	

Stand Description:		Soil Analysis:	
Community Age Pioneer <input type="checkbox"/> Young Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor <input type="checkbox"/> Very Poor
Standing Snags Rare Occasional <input type="checkbox"/> Abundant Dominant		Soil Moisture Regime Dry Fresh Moist <input type="checkbox"/> Wet	
Deadfall Logs Rare Occasional <input type="checkbox"/> Abundant Dominant		Effective Soil Texture	
Health Low Medium <input type="checkbox"/> High	Sensitivity Low Medium <input type="checkbox"/> High	Botanical Quality Low Medium <input type="checkbox"/> High	Depth to Mottles / Gley Sample: M - cm / G - cm
Slope <input type="checkbox"/> none gentle moderate steep (simple or complex)	Depth to Groundwater metres <input type="checkbox"/> at surface less than 1m more than 1 m		Depth to Bedrock metres at surface less than 1m more than 1 m

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	3	2	Thuja occidentalis = Larix laricina
2 Subcanopy	4	2	Thuja occidentalis > Larix laricina
3 Understorey	5	3	Salix sp. > Cornus stolonifera = Typha angustifolia
4 Ground Layer	6	3	Carex sp. > Caltha palustris > Liparis loeselii > Impatiens capensis

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0% - 10%, 2 = 10% - 25%, 3 = 25% - 60%, 4 = >60%

Size Class Analysis ³ <small>³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant</small>	D	A	O	NA
	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:
Wildlife / Habitat Observations: Leopard-frog, muskrat den
Comments: Soil primarily organic > 50 cm with areas of calcareous mineral soil.

				Community Name	Code	% Coverage
Inclusion		Complex	X	Tamarack Organic Coniferous Swamp Type	SWOC2-2	15
Inclusion		Complex	X	Mixed Willow Organic Deciduous Thicket Swamp Type	SWTO2-6	10
		Complex	X	Cattail Graminoid Organic Meadow Marsh Type	MAMO1-2	20

ELC Community Description & Classification

Polygon: P14

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P15

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Polygon Description P15		Community Series: FO	Ecosite: FOD	Vegetation Type: FODM8-1 Fresh – Moist Poplar Deciduous Forest	
System Terrestrial Wetland Aquatic		Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff		Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed	
Cover Open Shrub Treed		History Natural Cultural		Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water	
Stand Description:			Soil Analysis:		
Community Age Pioneer Young Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor		
Standing Snags Rare Occasional Abundant Dominant		Soil Moisture Regime Dry Fresh Moist Wet			
Deadfall Logs Rare Occasional Abundant Dominant		Effective Soil Texture Mineral			
Health Low Medium High	Sensitivity Low Medium High	Botanical Quality Low Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm		
Slope none gentle moderate steep (simple or complex)		Depth to Groundwater metres at surface less than 1m more than 1m	Depth to Bedrock metres at surface less than 1m more than 1m		

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	4	4	Populus tremuloides > Populus balsamifera > Acer platanoides
2 Subcanopy	3	3	Populus tremuloides > Populus balsamifera > Acer platanoides = Acer negundo
3 Understorey	2	3	Cornus alternifolia > Thuja occidentalis = Prunus virginiana > Symphyotrichum novae-angliae
4 Ground Layer	1	2	Tussilago farfara > Aegopodium podagraria > Myosotis scorpioides = Fragaria vesca

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³	O	A	A	NA
	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant

Evidence of Disturbance:

Wildlife / Habitat Observations:

Comments:

Inclusion		Complex	Community Name	Code	% Coverage
Inclusion		Complex			
Inclusion		Complex			

ELC Community Description & Classification

Polygon: P16

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Polygon Description P16		Community Series: FO	Ecosite: FOD	Vegetation Type: FODM6 Fresh – Moist Sugar Maple Deciduous Forest Ecosite	
System Terrestrial Wetland Aquatic		Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff		Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed	
Cover Open Shrub Treed		History Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water		
Stand Description:			Soil Analysis:		
Community Age Pioneer Young Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor		
Standing Snags Rare Occasional Abundant Dominant		Soil Moisture Regime Dry Fresh Moist Wet			
Deadfall Logs Rare Occasional Abundant Dominant		Effective Soil Texture			
Health Low Medium High	Sensitivity Low Medium High	Botanical Quality Low Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm		
Slope none gentle moderate steep (simple or complex)		Depth to Groundwater metres at surface less than 1m more than 1 m		Depth to Bedrock metres at surface less than 1m more than 1 m	

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	3	Acer saccharum ssp. saccharum > Acer negundo > Fraxinus pennsylvanica > Juglans nigra
2 Subcanopy	3	4	Acer negundo > Fraxinus pennsylvanica > Vitis riparia
3 Understorey	4	4	Lonicera tatarica > Rubus allegheniensis > Vitis riparia = Echinocystis lobata
4 Ground Layer	6	4	Alliaria petiolate >> Anemone Canadensis = Vicia cracca = Equisetum arvense

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³	A	A	O	NA
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:

Wildlife / Habitat Observations:

Comments: Cut-grass, planted non-native plants, occupied residence. ELC done from property edge.

Community Name		Code	% Coverage
Inclusion	Complex		
Inclusion	Complex		

ELC Community Description & Classification

Polygon: P16

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Representative Photographs of Vegetation Community:



DR

ELC Community Description & Classification

Polygon: P17

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Polygon Description P17	Community Series: SW	Ecosite: SWC	Vegetation Type: SWCM1-2 White Cedar – Conifer Mineral Coniferous Swamp		
System Terrestrial <input type="checkbox"/> Wetland Aquatic	Topographic Feature <input type="checkbox"/> Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland <input type="checkbox"/> Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff		Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous <input type="checkbox"/> Coniferous Mixed		
Cover Open Shrub <input type="checkbox"/> Treed	History <input type="checkbox"/> Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural <input type="checkbox"/> Swamp Fen Bog Marsh Open Water Shallow Water			

Stand Description:		Soil Analysis:	
Community Age Pioneer Young Mid-Aged <input type="checkbox"/> Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well <input type="checkbox"/> Imperfect Poor Very Poor
Standing Snags Rare Occasional <input type="checkbox"/> Abundant Dominant		Soil Moisture Regime Dry Fresh <input type="checkbox"/> Moist Wet	
Deadfall Logs Rare Occasional <input type="checkbox"/> Abundant Dominant		Effective Soil Texture	
Health Low Medium <input type="checkbox"/> High	Sensitivity Low Medium <input type="checkbox"/> High	Botanical Quality Low Medium <input type="checkbox"/> High	Depth to Mottles / Gley Sample: M - cm / G - cm
Slope none <input type="checkbox"/> gentle moderate steep (simple or complex)		Depth to Groundwater metres at surface <input type="checkbox"/> less than 1m more than 1 m	Depth to Bedrock metres at surface less than 1m more than 1 m

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	4	Thuja occidentalis >> Fraxinus pennsylvanica > Populus tremuloides
2 Subcanopy	3	2	Thuja occidentalis >> Abies balsamea > Fraxinus pennsylvanica
3 Understorey	4	2	Ribes americanum = Cornus alternifolia > Cornus stolonifera
4 Ground Layer	6	3	Caltha palustris > Equisetum palustre > Impatiens capensis

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7= <0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³ <small>³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant</small>	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH
--	-------------	-----------------	-----------------	-------------

Evidence of Disturbance: Trails, sand bags, water control structures.
Wildlife / Habitat Observations: Canada goose nest, snapping turtle along edge of pond.
Comments: Mostly Wetland, with areas of upland cedar forest on slopes around pond.

Community Name				Code	% Coverage
Inclusion	Complex	X	Fresh – Moist White Cedar Coniferous Forest Type	FOCM4-1	20
Inclusion	Complex				

ELC Community Description & Classification

Polygon: P17

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P18

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Polygon Description P18	Community Series: SA	Ecosite: SAM	Vegetation Type: SAM_1-8 Water Lily – Bullhead Lily Mixed Shallow Aquatic		
System Terrestrial Wetland Aquatic	Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff			Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed	
Cover Open Shrub Treed	History Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water			
Stand Description:			Soil Analysis:		
Community Age Pioneer Young Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor		
Standing Snags Rare Occasional Abundant Dominant			Soil Moisture Regime Dry Fresh Moist Wet		
Deadfall Logs Rare Occasional Abundant Dominant			Effective Soil Texture		
Health Low Medium High	Sensitivity Low Medium High	Botanical Quality Low Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm		
Slope none gentle moderate steep (simple or complex)			Depth to Groundwater metres at surface less than 1m more than 1 m	Depth to Bedrock metres at surface less than 1m more than 1 m	

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy			
2 Subcanopy			
3 Understorey			
4 Ground Layer	6	2	Nymphaea odorata ssp. odorata > Typha latifolia > Sagittaria latifolia

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4 = >60%

Size Class Analysis ³	NA	NA	NA	NA
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:

Wildlife / Habitat Observations:

Comments:

Community Name			Code	% Coverage
Inclusion	Complex			
Inclusion	Complex			

ELC Community Description & Classification

Polygon: P18

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Representative Photographs of Vegetation Community:



DRAFT

ELC Community Description & Classification

Polygon: P19

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Polygon Description P19	Community Series: SA	Ecosite: SAM	Vegetation Type: SAM_1-8 Water Lily – Bullhead Lily Mixed Shallow Aquatic		
System Terrestrial Wetland Aquatic	Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff			Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed	
Cover Open Shrub Treed	History Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water			
Stand Description:			Soil Analysis:		
Community Age Pioneer Young Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor		
Standing Snags Rare Occasional Abundant Dominant			Soil Moisture Regime Dry Fresh Moist Wet		
Deadfall Logs Rare Occasional Abundant Dominant			Effective Soil Texture		
Health Low Medium High	Sensitivity Low Medium High	Botanical Quality Low Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm		
Slope none gentle moderate steep (simple or complex)			Depth to Groundwater metres at surface less than 1m more than 1m	Depth to Bedrock metres at surface less than 1m more than 1m	

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy			
2 Subcanopy			
3 Understorey			
4 Ground Layer	6	2	Nymphaea odorata ssp. odorata > Typha latifolia > Sagittaria latifolia

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³	NA	NA	NA	NA
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:

Wildlife / Habitat Observations:

Comments:

Community Name			Code	% Coverage
Inclusion	Complex			
Inclusion	Complex			

ELC Community Description & Classification

Polygon: P19


Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P20

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Polygon Description P20	Community Series:	Ecosite: OAW Open Aquatic	Vegetation Type:
System Terrestrial Wetland Aquatic	Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff		Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed
Cover Open Shrub Treed	History Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Barren Tallgrass Prairie Savannah Woodland Open Water Shallow Water	Cliff Talus Alvar Rock Barren Crevice-Cave Sand
Stand Description:		Soil Analysis:	
Community Age Pioneer Young Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor
Standing Snags Rare Occasional Abundant Dominant		Soil Moisture Regime Dry Fresh Moist Wet	
Deadfall Logs Rare Occasional Abundant Dominant		Effective Soil Texture	
Health Low Medium High	Sensitivity Low Medium High	Botanical Quality Low Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm
Slope none gentle moderate steep (simple or complex)		Depth to Groundwater metres at surface less than 1m more than 1m	Depth to Bedrock metres at surface less than 1m more than 1m

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy			
2 Subcanopy			
3 Understorey			
4 Ground Layer	6	2	Nymphaea odorata ssp. odorata

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4 = >60%

Size Class Analysis ³	NA	NA	NA	NA
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:

Wildlife / Habitat Observations:

Comments:
Deep open water portion of main pond.

			Community Name	Code	% Coverage
Inclusion		Complex			
Inclusion		Complex			

ELC Community Description & Classification

Polygon: P20

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 22; Aug 5; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P21

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date:

Polygon Description P21	Community Series: SW	Ecosite: SWC	Vegetation Type: SWCM1-2 White Cedar – Conifer Mineral Coniferous Swamp		
System Terrestrial <input type="checkbox"/> Wetland Aquatic	Topographic Feature Lacustrine Riverine <input type="checkbox"/> Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff		Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous <input type="checkbox"/> Coniferous Mixed		
Cover Open Shrub <input type="checkbox"/> Treed	History <input type="checkbox"/> Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural <input type="checkbox"/> Swamp Fen Bog Marsh Open Water Shallow Water			

Stand Description:		Soil Analysis:	
Community Age Pioneer Young <input type="checkbox"/> Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well <input type="checkbox"/> Imperfect Poor Very Poor
Standing Snags Rare <input type="checkbox"/> Occasional Abundant Dominant		Soil Moisture Regime Dry Fresh <input type="checkbox"/> Moist Wet	
Deadfall Logs Rare Occasional <input type="checkbox"/> Abundant Dominant		Effective Soil Texture Organic (D) / Mineral (O)	
Health Low Medium <input type="checkbox"/> High	Sensitivity Low <input type="checkbox"/> Medium High	Botanical Quality Low Medium <input type="checkbox"/> High	Depth to Mottles / Gley Sample: M - cm / G - cm
Slope none <input type="checkbox"/> gentle moderate steep (simple or complex)		Depth to Groundwater metres at surface <input type="checkbox"/> less than 1m more than 1 m	Depth to Bedrock metres at surface less than 1m more than 1 m

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	3	Thuja occidentalis > Abies balsamea = Picea mariana = Picea glauca
2 Subcanopy	3	2	Thuja occidentalis > Abies balsamea = Populus balsamifera > Betula papyrifera
3 Understorey	4	2	Abies balsamea = Thuja occidentalis > Cornus alternifolia
4 Ground Layer	6	2	Caltha palustris > Onoclea sensibilis > Anemone canadensis

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³ <small>³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant</small>	A	A	A	R
	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:
Wildlife / Habitat Observations:
Comments: Most of the community is on private land without access. Evaluation from trail edge.

Community Name				Code	% Coverage
Inclusion	Complex	X	Fresh – Moist White Cedar – Hardwood Mixed forest Type	FOMM7-2	20
Inclusion	Complex				

ELC Community Description & Classification

Polygon: P21

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date:

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P22

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 24, 2015

Polygon Description P22		Community Series: SW		Ecosite: SWT		Vegetation Type: SWT02-6 Mixed Willow Organic Thicket Swamp Type			
System Terrestrial <input checked="" type="checkbox"/> Wetland Aquatic		Topographic Feature Lacustrine Riverine <input checked="" type="checkbox"/> Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff				Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous <input checked="" type="checkbox"/> Coniferous Mixed			
Cover <input checked="" type="checkbox"/> Open Shrub Treed		History <input checked="" type="checkbox"/> Natural Cultural		Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest <input checked="" type="checkbox"/> Thicket Cultural <input checked="" type="checkbox"/> Swamp Fen Bog Marsh Open Water Shallow Water					
Stand Description:					Soil Analysis:				
Community Age Pioneer <input checked="" type="checkbox"/> Young Mid-Aged Mature Old Growth				Basal Area (m²/ha)		Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect <input checked="" type="checkbox"/> Poor <input checked="" type="checkbox"/> Very Poor			
Standing Snags Rare Occasional <input checked="" type="checkbox"/> Abundant Dominant					Soil Moisture Regime Dry Fresh Moist <input checked="" type="checkbox"/> Wet				
Deadfall Logs Rare Occasional <input checked="" type="checkbox"/> Abundant Dominant					Effective Soil Texture				
Health Low <input checked="" type="checkbox"/> Medium High		Sensitivity Low <input checked="" type="checkbox"/> Medium High		Botanical Quality Low <input checked="" type="checkbox"/> Medium High		Depth to Mottles / Gley Sample: M - cm / G - cm			
Slope <input checked="" type="checkbox"/> none gentle moderate steep (simple or complex)					Depth to Groundwater metres at surface less than 1m more than 1 m		Depth to Bedrock metres at surface less than 1m more than 1 m		

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	2	Picea glauca > Betula papyrifera = Populus tremuloides
2 Subcanopy	3	1	Picea glauca > Thuja occidentalis = Picea mariana
3 Understorey	4	3	Salix sp. > Cornus stolonifera > Typha latifolia > Phalaris arundinacea
4 Ground Layer	6	4	Caltha palustris > Impatiens capensis > Sedge sp.

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4 = >60%

Size Class Analysis ³	O	O	O	NA
	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant

Evidence of Disturbance:

Wildlife / Habitat Observations:

Comments: No access. Evaluation from edge.

				Community Name	Code	% Coverage
Inclusion		Complex	X	Mixed Shallow Water	SAM	15
Inclusion		Complex	X	Mixed Mineral Meadow Marsh	MAMM3-1	15

ELC Community Description & Classification

Polygon: P22

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P23

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 24, 2015

Polygon Description P23		Community Series: FO		Ecosite: FOM		Vegetation Type: FOMM7-2 Fresh – Moist White Cedar – Hardwood Mixed Forest			
System Terrestrial Wetland Aquatic		Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff				Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed			
Cover Open Shrub Treed		History Natural Cultural		Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water					
Stand Description:					Soil Analysis:				
Community Age Pioneer Young Mid-Aged Mature Old Growth				Basal Area (m²/ha)		Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor			
Standing Snags Rare Occasional Abundant Dominant					Soil Moisture Regime Dry Fresh Moist Wet				
Deadfall Logs Rare Occasional Abundant Dominant					Effective Soil Texture				
Health Low Medium High		Sensitivity Low Medium High		Botanical Quality Low Medium High		Depth to Mottles / Gley Sample: M - cm / G - cm			
Slope none gentle moderate steep (simple or complex)					Depth to Groundwater metres at surface less than 1m more than 1 m			Depth to Bedrock metres at surface less than 1m more than 1 m	

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	3	White birch > white cedar > black cherry = trembling aspen
2 Subcanopy	3	3	Tremblin aspen = black cherry = white birch > alt leave dogwood
3 Understorey	4	3	Alt leave dogwood = Ash sp.
4 Ground Layer	6	3	Yellow trout lilly = Smooth yellow violet

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³					
		A	A	A	R
		< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant

Evidence of Disturbance:

Wildlife / Habitat Observations:

Comments: Small area of pure White Cedar Stand.

				Community Name	Code	% Coverage
Inclusion		Complex				
Inclusion		Complex				

ELC Community Description & Classification

Polygon: P23

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 24, 2015

Representative Photographs of Vegetation Community:



DR

ELC Community Description & Classification

Polygon: P24

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 24, 2015

Polygon Description P24	Community Series: SW	Ecosite: SWD	Vegetation Type: SWDM4-5 Poplar Mineral Deciduous Swamp		
System Terrestrial <input type="checkbox"/> Wetland Aquatic	Topographic Feature Lacustrine Riverine <input type="checkbox"/> Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff		Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte <input type="checkbox"/> Deciduous Coniferous Mixed		
Cover Open Shrub <input type="checkbox"/> Treed	History <input type="checkbox"/> Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland <input type="checkbox"/> Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water			

Stand Description:		Soil Analysis:	
Community Age Pioneer Young <input type="checkbox"/> Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well <input type="checkbox"/> Imperfect Poor Very Poor
Standing Snags <input type="checkbox"/> Rare Occasional Abundant Dominant		Soil Moisture Regime Dry Fresh <input type="checkbox"/> Moist Wet	
Deadfall Logs Rare <input type="checkbox"/> Occasional Abundant Dominant		Effective Soil Texture Mineral	
Health Low <input type="checkbox"/> Medium High	Sensitivity Low <input type="checkbox"/> Medium High	Botanical Quality Low <input type="checkbox"/> Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm
Slope none <input type="checkbox"/> gentle moderate steep (simple or complex)		Depth to Groundwater metres at surface less than 1m more than 1 m	Depth to Bedrock metres at surface less than 1m more than 1 m

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	3	Populus tremuloides > Betula papyrifera > Thuja occidentalis
2 Subcanopy	4	3	Populus tremuloides > Betula papyrifera > Thuja occidentalis
3 Understorey	3	3	Salix sp. > Cornus alternifolia > Eutrochium maculatum
4 Ground Layer	6	3	Onoclea sensibilis > Equisetum sp. = Carex sp. > Fragaria vesca

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³ <small>³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant</small>	A	A	O	R
	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:
Wildlife / Habitat Observations:
Comments:

			Community Name	Code	% Coverage
Inclusion		Complex			
Inclusion		Complex			

ELC Community Description & Classification

Polygon: P24

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P25

Project No: 12-137A		Project Name: Hillsburgh Dam EA		Surveyor(s): RH		Date: May 21; July 30; Sept 24, 2015	
Polygon Description P25		Community Series: FO		Ecosite: FOD		Vegetation Type: FODM8-1 Fresh – Moist Poplar Deciduous Forest	
System Terrestrial Wetland Aquatic		Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff				Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed	
Cover Open Shrub Treed		History Natural Cultural		Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water			
Stand Description:				Soil Analysis:			
Community Age Pioneer Young Mid-Aged Mature Old Growth			Basal Area (m²/ha)		Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor		
Standing Snags Rare Occasional Abundant Dominant				Soil Moisture Regime Dry Fresh Moist Wet			
Deadfall Logs Rare Occasional Abundant Dominant				Effective Soil Texture			
Health Low Medium High		Sensitivity Low Medium High		Botanical Quality Low Medium High		Depth to Mottles / Gley Sample: M - cm / G - cm	
Slope none gentle moderate steep (simple or complex)				Depth to Groundwater metres at surface less than 1m more than 1m		Depth to Bedrock metres at surface less than 1m more than 1m	

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	3	Populus tremuloides = Acer saccharum ssp. saccharum > Thuja occidentalis = Fraxinus americana
2 Subcanopy	3	3	Populus tremuloides = Acer saccharum ssp. saccharum > Fraxinus americana = Cornus alternifolia > Rhus typhina
3 Understorey	4	3	Rubus idaeus ssp. strigosus > Ribes americanum = Inserted Virginia Creeper
4 Ground Layer	6	3	Alliaria petiolate > Fragaria vesca > Erythronium americanum

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7= < 0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³	O	A	O	
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:
Wildlife / Habitat Observations:
Comments:

			Community Name	Code	% Coverage
Inclusion		Complex			
Inclusion		Complex			

ELC Community Description & Classification

Polygon: P25


Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P26

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 24, 2015

Polygon Description P26	Community Series: SW	Ecosite: SWD	Vegetation Type: SWDM2-1 Black Ash Mineral Deciduous Swamp		
System Terrestrial <input type="checkbox"/> Wetland Aquatic	Topographic Feature Lacustrine Riverine <input type="checkbox"/> Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff		Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte <input type="checkbox"/> Deciduous Coniferous Mixed		
Cover Open Shrub <input type="checkbox"/> Treed	History Natural Cultural	Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah <input type="checkbox"/> Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water			

Stand Description:		Soil Analysis:	
Community Age Pioneer Young <input type="checkbox"/> Mid-Aged Mature Old Growth		Basal Area (m²/ha)	Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor
Standing Snags <input type="checkbox"/> Rare Occasional Abundant Dominant		Soil Moisture Regime Dry Fresh Moist <input type="checkbox"/> Wet	
Deadfall Logs Rare <input type="checkbox"/> Occasional Abundant Dominant		Effective Soil Texture	
Health Low <input type="checkbox"/> Medium High	Sensitivity Low <input type="checkbox"/> Medium High	Botanical Quality Low <input type="checkbox"/> Medium High	Depth to Mottles / Gley Sample: M - cm / G - cm
Slope none <input type="checkbox"/> gentle moderate steep (simple or complex)		Depth to Groundwater metres at surface less than 1m more than 1 m	Depth to Bedrock metres at surface less than 1m more than 1 m

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	3	Fraxinus nigra > Populus tremuloides > Acer saccharum ssp. saccharum
2 Subcanopy	3	2	Fraxinus nigra > Populus tremuloides > Ulmus americana > Parthenocissus inserta
3 Understorey	4	2	Ribes americanum > Parthenocissus inserta > Ribes hirtellum
4 Ground Layer	6	2	Onoclea sensibilis > Impatiens capensis = Alliaria petiolate > Anemone canadensis

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³	O	D	O	
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:

Wildlife / Habitat Observations:

Comments:

Inclusion		Complex	Community Name	Code	% Coverage
Inclusion		Complex			
Inclusion		Complex			

ELC Community Description & Classification

Polygon: P26


Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P27

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: July 30; Sept 24, 2015

Polygon Description P27		Community Series: Fo		Ecosite: FOC		Vegetation Type: FOCM6			
System Terrestrial Wetland Aquatic		Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff				Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed			
Cover Open Shrub Treed		History Natural Cultural		Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water					
Stand Description:					Soil Analysis:				
Community Age Pioneer Young Mid-Aged Mature Old Growth				Basal Area (m²/ha)		Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor			
Standing Snags Rare Occasional Abundant Dominant					Soil Moisture Regime Dry Fresh Moist Wet				
Deadfall Logs Rare Occasional Abundant Dominant					Effective Soil Texture				
Health Low Medium High		Sensitivity Low Medium High		Botanical Quality Low Medium High		Depth to Mottles / Gley Sample: M - cm / G - cm			
Slope none gentle moderate steep (simple or complex)					Depth to Groundwater metres at surface less than 1m more than 1m		Depth to Bedrock metres at surface less than 1m more than 1m		

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	4	Larix sp. = Picea abies = Picea glauca = Pinus strobus = Thuja occidentalis
2 Subcanopy			
3 Understorey			
4 Ground Layer			

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7= < 0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant				

Evidence of Disturbance:
Wildlife / Habitat Observations:
Comments: Surveyed from a distance, from road side.

Community Name				Code	% Coverage
Inclusion		Complex			
Inclusion		Complex			

ELC Community Description & Classification

Polygon: P27

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: July 30; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P28

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 24, 2015

Polygon Description P28		Community Series: SW		Ecosite: SWT		Vegetation Type: SWT02-3 Meadow Willow Organic Deciduous Thicket Swamp			
System Terrestrial <input checked="" type="checkbox"/> Wetland Aquatic		Topographic Feature Lacustrine Riverine <input checked="" type="checkbox"/> Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff				Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte <input checked="" type="checkbox"/> Deciduous Coniferous Mixed			
Cover Open <input checked="" type="checkbox"/> Shrub <input type="checkbox"/> Treed		History <input checked="" type="checkbox"/> Natural Cultural		Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest <input checked="" type="checkbox"/> Thicket Cultural <input checked="" type="checkbox"/> Swamp Fen Bog Marsh Open Water Shallow Water					
Stand Description:					Soil Analysis:				
Community Age Pioneer Young <input checked="" type="checkbox"/> Mid-Aged Mature Old Growth				Basal Area (m²/ha)		Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect <input checked="" type="checkbox"/> Poor Very Poor			
Standing Snags Rare <input checked="" type="checkbox"/> Occasional Abundant Dominant					Soil Moisture Regime Dry Fresh <input checked="" type="checkbox"/> Moist Wet				
Deadfall Logs Rare Occasional <input checked="" type="checkbox"/> Abundant Dominant					Effective Soil Texture Organic				
Health Low <input checked="" type="checkbox"/> Medium High		Sensitivity Low <input checked="" type="checkbox"/> Medium High		Botanical Quality Low <input checked="" type="checkbox"/> Medium High		Depth to Mottles / Gley Sample: M - cm / G - cm			
Slope none <input checked="" type="checkbox"/> gentle moderate steep (simple or complex)					Depth to Groundwater metres at surface <input checked="" type="checkbox"/> less than 1m more than 1 m		Depth to Bedrock metres at surface less than 1m more than 1 m		

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	3	2	Betula papyrifera > Populus tremuloides = Populus balsamifera
2 Subcanopy	4	4	Cornus stolonifera = Salix discolor = Salix petiolaris
3 Understorey	5	3	Cornus stolonifera = Salix sp. > Rubus idaeus ssp. strigosus > Scirpus cyperinus
4 Ground Layer	6	3	Carex lacustris > Onoclea sensibilis = Sedge sp.

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4 = >60%

Size Class Analysis ³	D	O	O	NA
	³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH

Evidence of Disturbance:

Wildlife / Habitat Observations:

Comments: Salix petiolaris dominated community

Community Name				Code	% Coverage
Inclusion		Complex			
Inclusion		Complex			

ELC Community Description & Classification

Polygon: P28


Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 24, 2015

Representative Photographs of Vegetation Community:



ELC Community Description & Classification

Polygon: P29

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 24, 2015

Polygon Description P29		Community Series: SW		Ecosite: SWD		Vegetation Type: SWDM4-5 Poplar Mineral Deciduous Swamp			
System Terrestrial Wetland Aquatic		Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff				Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed			
Cover Open Shrub Treed		History Natural Cultural		Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water					
Stand Description:					Soil Analysis:				
Community Age Pioneer Young Mid-Aged Mature Old Growth				Basal Area (m²/ha)		Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor			
Standing Snags Rare Occasional Abundant Dominant				Soil Moisture Regime Dry Fresh Moist Wet					
Deadfall Logs Rare Occasional Abundant Dominant				Effective Soil Texture					
Health Low Medium High		Sensitivity Low Medium High		Botanical Quality Low Medium High		Depth to Mottles / Gley Sample: M - cm / G - cm			
Slope none gentle moderate steep (simple or complex)				Depth to Groundwater metres at surface less than 1m more than 1m			Depth to Bedrock metres at surface less than 1m more than 1m		

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	3	Populus tremuloides > Acer saccharum ssp. saccharum > Fraxinus pennsylvanica
2 Subcanopy	3	2	Populus tremuloides > Acer saccharum ssp. saccharum > Fraxinus pennsylvanica > Thuja occidentalis
3 Understorey	4	2	Acer negundo > Cornus alternifolia > Rubus idaeus ssp. strigosus > Vitis riparia
4 Ground Layer	5	2	Impatiens capensis > Alliaria petiolate > Solidago canadensis

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7= < 0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³	O	A	R	R
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance: Residence, driveway, lawn, planted species.

Wildlife / Habitat Observations:

Comments: Road side survey.
Complex of Poplar Mineral Deciduous Swamp and Fresh – Moist Sugar Maple Deciduous Forest. Boundary between community and surrounding community(ies) is unclear.

				Community Name	Code	% Coverage
Inclusion		Complex	X	Fresh – Moist Sugar Maple Deciduous Forest	FODM6	40
Inclusion		Complex				

ELC Community Description & Classification

Polygon: P30

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 25, 2015

Polygon Description P30		Community Series: FO		Ecosite: FOD		Vegetation Type: FODM7-7 Fresh – Moist Manitoba Maple Lowland Deciduous Forest			
System Terrestrial Wetland Aquatic		Topographic Feature Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff				Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid Forb Lichen Bryophyte Deciduous Coniferous Mixed			
Cover Open Shrub Treed		History Natural Cultural		Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog Marsh Open Water Shallow Water					
Stand Description:					Soil Analysis:				
Community Age Pioneer Young Mid-Aged Mature Old Growth				Basal Area (m²/ha)		Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect Poor Very Poor			
Standing Snags Rare Occasional Abundant Dominant					Soil Moisture Regime Dry Fresh Moist Wet				
Deadfall Logs Rare Occasional Abundant Dominant					Effective Soil Texture Mineral				
Health Low Medium High		Sensitivity Low Medium High		Botanical Quality Low Medium High		Depth to Mottles / Gley Sample: M - cm / G - cm			
Slope none gentle moderate steep (simple or complex)					Depth to Groundwater metres at surface less than 1m more than 1 m		Depth to Bedrock metres at surface less than 1m more than 1 m		

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	3	3	Acer negundo > Salix fragilis
2 Subcanopy	4	2	Acer negundo > Salix fragilis
3 Understorey	5	3	Prunus virginiana > Cornus stolonifera
4 Ground Layer	6-7	4	Impatiens capensis > Alliaria petiolate > Dactylis glomerata > Solidago canadensis

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7=<0.2m. ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4= >60%

Size Class Analysis ³	A	O	R	NA
	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant

Evidence of Disturbance: Garbage and trash in community and in river. Trails along river and creek.

Wildlife / Habitat Observations:

Comments: Highly disturbed, cultural origin community.

			Community Name	Code	% Coverage
Inclusion		Complex			
Inclusion		Complex			

ELC Community Description & Classification

Polygon: P30

Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 25, 2015

Representative Photographs of Vegetation Community:



DR

ELC Community Description & Classification

Polygon: P31

Project No: 12-137A		Project Name: Hillsburgh Dam EA		Surveyor(s): RH		Date: May 21; July 30; Sept 25, 2015	
Polygon Description P31		Community Series: MA		Ecosite: MAM		Vegetation Type: MAMM1-1 Cattail Graminoid Mineral Meadow Marsh Type	
System Terrestrial <input checked="" type="checkbox"/> Wetland Aquatic		Topographic Feature <input checked="" type="checkbox"/> Lacustrine Riverine Bottomland Terrace Valley slope Tableland Rolling upland Cliff Talus Crevice Cave Alvar Rockland Beach Bar Sand dune Bluff				Dominant Plant Form Plankton Submerged Floating-lvd. Graminoid <input checked="" type="checkbox"/> Forb Lichen Bryophyte Deciduous Coniferous Mixed	
Cover <input checked="" type="checkbox"/> Open Shrub Treed		History <input checked="" type="checkbox"/> Natural Cultural		Community Class Beach-Bar Sand Dune Bluff Cliff Talus Alvar Rock Barren Crevice-Cave Sand Barren Tallgrass Prairie Savannah Woodland Forest Thicket Cultural Swamp Fen Bog <input checked="" type="checkbox"/> Marsh Open Water Shallow Water			
Stand Description:				Soil Analysis:			
Community Age Pioneer <input checked="" type="checkbox"/> Young Mid-Aged Mature Old Growth			Basal Area (m²/ha)		Soil Drainage Very Rapid Rapid Well Moderately Well Imperfect <input checked="" type="checkbox"/> Poor Very Poor		
Standing Snags <input checked="" type="checkbox"/> Rare Occasional Abundant Dominant				Soil Moisture Regime Dry Fresh Moist <input checked="" type="checkbox"/> Wet			
Deadfall Logs Rare <input checked="" type="checkbox"/> Occasional Abundant Dominant				Effective Soil Texture Mineral			
Health Low <input checked="" type="checkbox"/> Medium High		Sensitivity Low <input checked="" type="checkbox"/> Medium High		Botanical Quality <input checked="" type="checkbox"/> Low Medium High		Depth to Mottles / Gley Sample: M - cm / G - cm	
Slope <input checked="" type="checkbox"/> none gentle moderate steep (simple or complex)				Depth to Groundwater metres <input checked="" type="checkbox"/> at surface less than 1m more than 1m		Depth to Bedrock metres at surface less than 1m more than 1m	

Vegetation Layer	Height ¹	Cover ²	Dominant Species per Vegetation Layer
1 Canopy	2	1	Ulmus americana
2 Subcanopy	3	2	Acer negundo > Ulmus americana > Cornus stolonifera
3 Understorey	4	4	Typha angustifolia > Phalaris arundinacea = Agrostis gigantea > Symphyotrichum puniceum
4 Ground Layer	5 - 6	4	Impatiens capensis > Verbena hastata

¹ Height Code: 1=>20m, 2=10m-20m, 3=2m-10m, 4=1m-2m, 5=0.5m-1m, 6=0.2m-0.5m, 7= < 0.2m ² Cover Codes: 0 = none, 1 = 0%- 10%, 2 = 10%- 25%, 3 = 25%-60%, 4 = >60%

Size Class Analysis ³	O	O	NA	NA
³ Abundance Code: RS=Rare, O=Occasional, A=Abundant, D=Dominant	< 10 cm DBH	10 to 24 cm DBH	25 to 50 cm DBH	> 50 cm DBH

Evidence of Disturbance:
Wildlife / Habitat Observations:
Comments:

			Community Name	Code	% Coverage
Inclusion		Complex			
Inclusion		Complex			

ELC Community Description & Classification

Polygon: P31


Project No: 12-137A

Project Name: Hillsburgh Dam EA

Surveyor(s): RH

Date: May 21; July 30; Sept 25, 2015

Representative Photographs of Vegetation Community:



APPENDIX 7. BOTANICAL INVENTORY (About Associate Inc. and CVC data)
 Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

Project: AA12-137A

AA ¹	CVC ²	PLANT TYPE ³	COMMON NAME	SCIENTIFIC NAME	FAMILY	CC ⁴	CW ⁵	SARO ⁶	SARA ⁷	S-Rank ⁸	G-Rank ⁹	CVC 2010 ¹⁰	Wellington County ¹¹
X		FE	Bracken Fern	<i>Pteridium aquilinum</i>	Dennstaedtiaceae	0	3	NL	NL	S5	G5T	4	
X	X	FE	Lady Fern	<i>Athyrium filix-femina var. angustum</i>	Dryopteridaceae	4	0	NL	NL	S5	G5	4	
X	X	FE	Bulblet Fern	<i>Cystopteris bulbifera</i>	Dryopteridaceae	5	-2	NL	NL	S5	G5	4	
X	X	FE	Spinulose Shield Fern	<i>Dryopteris carthusiana</i>	Dryopteridaceae	5	-2	NL	NL	S5	G5	4	
X	X	FE	Crested Shield-fern	<i>Dryopteris cristata</i>	Dryopteridaceae	7	-5	NL	NL	S5	G5	3	
X	X	FE	Evergreen Woodfern	<i>Dryopteris intermedia</i>	Dryopteridaceae	5	0	NL	NL	S5	G5	4	
X	X	FE	Marginal Wood-fern	<i>Dryopteris marginalis</i>	Dryopteridaceae	5	3	NL	NL	S5	G5	4	
X	X	FE	Oak Fern	<i>Gymnocarpium dryopteris</i>	Dryopteridaceae	7	0	NL	NL	S5	G5	3	
X		FE	Ostrich Fern	<i>Matteuccia struthiopteris</i>	Dryopteridaceae	3	0	NL	NL	S5	G5	4	
X	X	FE	Sensitive Fern	<i>Onoclea sensibilis</i>	Dryopteridaceae	4	-3	NL	NL	S5	G5	4	
X	X	FE	Field Horsetail	<i>Equisetum arvense</i>	Equisetaceae	0	0	NL	NL	S5	G5	4	
X	X	FE	Water Horsetail	<i>Equisetum fluviatile</i>	Equisetaceae	7	-5	NL	NL	S5	G5	2	
X	X	FE	Marsh Horsetail	<i>Equisetum palustre</i>	Equisetaceae	10	-3	NL	NL	S5	G5	2	✓
X	X	FE	Dwarf Scouring Rush	<i>Equisetum scirpoides</i>	Equisetaceae	7	-1	NL	NL	S5	G5	2	
	X	FE	Variegated Horsetail	<i>Equisetum variegatum</i>	Equisetaceae	6	-3	NL	NL	S5	G5T	2	✓
X	X	FE	Cinnamon Fern	<i>Osmundastrum cinnamomeum</i>	Osmundaceae	7	-3	NL	NL	S5	G5	3	
	X	FE	Hidden Spike-moss	<i>Selaginella eclipes</i>	Selaginellaceae	7	-4	NL	NL	S4	G4	2	
	X	FE	Northern Beech Fern	<i>Phegopteris connectilis</i>	Thelypteridaceae	8	5	NL	NL	S5	G5	2	
X		FE	New York Fern	<i>Thelypteris noveboracensis</i>	Thelypteridaceae	7	-1	NL	NL	S4S5	G5	2	
X	X	FE	Marsh Fern	<i>Thelypteris palustris</i>	Thelypteridaceae	2	-4	NL	NL	S5	G5T?	4	
X		FO	American Water-plantain	<i>Alisma subcordatum</i>	Alismataceae	1	-5	NL	NL	S4?	G4G5	4	
X	X	FO	Broadleaf Arrowhead	<i>Sagittaria latifolia</i>	Alismataceae	4	-5	NL	NL	S5	G5	3	
X		FO	Goutweed	<i>Aegopodium podagraria</i>	Apiaceae	0	0	NL	NL	SE5	G?	5	
X	X	FO	Bulb-bearing Water-hemlock	<i>Cicuta bulbifera</i>	Apiaceae	5	-5	NL	NL	S5	G5	3	
X	X	FO	Wild Carrot	<i>Daucus carota</i>	Apiaceae	0	5	NL	NL	SE5	G?	5	
	X	FO	American Water-pennywort	<i>Hydrocotyle americana</i>	Apiaceae	7	-5	NL	NL	S5	G5	2	
X		FO	Spreading Dogbane	<i>Apocynum androsaemifolium</i>	Apocynaceae	3	5	NL	NL	S5	G5T?	4	
X	X	FO	Periwinkle	<i>Vinca minor</i>	Apocynaceae	0	5	NL	NL	SE5	G?	5	
X	X	FO	Jack-in-the-pulpit	<i>Arisaema triphyllum</i>	Araceae	5	-2	NL	NL	S5	G5T5	4	
	X	FO	Wild Calla	<i>Calla palustris</i>	Araceae	8	-5	NL	NL	S5	G5	2	
X	X	FO	Wild Sarsaparilla	<i>Aralia nudicaulis</i>	Araliaceae	4	3	NL	NL	S5	G5	4	
	X	FO	American Spikenard	<i>Aralia racemosa</i>	Araliaceae	8	3	NL	NL	S5	G5T?	2	
X	X	FO	Canada Wild-ginger	<i>Asarum canadense</i>	Aristolochiaceae	6	5	NL	NL	S5	G5	4	
X	X	FO	Common Milkweed	<i>Asclepias syriaca</i>	Asclepiadaceae	0	5	NL	NL	S5	G5	4	
X	X	FO	Yarrow	<i>Achillea millefolium</i>	Asteraceae	1	3	NL	NL	SE?	G5T?	5	
	X	FO	White Snakeroot	<i>Ageratina altissima</i>	Asteraceae	4	3	NL	NL	S5	G5	4	
X	X	FO	Annual Ragweed	<i>Ambrosia artemisiifolia</i>	Asteraceae	0	3	NL	NL	S5	G5	4	
X		FO	Great Ragweed	<i>Ambrosia trifida</i>	Asteraceae	0	-1	NL	NL	S5	G5	4	
	X	FO	Pearly Everlasting	<i>Anaphalis margaritacea</i>	Asteraceae	3	5	NL	NL	S5	G5	3	
X		FO	Greater Burdock	<i>Arctium lappa</i>	Asteraceae	0	5	NL	NL	SE5	G?	5	
X	X	FO	Lesser Burdock	<i>Arctium minus</i>	Asteraceae	0	5	NL	NL	SE5	G?T?	5	
X	X	FO	Nodding Beggar-ticks	<i>Bidens cernua</i>	Asteraceae	2	-5	NL	NL	S5	G5	4	
X		FO	Devil's Beggar-ticks	<i>Bidens frondosa</i>	Asteraceae	3	-3	NL	NL	S5	G5	4	
X	X	FO	Beggar-ticks	<i>Bidens tripartita</i>	Asteraceae	4	-3	NL	NL	S5	G5	3	
X		FO	Chicory	<i>Cichorium intybus</i>	Asteraceae	0	5	NL	NL	SE5	G?	5	
X	X	FO	Canada Thistle	<i>Cirsium arvense</i>	Asteraceae	0	3	NL	NL	SE5	G?	5	
	X	FO	Bull Thistle	<i>Cirsium vulgare</i>	Asteraceae	0	4	NL	NL	SE5	G5	5	

APPENDIX 7. BOTANICAL INVENTORY (About Associate Inc. and CVC data)
 Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

Project: AA12-137A

AA ¹	CVC ²	PLANT TYPE ³	COMMON NAME	SCIENTIFIC NAME	FAMILY	CC ⁴	CW ⁵	SARO ⁶	SARA ⁷	S-Rank ⁸	G-Rank ⁹	CVC 2010 ¹⁰	Wellington County ¹¹
X	X	FO	Annual Fleabane	<i>Erigeron annuus</i>	Asteraceae	0	1	NL	NL	S5	G5	4	
X	X	FO	Philadelphia Fleabane	<i>Erigeron philadelphicus</i>	Asteraceae	2	-3	NL	NL	S5	G5T?	4	
X	X	FO	Common Boneset	<i>Eupatorium perfoliatum</i>	Asteraceae	2	-4	NL	NL	S5	G5	4	
X		FO	Large-leaf Wood-aster	<i>Eurybia macrophylla</i>	Asteraceae	5	5	NL	NL	S5	G5	4	
X	X	FO	Grass-leaved goldenrod	<i>Euthamia graminifolia</i>	Asteraceae	2	-2	NL	NL	S5	G5	4	
	X	FO	Spotted Joe-pye Weed	<i>Eutrochium maculatum</i> var. <i>foliosum</i>	Asteraceae			NL	NL	S5		4	
X		FO	Spotted Joe-pye Weed	<i>Eutrochium maculatum</i> var. <i>maculatum</i>	Asteraceae	4	-5	NL	NL	S5	G5T5	4	
	X	FO	Elecampane Flower	<i>Inula helenium</i>	Asteraceae	0	5	NL	NL	SE5	G?	5	
X		FO	Canada Lettuce	<i>Lactuca canadensis</i>	Asteraceae	3	2	NL	NL	S5	G5	4	
	X	FO	Common Nipplewort	<i>Lapsana communis</i>	Asteraceae	0	5	NL	NL	SE5	G?	5	
X	X	FO	Oxeye Daisy	<i>Leucanthemum vulgare</i>	Asteraceae	0	5	NL	NL	SE5	G?	5	
	X	FO	Tall Rattlesnake-root	<i>Nabalus altissimus</i>	Asteraceae	5	3	NL	NL	S5	G5?	3	
	X	FO	Golden Ragwort	<i>Packera aurea</i>	Asteraceae	7	-3	NL	NL	S5	G5	2	
	X	FO	Hawkweed Oxtongue	<i>Picris hieracioides</i>	Asteraceae	0	5	NL	NL	SE5	G5T?	5	
X		FO	Meadow Hawkweed	<i>Pilosella caespitosa</i>	Asteraceae	0	5	NL	NL	SE		5	
X	X	FO	Black-eyed Susan	<i>Rudbeckia hirta</i> var. <i>hirta</i>	Asteraceae	1	3	NL	NL	S5	G5	4	
X	X	FO	Tall Goldenrod	<i>Solidago altissima</i>	Asteraceae	1	3	NL	NL	S5	G5T5	4	
X	X	FO	Canada Goldenrod	<i>Solidago canadensis</i> var. <i>canadensis</i>	Asteraceae	1	3	NL	NL	S5		4	
X	X	FO	Broad-leaved Goldenrod	<i>Solidago flexicaulis</i>	Asteraceae	6	3	NL	NL	S5	G5	4	
X	X	FO	Roughleaf Goldenrod	<i>Solidago rugosa</i> var. <i>rugosa</i>	Asteraceae	3	-1	NL	NL	S5	G5T?	3	
X	X	FO	Bog Goldenrod	<i>Solidago uliginosa</i>	Asteraceae	9	-5	NL	NL	S5	G4G5	2	
X	X	FO	Field Sow-thistle	<i>Sonchus arvensis</i> ssp. <i>arvensis</i>	Asteraceae	0	1	NL	NL	SE5	G?	5	
X		FO	White Heath Aster	<i>Symphotrichum ericoides</i> var. <i>ericoides</i>	Asteraceae	4	4	NL	NL	S5	G5	4	
X	X	FO	Panicked Aster	<i>Symphotrichum lanceolatum</i> ssp. <i>lanceolatum</i>	Asteraceae	3	-3	NL	NL	S5	G5	4	
X	X	FO	Calico Aster	<i>Symphotrichum lateriflorum</i>	Asteraceae	2	0	NL	NL	S5		3	
X	X	FO	New England Aster	<i>Symphotrichum novae-angliae</i>	Asteraceae	2	-3	NL	NL	S5	G5	4	
X	X	FO	Purple-stemmed Aster	<i>Symphotrichum puniceum</i>	Asteraceae	5	-5	NL	NL	S5		4	
X		FO	Common Tansy	<i>Tanacetum vulgare</i>	Asteraceae	0	5	NL	NL	SE5	G?	5	
X	X	FO	Brown-seed Dandelion	<i>Taraxacum officinale</i>	Asteraceae	0	3	NL	NL	SE5	G5	5	
X		FO	Meadow Goat's-beard	<i>Tragopogon dubius</i>	Asteraceae	0	5	NL	NL	SE5	G?	5	
X	X	FO	Colt's Foot	<i>Tussilago farfara</i>	Asteraceae	0	3	NL	NL	SE5	G?	5	
X	X	FO	Spotted Jewel-weed	<i>Impatiens capensis</i>	Balsaminaceae	4	-3	NL	NL	S5	G5	4	
X		FO	Pale Jewel-weed	<i>Impatiens pallida</i>	Balsaminaceae	7	-3	NL	NL	S5	G5	2	✓
X	X	FO	Giant Blue Cohosh	<i>Caulophyllum thalictroides</i>	Berberidaceae	6	5	NL	NL	S4?	G?	3	
X	X	FO	May Apple	<i>Podophyllum peltatum</i>	Berberidaceae	5	3	NL	NL	S5	G5	4	
X		FO	Common Viper's-bugloss	<i>Echium vulgare</i>	Boraginaceae	0	5	NL	NL	SE5	G?	5	
X	X	FO	Small Forget-me-not	<i>Myosotis laxa</i>	Boraginaceae	6	-5	NL	NL	S5	G5	3	
X		FO	True Forget-me-not	<i>Myosotis scorpioides</i>	Boraginaceae	0	-5	NL	NL	SE5	G5	5	
X	X	FO	Garlic Mustard	<i>Alliaria petiolata</i>	Brassicaceae	0	0	NL	NL	SE5	G?	5	
X		FO	Yellow Rocket	<i>Barbarea vulgaris</i>	Brassicaceae	0	0	NL	NL	SE5	G?	5	
X	X	FO	Two-leaf Toothwort	<i>Cardamine diphylla</i>	Brassicaceae	7	5	NL	NL	S5	G5	3	
X		FO	Dame's Rocket	<i>Hesperis matronalis</i>	Brassicaceae	0	5	NL	NL	SE5	G4G5	5	
X	X	FO	Small-leaved Watercress	<i>Nasturtium microphyllum</i>	Brassicaceae	0	-5	NL	NL	SE5	G?	5	

Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

AA ¹	CVC ²	PLANT TYPE ³	COMMON NAME	SCIENTIFIC NAME	FAMILY	CC ⁴	CW ⁵	SARO ⁶	SARA ⁷	S-Rank ⁸	G-Rank ⁹	CVC 2010 ¹⁰	Wellington County ¹¹
X	X	FO	Marsh Bellflower	<i>Campanula aparinoides</i>	Campanulaceae	7	-5	NL	NL	S5	G5	2	
X	X	FO	Kalm's Lobelia	<i>Lobelia kalmii</i>	Campanulaceae	9	-5	NL	NL	S5	G5	2	✓
X		FO	Common Mouse-ear Chickweed	<i>Cerastium fontanum</i>	Caryophyllaceae	0	3	NL	NL	SE5	G?	5	
X		FO	Bouncing-bet	<i>Saponaria officinalis</i>	Caryophyllaceae	0	3	NL	NL	SE5	G?	5	
X		FO	A Catchfly	<i>Silene latifolia</i>	Caryophyllaceae	0	5	NL	NL	SE5	G?	5	
X		FO	Maiden's Tears	<i>Silene vulgaris</i>	Caryophyllaceae	0	5	NL	NL	SE5	G?	5	
X		FO	Common Lamb's-quarters	<i>Chenopodium album</i>	Chenopodiaceae	0	1	NL	NL	SE5	G5	5	
X	X	FO	St. John's-wort	<i>Hypericum perforatum</i>	Clusiaceae	0	5	NL	NL	SE5	G?	5	
X	X	FO	Common St. John's-wort	<i>Hypericum punctatum</i>	Clusiaceae	5	-1	NL	NL	S5	G5	2	
	X	FO	Marsh St. John's-wort	<i>Triadenum fraseri</i>	Clusiaceae	7	-5	NL	NL	S5	G4G5	2	
X		FO	Teasel	<i>Dipsacus fullonum</i>	Dipsacaceae	0	3	NL	NL	SE5	G?T?	5	
	X	FO	Roundleaf Sundew	<i>Drosera rotundifolia</i>	Droseraceae	7	-5	NL	NL	S5	G5	2	
X		FO	Birds-foot Trefoil	<i>Lotus corniculatus</i>	Fabaceae	0	1	NL	NL	SE5	G?	5	
X	X	FO	Black Medic	<i>Medicago lupulina</i>	Fabaceae	0	1	NL	NL	SE5	G?	5	
X		FO	White Sweet Clover	<i>Melilotus albus</i>	Fabaceae	0	3	NL	NL	SE5	G5	5	
X		FO	Yellow Sweetclover	<i>Melilotus officinalis</i>	Fabaceae	0	3	NL	NL	SE5	G?	5	
X	X	FO	Rabbit-foot Clover	<i>Trifolium arvense</i>	Fabaceae	0	5	NL	NL	SE4	G?	5	
X		FO	Red Clover	<i>Trifolium pratense</i>	Fabaceae	0	2	NL	NL	SE5	G?	5	
X	X	FO	White Clover	<i>Trifolium repens</i>	Fabaceae	0	2	NL	NL	SE5	G?	5	
X	X	FO	Herb-robert	<i>Geranium robertianum</i>	Geraniaceae	0	5	NL	NL	SE5	G5	5	
X		FO	Eel-grass	<i>Vallisneria americana</i>	Hydrocharitaceae	6	-5	NL	NL	S5	G5	2	
X	X	FO	Virginia waterleaf	<i>Hydrophyllum virginianum</i>	Hydrophyllaceae	6	-2	NL	NL	S5	G5	4	
X		FO	Yellow Iris	<i>Iris pseudacorus</i>	Iridaceae	0	-5	NL	NL	SE3	G?	5	
X	X	FO	Blueflag	<i>Iris versicolor</i>	Iridaceae	5	-5	NL	NL	S5	G5	3	
X		FO	Strict Blue-eyed-grass	<i>Sisyrinchium montanum</i>	Iridaceae	4	-1	NL	NL	S5	G5	2	
	X	FO	Field Basil	<i>Clinopodium vulgare</i>	Lamiaceae	4	5	NL	NL	S5	G?	4	
X	X	FO	Ground Ivy	<i>Glechoma hederacea</i>	Lamiaceae	0	3	NL	NL	SE5	G?	5	
X		FO	Common Mother-wort	<i>Leonurus cardiaca</i>	Lamiaceae	0	5	NL	NL	SE5	G?T?	5	
X	X	FO	American Bugleweed	<i>Lycopus americanus</i>	Lamiaceae	4	-5	NL	NL	S5	G5	3	
X	X	FO	Northern Bugleweed	<i>Lycopus uniflorus</i>	Lamiaceae	5	-5	NL	NL	S5	G5	4	
	X	FO	Corn Mint	<i>Mentha canadensis</i>	Lamiaceae	3	-3	NL	NL	S5		4	
	X	FO	Spearmint	<i>Mentha spicata</i>	Lamiaceae	0	-4	NL	NL	SE4	G?	5	
X	X	FO	Peppermint	<i>Mentha x piperita</i>	Lamiaceae	0	-5	NL	NL	SE4	G?	5	
X		FO	Catnip	<i>Nepeta cataria</i>	Lamiaceae	0	1	NL	NL	SE5	G?	5	
X	X	FO	Self-heal	<i>Ranunculus acris</i> ssp. <i>lanceolata</i>	Lamiaceae	5	5	NL	NL	S5	G5	4	
X	X	FO	Hooded Skullcap	<i>Scutellaria galericulata</i>	Lamiaceae	6	-5	NL	NL	S5	G5	3	
	X	FO	Mad Dog Skullcap	<i>Scutellaria lateriflora</i>	Lamiaceae	5	-5	NL	NL	S5	G5	3	
X	X	FO	Lesser Duckweed	<i>Lemna minor</i>	Lemnaceae	2	-5	NL	NL	S5	G5	4	
	X	FO	Lesser Bladderwort	<i>Utricularia minor</i>	Lentibulariaceae	8	-5	NL	NL	S5	G5	2	
X		FO	Wild Leek	<i>Allium tricoccum</i> var. <i>tricoccum</i>	Liliaceae	5	2	NL	NL	S4	G5	3	
X		FO	Garden Asparagus-fern	<i>Asparagus officinalis</i>	Liliaceae	0	3	NL	NL	SE5	G5?	5	
X	X	FO	Blue Bead-lily	<i>Clintonia borealis</i>	Liliaceae	7	-1	NL	NL	S5	G5	3	
	X	FO	European Lily-of-the-valley	<i>Convallaria majalis</i>	Liliaceae	0	5	NL	NL	SE5	G5	5	
X	X	FO	Yellow Trout-lily	<i>Erythronium americanum</i>	Liliaceae	5	5	NL	NL	S5	G5T5	4	
X		FO	Orange Daylily	<i>Hemerocallis fulva</i>	Liliaceae	0	5	NL	NL	SE5	G?	5	
X	X	FO	Wild-lily-of-the-valley	<i>Maianthemum canadense</i>	Liliaceae	5	0	NL	NL	S5	G5	3	

APPENDIX 7. BOTANICAL INVENTORY (About Associate Inc. and CVC data)
Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

Project: AA12-137A

AA ¹	CVC ²	PLANT TYPE ³	COMMON NAME	SCIENTIFIC NAME	FAMILY	CC ⁴	CW ⁵	SARO ⁶	SARA ⁷	S-Rank ⁸	G-Rank ⁹	CVC 2010 ¹⁰	Wellington County ¹¹
X	X	FO	False Solomon's-seal	<i>Maianthemum racemosum</i>	Liliaceae	5	3	NL	NL	S5	G5T	3	
X	X	FO	Starflower False Solomon's-seal	<i>Maianthemum stellatum</i>	Liliaceae	6	1	NL	NL	S5	G5	4	
X		FO	Common Daffodil	<i>Narcissus pseudonarcissus</i>	Liliaceae	0	0	NL	NL	SE2	G?	5	
	X	FO	Downy Solomon's-seal	<i>Polygonatum pubescens</i>	Liliaceae	5	5	NL	NL	S5	G5	4	
	X	FO	Rose Twisted-stalk	<i>Streptopus lanceolatus</i>	Liliaceae	7	0	NL	NL	S5	G5	2	
X	X	FO	Red Trillium	<i>Trillium erectum</i>	Liliaceae	6	1	NL	NL	S5	G5	3	
X	X	FO	White Trillium	<i>Trillium grandiflorum</i>	Liliaceae	5	5	NL	NL	S5	G5	4	
X	X	FO	Purple Loosestrife	<i>Lythrum salicaria</i>	Lythraceae	0	-5	NL	NL	SE5	G5	5	
X		FO	Velvet-leaf	<i>Abutilon theophrasti</i>	Malvaceae	0	4	NL	NL	SE5	G?	5	
X		FO	Musk Mallow	<i>Malva moschata</i>	Malvaceae	0	5	NL	NL	SE5	G?	5	
X		FO	Indian-pipe	<i>Monotropa uniflora</i>	Monotropaceae	6	3	NL	NL	S5	G5	3	
	X	FO	Slender Naiad	<i>Najas flexilis</i>	Najadaceae	5	-5	NL	NL	S5	G5	2	✓
X		FO	Yellow Cowwily	<i>Nuphar variegata</i>	Nymphaeaceae	4	-5	NL	NL	S5	G5	2	
X	X	FO	White Water-lily	<i>Nymphaea odorata ssp.</i>	Nymphaeaceae	5	-5	NL	NL	S5?	G5	2	
	X	FO	Tuberous White Water-lily	<i>Nymphaea odorata ssp. tuberosa</i>	Nymphaeaceae	5	-5	NL	NL	SU	G5	2	
X		FO	Fireweed	<i>Chamerion angustifolium</i>	Onagraceae	3	0	NL	NL	S5	G5	2	✓
X	X	FO	Small Enchanter's Nightshade	<i>Circaea alpina</i>	Onagraceae	6	-3	NL	NL	S5	G5	3	
X	X	FO	Broad-leaved Enchanter's Nightshade	<i>Circaea canadensis</i>	Onagraceae	3	3	NL	NL	S5	G5	4	
X		FO	Hairy Willow-herb	<i>Epilobium ciliatum</i>	Onagraceae	3	-3	NL	NL	S5	G5T?	4	
X	X	FO	Purple-leaf Willow-herb	<i>Epilobium coloratum</i>	Onagraceae	3	-5	NL	NL	S5	G5	2	
X	X	FO	Great-hairy Willow-herb	<i>Epilobium hirsutum</i>	Onagraceae	0	-4	NL	NL	SE5	G?	5	
X	X	FO	Linear-leaved Willow-herb	<i>Epilobium leptophyllum</i>	Onagraceae	7	-5	NL	NL	S5	G5	2	
X	X	FO	Small-flower Willow-herb	<i>Epilobium parviflorum</i>	Onagraceae	0	3	NL	NL	SE4	G?	5	
X	X	FO	Common Evening-primrose	<i>Oenothera biennis</i>	Onagraceae	0	3	NL	NL	S5	G5	4	
	X	FO	Small Yellow Lady's-slipper	<i>Cypripedium parviflorum var. makasin</i>	Orchidaceae	5	0	NL	NL	S5	G5T	2	
	X	FO	Large Yellow Lady's-slipper	<i>Cypripedium parviflorum var. pubescens</i>	Orchidaceae	5	-1	NL	NL	S5	G5	2	
	X	FO	Showy Lady's-slipper	<i>Cypripedium reginae</i>	Orchidaceae	7	-4	NL	NL	S4	G4	2	
X	X	FO	Eastern Helleborine	<i>Epipactis helleborine</i>	Orchidaceae	0	5	NL	NL	SE5	G?	5	
X	X	FO	Loesel's Twayblade	<i>Liparis loeselii</i>	Orchidaceae	5	-4	NL	NL	S4S5	G5	2	
	X	FO	Leafy Northern Green Orchid	<i>Platanthera aquilonis</i>	Orchidaceae	5	-4	NL	NL	S5	G5	2	
X	X	FO	Hooded Ladies'-tresses	<i>Spiranthes romanzoffiana</i>	Orchidaceae	9	-4	NL	NL	S5	G5	2	✓
X	X	FO	Common Wood-sorrel	<i>Oxalis montana</i>	Oxalidaceae	8	3	NL	NL	S5	G5	2	
X	X	FO	Upright Yellow Wood-sorrel	<i>Oxalis stricta</i>	Oxalidaceae	0	3	NL	NL	S5	G5	5	
X	X	FO	Greater Celadine	<i>Chelidonium majus</i>	Papaveraceae	0	5	NL	NL	SE5	G?	5	
X		FO	Bloodroot	<i>Sanguinaria canadensis</i>	Papaveraceae	5	4	NL	NL	S5	G5	4	
X	X	FO	English Plantain	<i>Plantago lanceolata</i>	Plantaginaceae	0	0	NL	NL	SE5	G5	5	
X	X	FO	Common Plantain	<i>Plantago major</i>	Plantaginaceae	0	-1	NL	NL	SE5	G5	5	
X		FO	Fall Phlox	<i>Phlox paniculata</i>	Polemoniaceae	0	3	NL	NL	SE3	G5	5	
	X	FO	Water Smartweed	<i>Persicaria amphibia</i>	Polygonaceae	5	-5	NL	NL	S5	G5	2	
X	X	FO	Marshpepper Smartweed	<i>Persicaria hydropiper</i>	Polygonaceae	0	-5	NL	NL	SE5	G5	5	
X	X	FO	Lady's Thumb	<i>Persicaria maculosa</i>	Polygonaceae	0	-3	NL	NL	SE5	G?	5	
	X	FO	Pennsylvania Smartweed	<i>Persicaria pennsylvanica</i>	Polygonaceae	3	-4	NL	NL	S5	G5	2	
X	X	FO	Curly Dock	<i>Rumex crispus</i>	Polygonaceae	0	-1	NL	NL	SE5	G?	5	

Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

AA ¹	CVC ²	PLANT TYPE ³	COMMON NAME	SCIENTIFIC NAME	FAMILY	CC ⁴	CW ⁵	SARO ⁶	SARA ⁷	S-Rank ⁸	G-Rank ⁹	CVC 2010 ¹⁰	Wellington County ¹¹
X	X	FO	Bitter Dock	<i>Rumex obtusifolius</i>	Polygonaceae	0	-3	NL	NL	SE5	G5	5	
X	X	FO	Water Dock	<i>Rumex orbiculatus</i>	Polygonaceae	6	-5	NL	NL	S4S5	G5	2	
	X	FO	Curly Pondweed	<i>Potamogeton crispus</i>	Potamogetonaceae	0	-5	NL	NL	SE5	G5	5	
	X	FO	Leafy Pondweed	<i>Potamogeton foliosus</i>	Potamogetonaceae	4	-5	NL	NL	S5	G5	2	
	X	FO	Berchtold's Pondweed	<i>Potamogeton pusillus ssp. tenuissimus</i>	Potamogetonaceae	4	-5	NL	NL	S4S5	G?	2	
X		FO	Pondweed sp.	<i>Potamogeton sp.</i>	Potamogetonaceae								
	X	FO	Sago Pondweed	<i>Stuckenia pectinata</i>	Potamogetonaceae	4	-5	NL	NL	S5	G5	4	
	X	FO	Fringed Loosestrife	<i>Lysimachia ciliata</i>	Primulaceae	4	-3	NL	NL	S5	G5	4	
X	X	FO	Northern Starflower	<i>Trientalis borealis</i>	Primulaceae	5	-1	NL	NL	S5	G5T?	3	
	X	FO	One-side Wintergreen	<i>Orthilia secunda</i>	Pyrolaceae	5	-1	NL	NL	S5	G5	2	
	X	FO	Pink Wintergreen	<i>Pyrola asarifolia</i>	Pyrolaceae	7	-3	NL	NL	S5	G5	2	
	X	FO	Shinleaf	<i>Pyrola elliptica</i>	Pyrolaceae	5	5	NL	NL	S5	G5	3	
X	X	FO	White Baneberry	<i>Actaea pachypoda</i>	Ranunculaceae	6	5	NL	NL	S5	G5	4	
X	X	FO	Red Baneberry	<i>Actaea rubra</i>	Ranunculaceae	5	5	NL	NL	S5	G5	4	
X	X	FO	Canada Anemone	<i>Anemone canadensis</i>	Ranunculaceae	3	-3	NL	NL	S5	G5	4	
X		FO	Marsh Marigold	<i>Caltha palustris</i>	Ranunculaceae	5	-5	NL	NL	S5	G5	3	
X	X	FO	Goldthread	<i>Coptis trifolia</i>	Ranunculaceae	5	-3	NL	NL	S5	G5T5	3	
X	X	FO	Kidney-leaved Buttercup	<i>Ranunculus abortivus</i>	Ranunculaceae	2	-2	NL	NL	S5	G5	4	
X	X	FO	Tall Butter-cup	<i>Ranunculus acris</i>	Ranunculaceae	0	-2	NL	NL	SE5	G5	5	
X	X	FO	White Water Buttercup	<i>Ranunculus aquatilis</i>	Ranunculaceae		-5	NL	NL	S5	G5T	2	
	X	FO	Bristly Buttercup	<i>Ranunculus hispidus var. caricetorum</i>	Ranunculaceae	5	-5	NL	NL	S5	G5T5	3	
X	X	FO	Hooked Crowfoot	<i>Ranunculus recurvatus</i>	Ranunculaceae	4	-3	NL	NL	S5	G5	3	
	X	FO	Creeping Butter-cup	<i>Ranunculus repens</i>	Ranunculaceae	0	-1	NL	NL	SE5	G?	5	
X	X	FO	Hooked Agrimony	<i>Agrimonia gryposepala</i>	Rosaceae	2	2	NL	NL	S5	G5	4	
X	X	FO	Woodland Strawberry	<i>Fragaria vesca</i>	Rosaceae	2	4	NL	NL	S5	G5T?	3	
X	X	FO	Virginia Strawberry	<i>Fragaria virginiana</i>	Rosaceae	2	1	NL	NL	S5	G5T?	4	
X	X	FO	Yellow Avens	<i>Geum aleppicum</i>	Rosaceae	2	-1	NL	NL	S5	G5	4	
X		FO	White Avens	<i>Geum canadense</i>	Rosaceae	3	0	NL	NL	S5	G5	4	
	X	FO	Purple Avens	<i>Geum rivale</i>	Rosaceae	7	-5	NL	NL	S5	G5	2	
X		FO	Old-field Cinquefoil	<i>Potentilla simplex</i>	Rosaceae	3	4	NL	NL	S5	G5	2	
X	X	FO	Rough Bedstraw	<i>Galium asprellum</i>	Rubiaceae	6	-5	NL	NL	S5	G5	4	
X		FO	Great Hedge Bedstraw	<i>Galium mollugo</i>	Rubiaceae	0	5	NL	NL	SE5	G?	5	
X	X	FO	Marsh Bedstraw	<i>Galium palustre</i>	Rubiaceae	5	-5	NL	NL	S5	G5	3	
	X	FO	Rough-fruit Corn Bedstraw	<i>Galium tricornutum</i>	Rubiaceae			NL	NL	SEH	G?	2	
X	X	FO	Sweet-scent Bedstraw	<i>Galium triflorum</i>	Rubiaceae	4	2	NL	NL	S5	G5	4	
X	X	FO	Partridge-berry	<i>Mitchella repens</i>	Rubiaceae	6	2	NL	NL	S5	G5	2	
	X	FO	American Golden-saxifrage	<i>Chrysosplenium americanum</i>	Saxifragaceae	8	-5	NL	NL	S5	G5	2	✓
	X	FO	Two-leaf Bishop's-cap	<i>Mitella diphylla</i>	Saxifragaceae	5	2	NL	NL	S5	G5	3	
	X	FO	Naked Bishop's-cap	<i>Mitella nuda</i>	Saxifragaceae	6	-3	NL	NL	S5	G5	3	
X	X	FO	Heart-leaved Foam-flower	<i>Tiarella cordifolia</i>	Saxifragaceae	6	1	NL	NL	S5	G5	3	
X	X	FO	White Turtlehead	<i>Chelone glabra</i>	Scrophulariaceae	7	-5	NL	NL	S5	G5	3	
X		FO	Butter-and-eggs	<i>Linaria vulgaris</i>	Scrophulariaceae	0	5	NL	NL	SE5	G?	5	
X		FO	Great Mullein	<i>Verbascum thapsus</i>	Scrophulariaceae	0	5	NL	NL	SE5	G?	5	
	X	FO	American Speedwell	<i>Veronica americana</i>	Scrophulariaceae	6	-5	NL	NL	S5	G5	2	
	X	FO	Brook-pimpernell	<i>Veronica anagallis-aquatica</i>	Scrophulariaceae	0	-5	NL	NL	SE5	G5	5	
	X	FO	Gypsy-weed	<i>Veronica officinalis</i>	Scrophulariaceae	0	5	NL	NL	SE5	G5	5	

APPENDIX 7. BOTANICAL INVENTORY (About Associate Inc. and CVC data)
 Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

Project: AA12-137A

AA ¹	CVC ²	PLANT TYPE ³	COMMON NAME	SCIENTIFIC NAME	FAMILY	CC ⁴	CW ⁵	SARO ⁶	SARA ⁷	S-Rank ⁸	G-Rank ⁹	CVC 2010 ¹⁰	Wellington County ¹¹
X	X	FO	Green-fruited Burreed	<i>Sparganium emersum</i>	Sparganiaceae	6	-5	NL	NL	S5		2	
	X	FO	Small Bur-reed	<i>Sparganium natans</i>	Sparganiaceae	8	-5	NL	NL	S5	G5	2	
X	X	FO	Narrow-leaved Cattail	<i>Typha angustifolia</i>	Typhaceae	3	-5	NL	NL	SE5	G5	5	
X	X	FO	Broad-leaf Cattail	<i>Typha latifolia</i>	Typhaceae	3	-5	NL	NL	S5	G5	4	
X		FO	(Typha angustifolia X Typha latifolia)	<i>Typha x glauca</i>	Typhaceae	3	-5	NL	NL	SE5	G?	5	
	X	FO	Springs Clearweed	<i>Pilea fontana</i>	Urticaceae	5	-3	NL	NL	S4	G5	2	
X		FO	Stinging Nettle	<i>Urtica dioica ssp. gracilis</i>	Urticaceae	2	-1	NL	NL	S5	G5T?	4	
	X	FO	Lopseed	<i>Phryma leptostachya</i>	Verbenaceae	6	5	NL	NL	S4S5	G5	5	
X	X	FO	Blue Vervain	<i>Verbena hastata</i>	Verbenaceae	4	-4	NL	NL	S5	G5	4	
	X	FO	White Vervain	<i>Verbena urticifolia</i>	Verbenaceae	4	-1	NL	NL	S5	G5	3	
X	X	FO	Marsh Blue Violet	<i>Viola cucullata</i>	Violaceae	5	-5	NL	NL	S5	G4G5	2	
X	X	FO	Labrador Violet	<i>Viola labradorica</i>	Violaceae	3	0	NL	NL	S4S5	G5	4	
X	X	FO	Smooth White Violet	<i>Viola maclouskeyi</i>	Violaceae	6	-5	NL	NL	S5	G5T5	2	
X	X	FO	Downy Yellow Violet	<i>Viola pubescens var. pubescens</i>	Violaceae	4	3	NL	NL	S5		4	
X		FO	Smooth Yellow Violet	<i>Viola pubescens var. scabriuscula</i>	Violaceae	4	3	NL	NL	S5		4	
X	X	GR	Redtop	<i>Agrostis gigantea</i>	Poaceae	0	0	NL	NL	SE5	G4G5	5	
	X	GR	Rough Bentgrass	<i>Agrostis scabra</i>	Poaceae	6	0	NL	NL	S5	G5	2	
X	X	GR	Fringed Brome	<i>Bromus ciliatus</i>	Poaceae	6	-3	NL	NL	S5	G5	2	
X		GR	Awnless Brome	<i>Bromus inermis</i>	Poaceae	0	5	NL	NL	SE5	G4G5	5	
X		GR	Brome sp.	<i>Bromus sp.</i>	Poaceae								
X	X	GR	Canada Blue-joint	<i>Calamagrostis canadensis</i>	Poaceae	4	-5	NL	NL	S5	G5	3	
	X	GR	Slender Wood Reedgrass	<i>Cinna latifolia</i>	Poaceae	7	-4	NL	NL	S5	G5	3	
X	X	GR	Orchard Grass	<i>Dactylis glomerata</i>	Poaceae	0	3	NL	NL	SE5	G?	5	
X	X	GR	American Mannagrass	<i>Glyceria grandis</i>	Poaceae	5	-5	NL	NL	S4S5	G5	4	
	X	GR	Reed Meadowgrass	<i>Glyceria maxima</i>	Poaceae	0	-5	NL	NL	SE4	G?	5	
X	X	GR	Fowl Manna-grass	<i>Glyceria striata</i>	Poaceae	3	-5	NL	NL	S4S5	G5	4	
X	X	GR	Rice Cutgrass	<i>Leersia oryzoides</i>	Poaceae	3	-5	NL	NL	S5	G5	4	
X		GR	Giant miscanthus	<i>Miscanthus x giganteus</i>	Poaceae	0		NL	NL	SE		5	
	X	GR	Mexican Muhly	<i>Muhlenbergia mexicana</i>	Poaceae	3	-3	NL	NL	S5	G5T?	2	
X	X	GR	Reed Canary Grass	<i>Phalaris arundinacea</i>	Poaceae	0	-4	NL	NL	S5	G5	4	
X	X	GR	Meadow Timothy	<i>Phleum pratense</i>	Poaceae	0	3	NL	NL	SE5	G?	5	
X	X	GR	European Reed	<i>Phragmites australis ssp. australis</i>	Poaceae	0	-3	NL	NL	S5	G5	5	
	X	GR	Canada Bluegrass	<i>Poa compressa</i>	Poaceae	0	2	NL	NL	SE5	G?	5	
	X	GR	Woods Bluegrass	<i>Poa nemoralis</i>	Poaceae	0	0	NL	NL	SE3	G5	5	
	X	GR	Fowl Bluegrass	<i>Poa palustris</i>	Poaceae	5	-4	NL	NL	S5	G5	3	
	X	GR	Kentucky Bluegrass	<i>Poa pratensis ssp. pratensis</i>	Poaceae	0	1	NL	NL	S5	G?	4	
X		GR	Grass sp.	<i>Poaceae sp.</i>	Poaceae								
	X	GR	Meadow Fescue	<i>Schedonorus pratensis</i>	Poaceae	0	4	NL	NL	SE5	G5	5	
	X	GR	Purple Oat	<i>Schizachne purpurascens</i>	Poaceae	5	3	NL	NL	S5	G5T?	3	
	X	GR	Slender Wedge Grass	<i>Sphenopholis intermedia</i>	Poaceae	6	0	NL	NL	S4S5	G5	3	
	X	MO	A Moss	<i>Fontinalis sullivantii</i>	Fontinalaceae			NL	NL	S1	G3G5		
X		MO	A Moss	<i>Sphagnum sp.</i>	Sphagnaceae			NL	NL				
	X	MO	A Moss	<i>Abietinella abietina</i>	Thuidiaceae			NL	NL	S4S5	G4G5		
	X	RU	Richardson Rush	<i>Juncus alpinoarticulatus</i>	Juncaceae	5	-5	NL	NL	S5	G5	2	

APPENDIX 7. BOTANICAL INVENTORY (About Associate Inc. and CVC data)
 Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

Project: AA12-137A

AA ¹	CVC ²	PLANT TYPE ³	COMMON NAME	SCIENTIFIC NAME	FAMILY	CC ⁴	CW ⁵	SARO ⁶	SARA ⁷	S-Rank ⁸	G-Rank ⁹	CVC 2010 ¹⁰	Wellington County ¹¹
	X	RU	Jointed Rush	<i>Juncus articulatus</i>	Juncaceae	5	-5	NL	NL	S5	G5	4	
	X	RU	Narrow-panicked Rush	<i>Juncus brevicaudatus</i>	Juncaceae	6	-5	NL	NL	S5	G5	2	
	X	RU	Toad Rush	<i>Juncus bufonius</i>	Juncaceae	1	-4	NL	NL	S5	G5	3	
X	X	RU	Dudley's Rush	<i>Juncus dudleyi</i>	Juncaceae	1	0	NL	NL	S5	G5	4	
X	X	RU	Soft Rush	<i>Juncus effusus</i>	Juncaceae	3	-5	NL	NL	S5	G5T?	4	
	X	RU	Knotted Rush	<i>Juncus nodosus</i>	Juncaceae	5	-5	NL	NL	S5	G5	4	
X	X	RU	Path Rush	<i>Juncus tenuis</i>	Juncaceae	0	0	NL	NL	S5	G5	4	
	X	SE	Black Sedge	<i>Carex arcata</i>	Cyperaceae	5	5	NL	NL	S5	G5?	3	
X	X	SE	Golden-fruited Sedge	<i>Carex aurea</i>	Cyperaceae	4	-4	NL	NL	S5	G5	3	
X	X	SE	Bebb's Sedge	<i>Carex bebbii</i>	Cyperaceae	3	-5	NL	NL	S5	G5	4	
	X	SE	Woodland Sedge	<i>Carex blanda</i>	Cyperaceae	3	0	NL	NL	S5	G5?	4	
	X	SE	Crested Sedge	<i>Carex cristatella</i>	Cyperaceae	3	-4	NL	NL	S5	G5	4	
	X	SE	Softleaf Sedge	<i>Carex disperma</i>	Cyperaceae	8	-5	NL	NL	S5	G5	3	
X	X	SE	Yellow Sedge	<i>Carex flava</i>	Cyperaceae	5	-5	NL	NL	S5	G5	3	
X	X	SE	Graceful Sedge	<i>Carex gracillima</i>	Cyperaceae	4	3	NL	NL	S5	G5	4	
X	X	SE	Porcupine Sedge	<i>Carex hystericina</i>	Cyperaceae	5	-5	NL	NL	S5	G5	4	
X	X	SE	Inland Sedge	<i>Carex interior</i>	Cyperaceae	6	-5	NL	NL	S5	G5	3	
X	X	SE	Bladder Sedge	<i>Carex intumescens</i>	Cyperaceae	6	-4	NL	NL	S5	G5	3	
X	X	SE	Lake-bank Sedge	<i>Carex lacustris</i>	Cyperaceae	5	-5	NL	NL	S5	G5	4	
	X	SE	Smooth-sheath Sedge	<i>Carex laevivaginata</i>	Cyperaceae	8	-5	NL	NL	S4	G5	2	
	X	SE	Bristly-stalk Sedge	<i>Carex leptalea</i>	Cyperaceae	8	-5	NL	NL	S5	G5	2	
	X	SE	Finely-nerved Sedge	<i>Carex leptoneuria</i>	Cyperaceae	5	0	NL	NL	S4	G4	2	✓
X	X	SE	White-tinged Sedge	<i>Carex peckii</i>	Cyperaceae	6	5	NL	NL	S5	G4G5	3	
	X	SE	Longstalk Sedge	<i>Carex pedunculata</i>	Cyperaceae	5	5	NL	NL	S5	G5	4	
X	X	SE	Pennsylvania Sedge	<i>Carex pennsylvanica</i>	Cyperaceae	5	5	NL	NL	S5	G5	4	
	X	SE	Prairie Sedge	<i>Carex prairea</i>	Cyperaceae	7	-4	NL	NL	S5	G5?	2	
	X	SE	Cyperus-like Sedge	<i>Carex pseudocyperus</i>	Cyperaceae	6	-5	NL	NL	S5	G5	4	
X	X	SE	Retrorse Sedge	<i>Carex retrosa</i>	Cyperaceae	5	-5	NL	NL	S5	G5	4	
	X	SE	Rough Sedge	<i>Carex scabrata</i>	Cyperaceae	8	-5	NL	NL	S5	G5	2	
X		SE	Sedge sp.	<i>Carex sp.</i>	Cyperaceae								
X	X	SE	Stalk-grain Sedge	<i>Carex stipata</i>	Cyperaceae	3	-5	NL	NL	S5	G5	4	
X	X	SE	Tussock Sedge	<i>Carex stricta</i>	Cyperaceae	4	-5	NL	NL	S5	G5	2	
	X	SE	Three-seed Sedge	<i>Carex trisperma</i>	Cyperaceae	9	-5	NL	NL	S5	G5	2	
X	X	SE	Bladder Sedge	<i>Carex utriculata</i>	Cyperaceae	7	-5	NL	NL	S5	G5	3	
X	X	SE	Fox Sedge	<i>Carex vulpinoidea</i>	Cyperaceae	3	-5	NL	NL	S5	G5	4	
	X	SE	Bald Spikerush	<i>Eleocharis erythropoda</i>	Cyperaceae	4	-5	NL	NL	S5	G5	4	
X	X	SE	Green Keeled Cottongrass	<i>Eriophorum viridicarinatum</i>	Cyperaceae	9	-5	NL	NL	S5	G5	2	
X	X	SE	Soft-stem Club-rush	<i>Schoenoplectus tabernaemontani</i>	Cyperaceae	5	-5	NL	NL	S5	G?	3	
X	X	SE	Dark-green Bulrush	<i>Scirpus atrovirens</i>	Cyperaceae	3	-5	NL	NL	S5	G5?	4	
X	X	SE	Common Woolly Bulrush	<i>Scirpus cyperinus</i>	Cyperaceae	4	-5	NL	NL	S5	G5	3	
X	X	SE	Red-tinge Bulrush	<i>Scirpus microcarpus</i>	Cyperaceae	4	-5	NL	NL	S5	G5	3	
	X	SH	Mountain Maple	<i>Acer spicatum</i>	Aceraceae	6	3	NL	NL	S5	G5	3	
X		SH	Staghorn Sumac	<i>Rhus typhina</i>	Anacardiaceae	1	5	NL	NL	S5	G5	4	
X	X	SH	Rydberg's Poison Ivy	<i>Toxicodendron rydbergii</i>	Anacardiaceae	0	0	NL	NL	S5	G5	4	
	X	SH	Black Holly	<i>Ilex verticillata</i>	Aquifoliaceae	5	-4	NL	NL	S5	G5	2	
X	X	SH	European Alder	<i>Alnus glutinosa</i>	Betulaceae	0	-2	NL	NL	SE4	G?	5	
	X	SH	Twinflower	<i>Linnaea borealis</i>	Caprifoliaceae	6	0	NL	NL	S5	G5T?	2	

APPENDIX 7. BOTANICAL INVENTORY (About Associate Inc. and CVC data)
Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

Project: AA12-137A

AA ¹	CVC ²	PLANT TYPE ³	COMMON NAME	SCIENTIFIC NAME	FAMILY	CC ⁴	CW ⁵	SARO ⁶	SARA ⁷	S-Rank ⁸	G-Rank ⁹	CVC 2010 ¹⁰	Wellington County ¹¹
X	X	SH	American Fly-honeysuckle	<i>Lonicera canadensis</i>	Caprifoliaceae	6	3	NL	NL	S5	G5	3	
X		SH	Morrow Honeysuckle	<i>Lonicera morrowii</i>	Caprifoliaceae	0	5	NL	NL	SE3	G?	5	
	X	SH	Swamp Fly-honeysuckle	<i>Lonicera oblongifolia</i>	Caprifoliaceae	8	-5	NL	NL	S4S5	G4	2	
X	X	SH	Tartarian Honeysuckle	<i>Lonicera tatarica</i>	Caprifoliaceae	0	3	NL	NL	SE5	G?	5	
X	X	SH	Common Elderberry	<i>Sambucus canadensis</i>	Caprifoliaceae	5	-2	NL	NL	S5	G5	4	
X	X	SH	Red Elderberry	<i>Sambucus racemosa</i>	Caprifoliaceae	3	3	NL	NL	S5		4	
X	X	SH	Nannyberry	<i>Viburnum lentago</i>	Caprifoliaceae	4	-1	NL	NL	S5	G5	4	
X	X	SH	European Highbush-cranberry	<i>Viburnum opulus</i>	Caprifoliaceae	0	-3	NL	NL	SE4	G5	5	
X	X	SH	Alternate-leaf Dogwood	<i>Cornus alternifolia</i>	Cornaceae	6	5	NL	NL	S5	G5	4	
X	X	SH	Bunchberry	<i>Cornus canadensis</i>	Cornaceae	7	0	NL	NL	S5	G5	2	
X	X	SH	Red-osier Dogwood	<i>Cornus stolonifera</i>	Cornaceae	2	-3	NL	NL	S5	G5	4	
	X	SH	Autum Olive	<i>Elaeagnus umbellata</i>	Elaeagnaceae	0	3	NL	NL	SE3	G?	5	
	X	SH	Creeping Snowberry	<i>Gaultheria hispida</i>	Ericaceae	8	-3	NL	NL	S5	G5	2	
X	X	SH	Wild Black Currant	<i>Ribes americanum</i>	Grossulariaceae	4	-3	NL	NL	S5	G5	4	
X	X	SH	Prickly Gooseberry	<i>Ribes cynosbati</i>	Grossulariaceae	4	5	NL	NL	S5	G5	4	
X	X	SH	Smooth Gooseberry	<i>Ribes hirtellum</i>	Grossulariaceae	6	-3	NL	NL	S5	G5	2	✓
	X	SH	Northern Red Currant	<i>Ribes rubrum</i>	Grossulariaceae	0	5	NL	NL	SE5	G4G5	5	
X		SH	Current sp.	<i>Ribes sp.</i>	Grossulariaceae								
X	X	SH	Swamp Red Currant	<i>Ribes triste</i>	Grossulariaceae	6	-5	NL	NL	S5	G5	3	
X	X	SH	Common Lilac	<i>Syringa vulgaris</i>	Oleaceae	0	5	NL	NL	SE5	G?	5	
X	X	SH	Alderleaf Buckthorn	<i>Rhamnus alnifolia</i>	Rhamnaceae	7	-5	NL	NL	S5	G5	2	
X	X	SH	Buckthorn	<i>Rhamnus cathartica</i>	Rhamnaceae	0	3	NL	NL	SE5	G?	5	
X	X	SH	Downy Serviceberry	<i>Amelanchier arborea</i>	Rosaceae	5	3	NL	NL	S5	G5	3	
X		SH	Hawthorn sp.	<i>Crataegus sp.</i>	Rosaceae								
X	X	SH	Common Apple	<i>Malus pumila</i>	Rosaceae	0	5	NL	NL	SE5	G5	5	
X	X	SH	Choke Cherry	<i>Prunus virginiana</i>	Rosaceae	2	3	NL	NL	S5	G5T?	4	
X	X	SH	Multiflora Rose	<i>Rosa multiflora</i>	Rosaceae	0	3	NL	NL	SE4	G?	5	
X		SH	Common Blackberry	<i>Rubus allegheniensis</i>	Rosaceae	2	2	NL	NL	S5	G5	4	
X	X	SH	Common Red Raspberry	<i>Rubus idaeus ssp. idaeus</i>	Rosaceae			NL	NL	SE1	G5T5	5	
X	X	SH	Wild Red Raspberry	<i>Rubus idaeus ssp. strigosus</i>	Rosaceae	0	-2	NL	NL	S5	G5	4	
X	X	SH	Dwarf Raspberry	<i>Rubus pubescens</i>	Rosaceae	4	-4	NL	NL	S5	G5	3	
X	X	SH	European Mountain-ash	<i>Sorbus aucuparia</i>	Rosaceae	0	5	NL	NL	SE4	G5	5	
X	X	SH	Narrow-leaved Meadow-sweet	<i>Spiraea alba</i>	Rosaceae	3	-4	NL	NL	S5	G5	3	
X	X	SH	Bebb's Willow	<i>Salix bebbiana</i>	Salicaceae	4	-4	NL	NL	S5	G5	3	
X	X	SH	Pussy Willow	<i>Salix discolor</i>	Salicaceae	3	-3	NL	NL	S5	G5	4	
X	X	SH	Heart-leaved Willow	<i>Salix eriocephala</i>	Salicaceae	4	-3	NL	NL	S5	G5	4	
X	X	SH	Shining Willow	<i>Salix lucida</i>	Salicaceae	5	-4	NL	NL	S5	G5	3	
X	X	SH	Meadow Willow	<i>Salix petiolaris</i>	Salicaceae	3	-4	NL	NL	S5	G5	3	

APPENDIX 7. BOTANICAL INVENTORY (About Associate Inc. and CVC data)
 Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

Project: AA12-137A

AA ¹	CVC ²	PLANT TYPE ³	COMMON NAME	SCIENTIFIC NAME	FAMILY	CC ⁴	CW ⁵	SARO ⁶	SARA ⁷	S-Rank ⁸	G-Rank ⁹	CVC 2010 ¹⁰	Wellington County ¹¹
X		SH	Purple Willow	<i>Salix purpurea</i>	Salicaceae	0	-3	NL	NL	SE4	G5	5	
X	X	SH	Canadian Yew	<i>Taxus canadensis</i>	Taxaceae	7	3	NL	NL	S4	G5	3	
X	X	SH	February Daphne	<i>Daphne mezereum</i>	Thymelaeaceae	0	0	NL	NL	SE2	G?	5	
X	X	TR	Manitoba Maple	<i>Acer negundo</i>	Aceraceae	0	-2	NL	NL	S5	G5	5	
X	X	TR	Norway Maple	<i>Acer platanoides</i>	Aceraceae	0	5	NL	NL	SE5	G?	5	
X	X	TR	Red Maple	<i>Acer rubrum</i>	Aceraceae	4	0	NL	NL	S5	G5	4	
X		TR	Silver Maple	<i>Acer saccharinum</i>	Aceraceae	5	3	NL	NL	S5	G5	3	
X	X	TR	Sugar Maple	<i>Acer saccharum ssp. saccharum</i>	Aceraceae	4	3	NL	NL	S5	G5	4	
X		TR	Freeman's Maple	<i>Acer x freemanii</i>	Aceraceae			NL	NL	S4	G?	4	
X	X	TR	Yellow Birch	<i>Betula alleghaniensis</i>	Betulaceae	6	0	NL	NL	S5	G5	3	
X	X	TR	White Birch	<i>Betula papyrifera</i>	Betulaceae	2	2	NL	NL	S5	G5	4	
X	X	TR	Eastern Hop-hornbeam	<i>Ostrya virginiana</i>	Betulaceae	4	4	NL	NL	S5	G5	4	
X	X	TR	Eastern White Cedar	<i>Thuja occidentalis</i>	Cupressaceae	4	-3	NL	NL	S5	G5	4	
X	X	TR	American Beech	<i>Fagus grandifolia</i>	Fagaceae	6	3	NL	NL	S4	G5	3	
X		TR	Northern Red Oak	<i>Quercus rubra</i>	Fagaceae	6	3	NL	NL	S5	G5	4	
X		TR	Black Walnut	<i>Juglans nigra</i>	Juglandaceae	5	3	NL	NL	S4	G5	4	
X	X	TR	White Ash	<i>Fraxinus americana</i>	Oleaceae	4	3	NL	NL	S5	G5	4	
X		TR	Black Ash	<i>Fraxinus nigra</i>	Oleaceae	7	-4	NL	NL	S5	G5	3	
X	X	TR	Green Ash	<i>Fraxinus pennsylvanica</i>	Oleaceae	3	-3	NL	NL	S5	G5	4	
X	X	TR	Balsam Fir	<i>Abies balsamea</i>	Pinaceae	5	-3	NL	NL	S5	G5	3	
X		TR	European Larch	<i>Larix decidua</i>	Pinaceae	0	5	NL	NL	SE2	G?	5	
X	X	TR	Tamarack	<i>Larix laricina</i>	Pinaceae	7	-3	NL	NL	S5	G5	3	
X		TR	Norway Spruce	<i>Picea abies</i>	Pinaceae	0	5	NL	NL	SE3	G?	5	
X	X	TR	White Spruce	<i>Picea glauca</i>	Pinaceae	6	3	NL	NL	S5	G5	2	
X	X	TR	Black Spruce	<i>Picea mariana</i>	Pinaceae	8	-3	NL	NL	S5	G5	2	
X	X	TR	Blue Spruce	<i>Picea pungens</i>	Pinaceae	0	3	NL	NL	SE1	G5	5	
X	X	TR	Eastern White Pine	<i>Pinus strobus</i>	Pinaceae	4	3	NL	NL	S5	G5	4	
X		TR	Scots Pine	<i>Pinus sylvestris</i>	Pinaceae	0	5	NL	NL	SE5	G?	5	
X	X	TR	Eastern Hemlock	<i>Tsuga canadensis</i>	Pinaceae	7	3	NL	NL	S5	G5	4	
X	X	TR	Wild Black Cherry	<i>Prunus serotina</i>	Rosaceae	3	3	NL	NL	S5	G5	4	
X	X	TR	Balsam Poplar	<i>Populus balsamifera</i>	Salicaceae	2	-3	NL	NL	S5	G5T?	4	
X	X	TR	Large-tooth Aspen	<i>Populus grandidentata</i>	Salicaceae	5	3	NL	NL	S5	G5	3	
X	X	TR	Trembling Aspen	<i>Populus tremuloides</i>	Salicaceae	2	0	NL	NL	S5	G5	4	
X	X	TR	White Willow	<i>Salix alba</i>	Salicaceae	0	-3	NL	NL	SE4	G5	5	
X		TR	Crack Willow	<i>Salix fragilis</i>	Salicaceae	0	0	NL	NL	SE	G5	5	
X	X	TR	American Basswood	<i>Tilia americana</i>	Tiliaceae	4	3	NL	NL	S5	G5	4	
X	X	TR	American Elm	<i>Ulmus americana</i>	Ulmaceae	3	-2	NL	NL	S5	G5?	4	
X	X	VI	Wild Mock-cucumber	<i>Echinocystis lobata</i>	Cucurbitaceae	3	-2	NL	NL	S5	G5	4	
X		VI	Broad-leaf Peavine	<i>Lathyrus latifolius</i>	Fabaceae	0	5	NL	NL	SE4	G?	5	
X	X	VI	Tufted Vetch	<i>Vicia cracca</i>	Fabaceae	0	5	NL	NL	SE5	G?	5	
	X	VI	Spring Vetch	<i>Vicia sativa</i>	Fabaceae	0	3	NL	NL	SE5	G?T?	5	
X	X	VI	Black Bindweed	<i>Fallopia convolvulus</i>	Polygonaceae	0	1	NL	NL	SE5	G?	5	
X	X	VI	Virginia Clematis	<i>Clematis virginiana</i>	Ranunculaceae	3	0	NL	NL	S5	G5	4	
	X	VW	Mountain Honeysuckle	<i>Lonicera dioica</i>	Caprifoliaceae	5	3	NL	NL	S5	G5	3	
X	X	VW	Climbing Nightshade	<i>Solanum dulcamara</i>	Solanaceae	0	0	NL	NL	SE5	G?	5	
X	X	VW	Inserted Virginia Creeper	<i>Parthenocissus inserta</i>	Vitaceae	3	3	NL	NL	S5	G5	4	
X	X	VW	Riverbank Grape	<i>Vitis riparia</i>	Vitaceae	0	-2	NL	NL	S5	G5	4	

1	AA: Botanical data collected by Aboud & Associates Inc. during 2014
2	CVC: Botanical data collected by Credit River Conservation from 2008 to 2009
3	Plant Types: AL = Algae; FE = Fern; FO = Forb; GR = Grass; LC = Lichen; LV = Liverwort; MO = Moss; RU = Rush; SE = Sedge; SH = Shrub; TR = Tree; VI = Herbaceous vine; VW =
4	CC: Coefficient of Conservatism reflects a species' fidelity to a specific habitat. Range from 0 to 10; 10 = very conservative, not likely in disturbed habitats, 1 = least conservative, likely
5	CW: Coefficient of Wetness reflects a species' affinity for wet soil conditions. Range from -5 to 5; -5 = obligate wetland species, 5 = obligate upland species.
6	SARO: Status under the Provincial Endangered Species Act, listed on the Species at Risk in Ontario (SARO) list. In order of severity, statuses include: EXP = Extirpated; END =
7	SARA: Status under the National Species at Risk Act (SARA), assessed by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC). In order of severity, statuses
8	S-Rank: Provincial rarity rank. Range from S1 to S5; S1 = Extremely rare, S5 = Very common. NR = Unranked; U = Unrankable.
9	questionable; T = Applies to subspecies or variety; Nothing = Rank not yet obtained.
10	CVC 2010: Species of Conservation Concern Teir Ranking. Range from 1 to 5; 1 = Species of Conservation Concern, 2 = Species of Interest, 3 = Urban Interest, 4 = Secure Species, 5 = Non-native & Non-native Hybrid Species.
11	Wellington County: Significant Flora Species within Wellington County as identified by Dougan & Associates, with Snell & Cecile. 2009. Guelph Natural Heritage Strategy. Phase 2, Volume 2 (Significant Plant List for Wellington County). Guelph, Ontario.

DRAFT

STATION	DATE	SPECIES				
		Gray Treefrog	Spring Peeper	Green Frog	Northern Leopard Frog	Wood Frog
A	15-Apr-15					
	28-May-15		1-3	1-2		
	24-Jun-15	1-1		1-4		
B1	15-Apr-15		2-17			2-10
	28-May-15		1-2			
	24-Jun-15	1-2				
B2	15-Apr-15		1-2			
	28-May-15		1-2			
	24-Jun-15					
C1	15-Apr-15		1-4*			
	28-May-15	2-7	1-3			
	24-Jun-15	2-5*		1-2*		
C2	15-Apr-15		2-8			
	28-May-15	1-2	1-4		1-2	1-1
	24-Jun-15	1-1		1-2		
D	15-Apr-15		3			1-3
	28-May-15	1-4	1-1	1-1		
	24-Jun-15			1-1		
E	15-Apr-15					
	28-May-15		1-1			
	24-Jun-15					
F	15-Apr-15					
	28-May-15	2-11	1-2			
	24-Jun-15	1-6		1-1		
G	15-Apr-15					
	28-May-15	2-9	1-1	1-3		
	24-Jun-15	1-3		1-3		
H	15-Apr-15					
	28-May-15	1-1				
	24-Jun-15					

* indicates call heard outside of survey area.

Amphibian Call Level codes:

- 1 - # :Calls not simultaneous, number of individuals can be accurately counted
- 2 - # :Some calls simultaneous, number of individuals can be reliably estimated
- 3 :Full chorus, calls continuous and overlapping, number of individuals cannot reliably be estimated

COMMON NAME	SCIENTIFIC NAME	COSARO	COSEWIC	SARA	S-RANK	G-RANK	Wellington	CVC	PIF	PC1 Deciduous forest		PC2 OA, Coniferous Swamp		PC3 Deciduous forest		PC4 Meadow Marsh, Mixed Swamp		PC5 Shallow Marsh, Mixed Swamp		PC6 Mixed Swamp, Coniferous Forest		PC7 Woodland, Deciduous Forest		PC8 Hillsburgh Pond and Dam		PC9 Cattail Marsh		PC10 Creek and Riparian		Site Totals #	Overall Highest Breeding Evidence
										#	HBE	#	HBE	#	HBE	#	HBE	#	HBE	#	HBE	#	HBE	#	HBE	#	HBE	#	HBE		
Great Blue Heron	<i>Ardea herodias</i>				S4	G5	✓	3	0		1 X	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	Observed	
Green Heron	<i>Butorides virescens</i>				S4B	G5	✓	2	0		1 H	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	Possible		
Canada Goose	<i>Branta canadensis</i>				S5	G5		4	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Observed	
Mallard	<i>Anas platyrhynchos</i>				S5	G5	✓	4	0		0	0	0	0	13 P	0	0	0	0	0	0	0	0	0	0	0	0	0	13	Probable	
Killdeer	<i>Charadrius vociferus</i>				S5B,S5N	G5		3	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Observed	
Ring-billed Gull	<i>Larus delawarensis</i>				S5B,S4N	G5	✓	2	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Observed	
Mourning Dove	<i>Zenaida macroura</i>				S5	G5	✓	4	0		0	0	0	1 S	0	0	0	0	0	0	0	0	0	0	0	0	1 S	2	Possible		
Belted Kingfisher	<i>Megaceryle alcyon</i>				S4B	G5	✓	3	✓	0	2 P	0	0	1 H	0	0	0	0	0	0	0	0	0	0	1 H	0	0	5	Probable		
Downy Woodpecker	<i>Picoides pubescens</i>				S5	G5	✓	4	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1 T	0	1	Probable		
Northern Flicker	<i>Colaptes auratus</i>				S4B	G5	✓	3	✓	0	1 S	0	1 T	1 S	1 S	1 S	2 T	1 T	0 S	0	0	0	0	0	0	0	0	7	Probable		
Eastern Wood-pewee	<i>Contopus virens</i>	SC	SC		S4B	G5	✓	1	✓	1 S	1 T	1 S	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	Probable		
Alder Flycatcher	<i>Empidonax alnorum</i>				S5B	G5	✓		0		0	0	0	0	1 S	0	0	0	0	0	0	0	0	0	0	0	0	1	Possible		
Eastern Phoebe	<i>Sayornis phoebe</i>				S5B	G5		3	0		0	0	0	0	0	0	1 S	0	0	0	0	0	0	0	0	0	0	1	Possible		
Great Crested Flycatcher	<i>Myiarchus crinitus</i>				S4B	G5		3	0		0	0	0	1 S	1 T	0	0	0	0	0	0	0	0	0	0	0	0	3	Probable		
Eastern Kingbird	<i>Tyrannus tyrannus</i>				S4B	G5		3	0		0	0	0	2 S	1 A	0	0	0	0	0	0	0	0	0	0	0	0	3	Probable		
Tree Swallow	<i>Tachycineta bicolor</i>				S4B	G5		3	0		2 H	0	0	1 H	0	0	0	0	0	0	0	0	0	1 H	1 H	0	0	5	Possible		
Blue Jay	<i>Cyanocitta cristata</i>				S5	G5		4	1 FY	2 A	0	0	0	0	0	0	0	0	1 H	1 S	0	0	0	0	0	1 H	6	confirmed			
American Crow	<i>Corvus brachyrhynchos</i>				S5B	G5	✓	2	1 A	1 A	0	0	0	2 A	1	0	0	0	0	0	0	0	0	0	3 A	2 H	12	Probable			
Black-capped Chickadee	<i>Poecile atricapillus</i>				S5	G5		4	1 S	1 S	2 A	0	0	2 S	2 T	2 S	3 FY	0	0	0	0	0	0	0	1 S	14	confirmed				
Red-breasted Nuthatch	<i>Sitta canadensis</i>				S5	G5		3	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	Possible		
House Wren	<i>Troglodytes aedon</i>				S5B	G5	✓	4	0		0	0	0	1 A	1 S	0	0	0	0	0	0	0	0	0	0	1 S	3	Probable			
Winter Wren	<i>Troglodytes troglodytes</i>				S5B	G5	✓	3	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	Possible		
American Robin	<i>Turdus migratorius</i>				S5B	G5		4	1 T	1 T	1 A	5 A	4 T	1 A	2 A	1 A	0	0	0	0	0	0	0	0	0	5 A	21	Probable			
Gray Catbird	<i>Dumetella carolinensis</i>				S4B	G5	✓	3	0		0	1 S	0	1 S	0	0	0	0	0	0	0	0	0	0	1 S	0	4	Possible			
Cedar Waxwing	<i>Bombycilla cedrorum</i>				S5B	G5	✓	3	0		1 H	0	3 H	2 H	0	0	0	0	0	0	0	0	0	0	0	0	0	12	Possible		
European Starling	<i>Sturnus vulgaris</i>				SNA	G5			0		0	0	0	0	0	0	0	0	0	0	0	0	0	3 S	1 H	1 H	5	Possible			
Warbling Vireo	<i>Vireo gilvus</i>				S5B	G5		5	0		0	0	0	1 T	1 S	0	0	0	0	0	0	0	0	0	1 T	1 S	0	4	Probable		
Red-eyed Vireo	<i>Vireo olivaceus</i>				S5B	G5	✓	4	1 T	1 T	1 S	1 S	2 S	0	0	0	0	0	0	0	0	0	0	0	0	0	0	7	Probable		
Yellow Warbler	<i>Dendroica petechia</i>				S5B	G5		4	0		0	0	0	1 T	2 S	0	0	0	0	0	0	0	0	0	0	1 T	0	5	Probable		
Chestnut-sided Warbler	<i>Dendroica pensylvanica</i>				S5B	G5	✓	2	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	Possible		
Northern Waterthrush	<i>Seiurus noveboracensis</i>				S5B	G5	✓	3	0		1 S	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	Possible		
Common Yellowthroat	<i>Geothlypis trichas</i>				S5B	G5	✓	4	0		1 S	0	0	2 S	2 T	0	0	0	0	0	0	0	0	0	0	0	0	7	Probable		
Northern Cardinal	<i>Cardinalis cardinalis</i>				S5	G5	✓	4	0		0	0	0	0	1 S	1 S	0	0	0	0	0	0	0	0	0	0	0	3	Possible		
Rose-breasted Grosbeak	<i>Pheucticus ludovicianus</i>				S4B	G5		3	0		0	0	0	1 S	1 H	1 S	0	0	0	0	0	0	0	0	0	0	0	3	Possible		
Indigo Bunting	<i>Passerina cyanea</i>				S4B	G5			0		1 S	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	Possible		
Chipping Sparrow	<i>Spizella passerina</i>				S5B	G5	✓	4	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	Possible		
Song Sparrow	<i>Melospiza melodia</i>				S5B	G5	✓	4	0		0	0	0	2 S	2 S	0	0	0	0	0	0	0	0	0	0	2 T	1 S	11	confirmed		
Swamp Sparrow	<i>Melospiza georgiana</i>				S5B	G5		4	0		1 S	0	0	1 S	2 S	0	0	0	0	0	0	0	0	0	0	0	0	6	Probable		
White-throated Sparrow	<i>Zonotrichia albicollis</i>				S5B	G5		3	0		0	0	0	0	1 S	0 S	0	0	0	0	0	0	0	0	0	0	0	1	Possible		
Red-winged Blackbird	<i>Agelaius phoeniceus</i>				S4	G5	✓	4	0		2 A	0	4 T	4 S	0	0	0	0	0	0	0	0	0	6 CF	12 FY	0	0	28	confirmed		
Eastern Meadowlark	<i>Sturnella magna</i>	THR	THR		S4B	G5	✓	1	✓	0	0	0	0	0 S	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Possible		
Common Grackle	<i>Quiscalus quiscula</i>				S5B	G5		4	0		0	0	0	1 A	6 FY	0	0	0	0	0	0	0	0	0	2 H	1 H	0	10	confirmed		
Brown-headed Cowbird	<i>Molothrus ater</i>				S4B	G5	✓	4	0		0	0	0	0	0	0	0	0	0	0	0	0	0	1 S	1 H	0	0	2	Possible		
Baltimore Oriole	<i>Icterus galbula</i>				S4B	G5		3	✓	0	1 S	1 S	2 S	0	0	0	0	0	0	0	0	0	0	0	1 A	1 H	0	6	Probable		
American Goldfinch	<i>Carduelis tristis</i>				S5B	G5		4	0		1 H	0	0	0	1 H	0	0	0	0	0	0	0	0	1 H	7 H	3 H	2 H	15	Possible		
House Sparrow	<i>Passer domesticus</i>				SNA	G5		5	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	Possible		

Legend:
 SARO: Species at Risk Ontario
 COSEWIC: Committee on the Status of Endangered wildlife in Canada
 SARA: Species at Risk Act
 ESA: Endangered Species Act
 END: Endangered
 THR: Threatened
 SC: special Concern
 NAR: Not At Risk
 NL: Not listed
 DD: Data Deficient
 HBE: Highest Breeding Evidence over 2 surveys
 PIF: Priority species in BCR13

G-Rank:
 G1: Extremely rare globally
 G1G2: Extremely rare to very rare globally
 G2: Very rare globally
 G2G3: Very rare to uncommon globally
 G3: Rare to uncommon globally
 G3G4: Rare to common globally
 G4: Common globally
 G4G5: Common to very common globally
 G5: Very common globally; demonstrably secure
 T: rank applies to a subspecies or variety

S-Rank:
 S1: Critically Imperiled—Critically imperiled in the province
 S2: Imperiled—Imperiled in the province
 S3: Vulnerable—Vulnerable in the province
 S4: Apparently Secure—Uncommon but not rare
 S5: Secure—Common, widespread, and abundant
 SX: Presumed extirpated
 SH: Possibly Extirpated (Historical)
 SNR: Unranked
 SU: Unrankable—Currently unrankable
 SNA: Not applicable—A conservation status rank is not applicable
 S#S#: Range Rank— indicates range of uncertainty about the status
 S#B- Breeding status rank
 S#N- Non Breeding status rank
 ?: Indicates uncertainty in the assigned rank

CVC Tiers:
 1 - Species of Conservation Concern
 2 - Species of Interest
 3 - Species of Urban Interest
 4 - Secure Species
 5 - Non-native & Non-native Hybrid Species

Breeding Evidence Codes
Observed
 X-no breeding evidence
 FO-flyover
Possible
 H-Suitable habitat
 S-Singing male
probable
 P-PairD-Display
 T-Territory (2 visits)
 D-Display
 V-Visiting nest
 A-Agitated
 B-Broodpatch
 N-Nest building or excavation

Confirmed
 DD-Distracted display
 NU-Used nest
 FY-Fledged young
 AE-Adult entering/leaving nest
 FS-Adult carrying fecal sac
 CF-Adult carrying food
 NE-Nest with eggs
 NY-Nest with young

Wellington County:
 ✓ Significant Species

APPENDIX 10. MARSH BREEDING BIRD RESULTS

Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

COMMON NAME	SCIENTIFIC NAME	STATION	ROUND	BE	in/out	tally
American Redstart	<i>Setophaga ruticilla</i>	MBB1	2	s	in	1
American Robin	<i>Turdus migratorius</i>	MBB1	1	H	in	2
Blue Jay	<i>Cyanocitta cristata</i>	MBB1	2	a	in	1
Common Grackle	<i>Quiscalus quiscula</i>	MBB1	2	h	in	1
Downy Woodpecker	<i>Picoides pubescens</i>	MBB1	2	t	in	1
Gray Catbird	<i>Dumetella carolinensis</i>	MBB1	1	S	in	1
Green Heron	<i>Butorides virescens</i>	MBB1	1	H	out	1
Mallard	<i>Anas platyrhynchos</i>	MBB1	2	h	in	3
Red-eyed Vireo	<i>Vireo olivaceus</i>	MBB1	1	S	in	1
Red-winged Blackbird	<i>Agelaius phoeniceus</i>	MBB1	1	P	in	8
Song Sparrow	<i>Melospiza melodia</i>	MBB1	1	S	in	1
Tree Swallow	<i>Tachycineta bicolor</i>	MBB1	1	fo	n/a	6
Yellow Warbler	<i>Dendroica petechia</i>	MBB1	1	S	in	1
American Black Duck	<i>Anas rubripes</i>	MBB2	2	h	in	1
Baltimore Oriole	<i>Icterus galbula</i>	MBB2	2	s	in	1
Black-and-white Warbler	<i>Mniotilta varia</i>	MBB2	1	s	in	2
Cliff Swallow	<i>Petrochelidon pyrrhonota</i>	MBB2	1	fo	n/a	2
Common Grackle	<i>Quiscalus quiscula</i>	MBB2	2	a	in	3
Great Crested Flycatcher	<i>Myiarchus crinitus</i>	MBB2	1	S	in	2
Northern Flicker	<i>Colaptes auratus</i>	MBB2	2	t	in	2
Red-breasted Nuthatch	<i>Sitta canadensis</i>	MBB2	2	s	in	1
White-throated Sparrow	<i>Zonotrichia albicollis</i>	MBB2	1	s	in	2
American Goldfinch	<i>Carduelis tristis</i>	MBB3	1	H	in	1
American Robin	<i>Turdus migratorius</i>	MBB3	1	A	in	2
American Robin	<i>Turdus migratorius</i>	MBB3	2	fy	in	1
Baltimore Oriole	<i>Icterus galbula</i>	MBB3	1	H	in	1
Belted Kingfisher	<i>Megasceryle alcyon</i>	MBB3	2	h	in	1
Brown-headed Cowbird	<i>Molothrus ater</i>	MBB3	2	s	in	1
Cedar Waxwing	<i>Bombcilla cedrorum</i>	MBB3	2	h	in	4
House Wren	<i>Troglodytes aedon</i>	MBB3	2	s	in	1
Northern Flicker	<i>Colaptes auratus</i>	MBB3	1	T	in	1
Red-winged Blackbird	<i>Agelaius phoeniceus</i>	MBB3	1	A	in	4
Song Sparrow	<i>Melospiza melodia</i>	MBB3	2	a	in	1
Warbling Vireo	<i>Vireo gilvus</i>	MBB3	1	S	in	1
Alder Flycatcher	<i>Empidonax alnorum</i>	MBB4	2	h	in	1
American Crow	<i>Corvus brachyrhynchos</i>	MBB4	1	h	in	2
American Crow	<i>Corvus brachyrhynchos</i>	MBB4	2	h	in	1
American Goldfinch	<i>Carduelis tristis</i>	MBB4	1	h	in	1
American Goldfinch	<i>Carduelis tristis</i>	MBB4	2	h	in	2
American Robin	<i>Turdus migratorius</i>	MBB4	2	a	in	2
Baltimore Oriole	<i>Icterus galbula</i>	MBB4	1	s	in	1
Belted Kingfisher	<i>Megasceryle alcyon</i>	MBB4	1	h	in	1
Belted Kingfisher	<i>Megasceryle alcyon</i>	MBB4	2	h	out	1
Black-capped Chickadee	<i>Poecile atricapillus</i>	MBB4	1	s	in	1
Black-capped Chickadee	<i>Poecile atricapillus</i>	MBB4	2	s	in	3
Blue Jay	<i>Cyanocitta cristata</i>	MBB4	2	fy	in	1
Chipping Sparrow	<i>Spizella passerina</i>	MBB4	1	s	in	1
Common Yellowthroat	<i>Geothlypis trichas</i>	MBB4	2	s	in	1
Downy Woodpecker	<i>Picoides pubescens</i>	MBB4	2	t	in	1
Eastern Kingbird	<i>Tyrannus tyrannus</i>	MBB4	1	a	in	1
Great Crested Flycatcher	<i>Myiarchus crinitus</i>	MBB4	1	s	in	1
Marsh Wren	<i>Cistothorus palustris</i>	MBB4	1	s	in	1
Mourning Dove	<i>Zenaid macroura</i>	MBB4	2	fo	in	1
Northern Waterthrush	<i>Seiurus noveboracensis</i>	MBB4	1	s	in	1
Red-eyed Vireo	<i>Vireo olivaceus</i>	MBB4	2	s	in	1
Red-winged Blackbird	<i>Agelaius phoeniceus</i>	MBB4	1	a	in	3
Red-winged Blackbird	<i>Agelaius phoeniceus</i>	MBB4	2	s	in	1
Rose-breasted Grosbeak	<i>Pheucticus ludovicianus</i>	MBB4	2	s	in	1
Song Sparrow	<i>Melospiza melodia</i>	MBB4	2	a	in	1
Yellow Warbler	<i>Dendroica petechia</i>	MBB4	1	s	in	1
Yellow Warbler	<i>Dendroica petechia</i>	MBB4	2	s	in	1

Breeding Evidence Codes

Observed

X-no breeding evidence
FO-flyover

Possible

H-Suitable habitat
S-Singing male

Probable

P-Pair
T-Territory (2 visits)
D-Display

V-Visiting nest

A-Agitated
B-Broodpatch
N-Nest building

Confirmed

DD-Distraction display

NU-Used nest

FY-Fledged young

AE-Adult entering/leaving nest

FS-Adult carrying fecal sac

CF-Adult carrying food

NE-Nest with eggs


NY-Nest with young

ROUND	TRANSECT	SPECIES or FEATURE	APPROX. LENGTH	COVER	COUNT	PHOTO	NOTES	HABITAT
1	none/scoping	Candidate Hibernacula	20m* 2m	Rock birm		yes	Linear rock feature, many crevices openings	open woodland
1	none/scoping	Candidate Hibernacula	5m* 5m	Rock birm		yes	circular rock pile, overgrown with many cracks.crevices	open woodland
1	none/scoping	Common gartersnake	30cm	long grass	1	yes	sunning on trail	open gravel and meadow
2	S1	none	-	-	-	-	-	-
2	S2	none	-	-	-	-	-	-
2	S3	none	-	-	-	-	-	-
2	S4	none	-	-	-	-	-	-
2	S5	none	-	-	-	-	-	-
3	S1	none	-	-	-	-	-	-
3	S2	none	-	-	-	-	-	-
3	S3	none	-	-	-	-	-	-
3	S4	none	-	-	-	-	-	-
3	S5	none	-	-	-	-	-	-

DRAFT

Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

STATION	DATE	SPECIES		
		Common Snapping Turtle	Midland Painted Turtle	Unknown Turtle Species
Turtle Habitat 1	29-Apr-15	2	26	
	08-May-15	3	25	
	14-May-15	2	19	
	28-May-15	2	12	
	11-Jun-15		22	
Turtle Habitat 1 total		9	104	0
Turtle Habitat 2	29-Apr-15			
	08-May-15			
	14-May-15			
	28-May-15			
	11-Jun-15		1	
Turtle Habitat 2 Total		0	1	0
Turtle Habitat 3	29-Apr-15		64	
	08-May-15	2	132	
	14-May-15		109	
	28-May-15	3	77	
	11-Jun-15	1	63	
Turtle Habitat 3 Total		6	445	0
Turtle Habitat 4	29-Apr-15		2	
	08-May-15		1	1
	14-May-15		2	
	28-May-15			
	11-Jun-15		2	
Turtle Habitat 4 Total		0	7	1
Turtle Habitat 5	29-Apr-15			
	08-May-15			
	14-May-15			
	28-May-15			
	11-Jun-15		1	
Turtle Habitat 5 total		0	1	0



COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-Rank	G-Rank	CVC (2010)
BIRDS							
Canada Goose	<i>Branta canadensis</i>				S5	G5	4
American Black Duck	<i>Anas rubripes</i>				S4	G5	2
Bald Eagle	<i>Haliaeetus leucocephalus</i>	SC			S2N,S4B	G5	1
Ruffed Grouse	<i>Bonasa umbellus</i>				S4	G5	2
Downy Woodpecker	<i>Picoides pubescens</i>				S5	G5	4
Pileated Woodpecker	<i>Dryocopus pileatus</i>				S5	G5	2
American Crow	<i>Corvus brachyrhynchos</i>				S5B	G5	2
Black-capped Chickadee	<i>Poecile atricapillus</i>				S5	G5	4
Pine Siskin	<i>Carduelis pinus</i>				S4B	G5	2
MAMMALS							
Eastern Cottontail	<i>Sylvilagus floridanus</i>				S5	G5	4
Eastern Gray Squirrel	<i>Sciurus carolinensis</i>				S5	G5	4
Red Squirrel	<i>Tamiasciurus hudsonicus</i>				S5	G5	3
Beaver	<i>Castor canadensis</i>				S5	G5	3
Meadow Vole	<i>Microtus pennsylvanicus</i>				S5	G5	3
Coyote	<i>Canis latrans</i>				S5	G5	3
American Mink	<i>Mustela vison</i>				S4	G5	2
White-tailed Deer	<i>Odocoileus virginianus</i>				S5	G5	3
Red Fox (cf.)	<i>Vulpes vulpes</i>				S5	G5	3

Legend:

SARO: Species at Risk Ontario
 COSEWIC: Committee on the status of endangered wildlife in Canada
 SARA: Species at Risk Act
 SC: Special Concern

Global Rank:


G1: Extremely rare globally
 G1G2: Extremely rare to very rare globally
 G2: Very rare globally
 G2G3: Very rare to uncommon globally
 G3: Rare to uncommon globally
 G3G4: Rare to common globally
 G4: Common globally
 G4G5: Common to very common globally
 G5: Very common globally; demonstrably secure

CVC Tiers:

1 - Species of Conservation Concern
 2 - Species of Interest
 3 - Species of Urban Interest
 4 - Secure Species
 5 - Non-native & Non-native Hybrid Species

Provincial Rank:

S1: Critically Imperiled—Critically imperiled in the province
 S2: Imperiled—Imperiled in the province, very few populations
 S3: Vulnerable—Vulnerable in the province, relatively few populations
 S4: Apparently Secure—Uncommon but not rare
 S5: Secure—Common, widespread, and abundant in the province
 S#B- Breeding status rank
 S#N- Non Breeding status rank



COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-Rank	G-Rank	CVC (2010)
Great Egret	<i>Ardea alba</i>				S2B	G5	1
Canada Goose	<i>Branta canadensis</i>				S5	G5	4
Wood Duck	<i>Aix sponsa</i>				S5	G5	2
American Black Duck	<i>Anas rubripes</i>				S4	G5	2
Mallard	<i>Anas platyrhynchos</i>				S5	G5	4
Belted Kingfisher	<i>Megaceryle alcyon</i>				S4B	G5	3
Northern Flicker	<i>Colaptes auratus</i>				S4B	G5	3
Blue Jay	<i>Cyanocitta cristata</i>				S5	G5	4
American Crow	<i>Corvus brachyrhynchos</i>				S5B	G5	2
Black-capped Chickadee	<i>Poecile atricapillus</i>				S5	G5	4
Red-breasted Nuthatch	<i>Sitta canadensis</i>				S5	G5	3
House Wren	<i>Troglodytes aedon</i>				S5B	G5	4
Golden-crowned Kinglet	<i>Regulus satrapa</i>				S5B	G5	2
Ruby-crowned Kinglet	<i>Regulus calendula</i>				S4B	G5	2
American Robin	<i>Turdus migratorius</i>				S5B	G5	4
European Starling	<i>Sturnus vulgaris</i>				SNA	G5	5
Song Sparrow	<i>Melospiza melodia</i>				S5B	G5	4
White-throated Sparrow	<i>Zonotrichia albicollis</i>				S5B	G5	3
American Goldfinch	<i>Carduelis tristis</i>				S5B	G5	4
Black bird species (mixed flock)							

Legend:

SARO: Species at Risk Ontario
 COSEWIC: Committee on the Status of Endangered wildlife in Canada
 SARA: Species at Risk Act
 ESA: Endangered Species Act
 END: Endangered
 THR: Threatened
 SC: special Concern
 NAR: Not At Risk
 NL: Not listed
 DD: Data Deficient

Global Rank:

G1: Extremely rare globally
 G1G2: Extremely rare to very rare globally
 G2: Very rare globally
 G2G3: Very rare to uncommon globally
 G3: Rare to uncommon globally
 G3G4: Rare to common globally
 G4: Common globally
 G4G5: Common to very common globally
 G5: Very common globally; demonstrably secure
 T: rank applies to a subspecies or variety

CVC Tiers:

1 - Species of Conservation Concern
 2 - Species of Interest
 3 - Species of Urban Interest
 4 - Secure Species
 5 - Non-native & Non-native Hybrid Species

Provincial Rank:

S1: Critically Imperiled—Critically imperiled in the province
 S2: Imperiled—Imperiled in the province, very few populations
 S3: Vulnerable—Vulnerable in the province, few populations
 S4: Apparently Secure—Uncommon but not rare
 S5: Secure—Common, widespread, and abundant in the province
 SX: Presumed extirpated
 SH: Possibly Extirpated (Historical)
 SNR: Unranked
 SU: Unrankable—Currently unrankable due to lack of information
 SNA: Not applicable—conservation status rank is not applicable
 S#S#: Range Rank—range of uncertainty about the status
 S#B- Breeding status rank
 S#N- Non Breeding status rank
 ?: Indicates uncertainty in the assigned rank

Appendix 14
Shorebird Habitat Assessment

DRAFT

Shorebird Habitat Assessment

Project: Hillsburgh Dam Project number: AA12-137A Observer(s): C.A. Ross Date: 05/08/2015

Location: Hillsburgh Dam Approximate Size of Census Area (X *X km): 0.5 x 0.1

Weather Conditions:

Temp (°C)	Wind*	Cloud Cover	Precipitation	Precipitation(24hrs)
17	3	19	None	none

*Beaufort Scale: 0-Calm (0 km/hr), 1-light Air (1-5km/hr), 2-Light Breeze (6-11km/hr), 3-gentle Breeze (12-19km/hr), 4-moderate Breeze (20-28km/hr), 5-fresh breeze (29-38km/hr), 6-strong breeze (39-49km/hr)

1. Habitat Availability (rank by area (ha) the habitats that are available to be used):

Sand Beach		Salt Marsh		Sewage Plant	
Sand Flat		Field		Mangrove	
Sand & Mud Flat		Brackish Pond		Other:	
Mud Flat	2	Temporary Pond		Other:	
Rocky Beach		Fresh Pond Or Lake	1	Other:	
Rocky Point		River		Other:	

2. Site is (rank by area (ha) if more than one):

On A Bay Or Estuary		An Inland Salt Lake Or Sea		Other:	
A Coastal Bay		On The Ocean Front			
A Lagoon		Principally An Inland Area	1		

3. Raptors observed:

Frequently		Infrequently		Never	x
Species observed:					

4. Disturbance by Humans:

>10 per day	x	5-10 per day		1-5 per day	
< 1 per day		Variable		unknown	

5. Rank major causes of disturbance:

People on foot	x	Vehicle Traffic	x	Pets	x
Boats		Hunting		Other	

6. Comments:

Very little mud or sand exposed, vegetated with pond lilies, cattails, No shorebirds observed during August survey.

APPENDIX 16. INCIDENTAL WILDLIFE LIST
Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

Project: AA12-137A

COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-Rank	G-Rank	CVC (2010)	Wellington County (2008)	Date(s) observed (2015)
BUTTERFLIES									
Canadian Tiger Swallowtail	<i>papilo canadensis</i>								May 28
AMPHIBIANS									
American Toad	<i>Anaxyrus americanus</i>				S5	G5	3		May 8
Spring Peeper	<i>Pseudacris crucifer</i>				S5	G5	3		April 15, April 29, May 8
Green Frog	<i>Lithobates clamitans</i>				S5	G5	3	✓	May 28
Northern Leopard Frog	<i>Lithobates pipiens</i>	NAR	NAR		S5	G5	3	✓	April 15, April 29, May 8, May 14, May 28
TURTLES									
Snapping Turtle	<i>Chelydra serpentina</i>	SC	SC	SC	S3	G5T5	1	✓	April 15, May 28
Midland Painted Turtle	<i>Chrysemys picta marginata</i>				S5	G5T5	3	✓	April 15
BIRDS									
Common Loon	<i>Gavia immer</i>	NAR	NAR		S5B, S5N	G5	2	✓	May 8
Great Blue Heron	<i>Ardea herodias</i>				S4	G5	3	✓	May 28
Great Egret	<i>Ardea alba</i>				S2B	G5	1	✓	May 28
Green Heron	<i>Butorides virescens</i>				S4B	G5	2	✓	May 28, August 5
Trumpeter Swan	<i>Cygnus buccinator</i>	NAR	NAR		S4	G4	1		April 29, May 28
Canada Goose	<i>Branta canadensis</i>				S5	G5	4		April 15, April 29, May 8, May 14, May 28
Wood Duck	<i>Aix sponsa</i>				S5	G5	2	✓	April 29, August 8
Mallard	<i>Anas platyrhynchos</i>				S5	G5	4	✓	April 29, May 8, May 28, Aug. 5, Sept. 25
Ring-necked Duck	<i>Aythya collaris</i>				S5	G5	3		April 15
Bufflehead	<i>Bucephala albeola</i>				S4	G5		✓	April 15
Common Merganser	<i>Mergus merganser</i>				S5B, S5N	G5	2	✓	April 15
Killdeer	<i>Charadrius vociferus</i>				S5B, S5N	G5	3		
Spotted Sandpiper	<i>Actitis macularia</i>				S5	G5	3	✓	
Ring-billed Gull	<i>Larus delawarensis</i>				S5B, S4N	G5	2	✓	August 5
Barred Owl	<i>Strix varia</i>				S5	G5	2		April 29
Belted Kingfisher	<i>Megasceryle alcyon</i>				S4B	G5	3	✓	April 15, April 29, August 5, September 29
Downy Woodpecker	<i>Picoides pubescens</i>				S5	G5	4	✓	April 15, May 8, May 28
Northern Flicker	<i>Colaptes auratus</i>				S4B	G5	3	✓	April 15
Alder Flycatcher	<i>Empidonax alnorum</i>				S5B	G5		✓	May 28
Eastern Phoebe	<i>Sayornis phoebe</i>				S5B	G5	3		April 15, May 8, May 28
Great Crested Flycatcher	<i>Myiarchus cineritus</i>				S4B	G5	3		May 28
Eastern Kingbird	<i>Tyrannus tyrannus</i>				S4B	G5	3		May 8, May 28, August 5
Tree Swallow	<i>Tachycineta bicolor</i>				S4B	G5	3		May 8, August 5
Blue Jay	<i>Cyanocitta cristata</i>				S5	G5	4		April 29, May 28
American Crow	<i>Corvus brachyrhynchos</i>				S5B	G5	2	✓	April 15, May 8, May 28
Black-capped Chickadee	<i>Poecile atricapillus</i>				S5	G5	4		April 15, April 29, May 8, May 28
White-breasted Nuthatch	<i>Sitta carolinensis</i>				S5	G5	3	✓	April 15

APPENDIX 16. INCIDENTAL WILDLIFE LIST
Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

Project: AA12-137A

COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-Rank	G-Rank	CVC (2010)	Wellington County (2008)	Date(s) observed (2015)
House Wren	<i>Troglodytes aedon</i>				S5B	G5	4	✓	May 8, May 28
Golden-crowned Kinglet	<i>Regulus satrapa</i>				S5B	G5	2	✓	April 15, April 29
Ruby-crowned Kinglet	<i>Regulus calendula</i>				S4B	G5	2	✓	April 15, April 29
Hermit Thrush	<i>Catharus guttatus</i>				S5B	G5	2		May 28
American Robin	<i>Turdus migratorius</i>				S5B	G5	4		April 15, April 29, May 8, May 28
Cedar Waxwing	<i>Bombycilla cedrorum</i>				S5B	G5	3	✓	May 28
Warbling Vireo	<i>Vireo gilvus</i>				S5B	G5	4		May 29
Red-eyed Vireo	<i>Vireo olivaceus</i>				S5B	G5	4	✓	May 30
Nashville Warbler	<i>Vermivora ruficapilla</i>				S5B	G5	2	✓	May 8
Yellow Warbler	<i>Dendroica petechia</i>				S5B	G5	4		May 8, May 25
Yellow-rumped Warbler	<i>Dendroica coronata</i>				S5B	G5	2	✓	May 8
Black-and-white Warbler	<i>Mniotilta varia</i>				S5B	G5	3	✓	May 8
American Redstart	<i>Setophaga ruticilla</i>				S5B	G5	3	✓	May 28
Northern Waterthrush	<i>Seiurus noveboracensis</i>				S5B	G5	3	✓	May 28
Common Yellowthroat	<i>Geothlypis trichas</i>				S5B	G5	4	✓	May 8, May 25
Northern Cardinal	<i>Cardinalis cardinalis</i>				S5	G5	4	✓	April 15, April 29, May 8, May 28
Rose-breasted Grosbeak	<i>Pheucticus ludovicianus</i>				S4B	G5	3		May 8
Indigo Bunting	<i>Passerina cyanea</i>				S4B	G5			May 28
Chipping Sparrow	<i>Spizella passerina</i>				S5B	G5	4	✓	April 15, May 8, May 28
Song Sparrow	<i>Melospiza melodia</i>				S5B	G5	4	✓	April 15, May 8, May 28
Swamp Sparrow	<i>Melospiza georgiana</i>				S5B	G5	4		April 29, May 8
White-throated Sparrow	<i>Zonotrichia albicollis</i>				S5B	G5	3		April 29, May 8, May 28
Red-winged Blackbird	<i>Agelaius phoeniceus</i>				S4	G5	4	✓	April 15, April 29, May 8, May 28
Common Grackle	<i>Quiscalus quiscula</i>				S5B	G5	4		April 15, April 29, May 28
Baltimore Oriole	<i>Icterus galbula</i>				S4B	G5	3		May 8, May 28
American Goldfinch	<i>Carduelis tristis</i>				S5B	G5	4		April 15, April 29, May 8
MAMMALS									
Eastern Chipmunk	<i>Tamias striatus</i>				S5	G5	3		May 8
Red Squirrel	<i>Tamiasciurus hudsonicus</i>				S5	G5	3	✓	April 15, May 8
Beaver	<i>Castor canadensis</i>				S5	G5	3		April 29, May 8, May 14
Muskrat	<i>Ondatra zibethicus</i>				S5	G5	3	✓	April 8
American Mink	<i>Mustela vison</i>				S4	G5	2		May 14

COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-Rank	G-Rank	CVC (2010)	Wellington County (2008)	Date(s) observed (2015)
-------------	-----------------	------	---------	------	--------	--------	------------	--------------------------	-------------------------

Legend:

SARO: Species at Risk Ontario
 COSEWIC: Committee on the Status of Endangered wildlife in Canada
 SARA: Species at Risk Act
 ESA: Endangered Species Act
 END: Endangered
 THR: Threatened
 SC: special Concern
 NAR: Not At Risk
 NL: Not listed
 DD: Data Deficient

Wellington County:

√ : Significant Species

PIF:

√: Priority Species

CVC Tiers:

- 1 - Species of Conservation Concern
- 2 - Species of Interest
- 3 - Species of Urban Interest
- 4 - Secure Species
- 5 - Non-native & Non-native Hybrid Species

Global Rank:

G1: Extremely rare globally
 G1G2: Extremely rare to very rare globally
 G2: Very rare globally
 G2G3: Very rare to uncommon globally
 G3: Rare to uncommon globally
 G3G4: Rare to common globally
 G4: Common globally
 G4G5: Common to very common globally
 G5: Very common globally; demonstrably secure
 T: rank applies to a subspecies or variety

Provincial Rank:

S1: Critically Imperiled—Critically imperiled in the province
 S2: Imperiled—Imperiled in the province, very few populations
 S3: Vulnerable—Vulnerable in the province, relatively few populations
 S4: Apparently Secure—Uncommon but not rare
 S5: Secure—Common, widespread, and abundant in the province
 SX: Presumed extirpated
 SH: Possibly Extirpated (Historical)
 SNR: Unranked
 SU: Unrankable—Currently unrankable due to lack of information
 SNA: Not applicable—A conservation status rank is not applicable
 S#S#: Range Rank— indicates range of uncertainty about the status
 S#B- Breeding status rank
 S#N- Non Breeding status rank
 ?: Indicates uncertainty in the assigned rank



APPENDIX 17. CANDIDATE SIGNIFICANT WILDLIFE HABITAT ASSESSMENT
Hillsburgh Dam Environmental Assessment, Natural Heritage – Existing Conditions

Project: A12-137AA

#	SIGNIFICANT WILDLIFE HABITAT (SWH)	CANDIDATE SWH CRITERIA	CRITERIA FOR SWH CONFIRMATION	SWH PROTECTED AREA	SITE ASSESSMENT DETAILS	CANDIDATE SWH	FIELD STUDIES REQUIRED/ COMPLETED	CONFIRMED SWH
SEASONAL CONCENTRATION AREAS OF ANIMALS								
1	Waterfowl stopover and Staging Areas (terrestrial)	- Fields with Sheet water in spring (incl. agricultural)	- Mixed species aggregations of 100 or more individuals confirms SWH	flooded field ecosite and 100-300m radius is the SWH	No Habitat matching Criteria identified in Study Area	No	None required.	No
2	Waterfowl Stopover and Staging (Aquatic)	- Ponds, marshes, lakes, bays, coastal inlets and watercourses and reservoirs SWTP & SWMP are not SWH - SHWH	- Aggregations of 100 or more listed species for 7 days (ie. >700 waterfowl use days) confirms SWH	Aquatic ecosite and 100m radius is the SWH	Hillsburgh pond is of sufficient size, shallow depth, and abundant aquatic vegetation.	Yes	Identified by CVC as SWH through spring surveys	Yes
3	Shorebird Migratory stopover	- Shorelines of Lakes, rivers, wetlands, beaches, bars; seasonally flooded, muddy and un-vegetated shoreline habitat	- 3 or more listed species and >1000 shorebird use days, or >100 whimbrel, confirms SWH	Shoreline ecosite and 100m radius is the SWH	No Habitat matching Criteria identified in Study Area, >5km from any Great Lake	No	Fall migration survey completed.	No
4	Raptor Wintering Area	- Combination of upland field and woodland habitat >20ha total (includes >15ha upland field) - least disturbed sites, idle, fallow or lightly grazed field/meadow best	- 1 or more Short-eared Owl, or, at least 10 individuals and 2 listed species for a minimum of 20 days, and 3 of 5 years, confirms SWH	Ecosite communities (field and woodland) is the SWH	No Habitat matching Criteria identified in Study Area	No	None required	No
5	Bat Hibernacula	- Caves, mine shafts, underground foundations, karsts - buildings are not SWH	- All sites with confirmed hibernating bats, confirms SWH	Ecosite and 200m radius is the SWH	No Habitat matching Criteria identified in Study Area	No	None required	No
6	Bat Maternity Colony	- All forested ecosites, FOD, FOC, FOM, SWD, SWM, SWC with >10/ha trees (>25cm DBH) in early stages of decay (class 1-3) - buildings are not SWH	- >10 Big Brown Bats, >20 Little Brown Myotis, >5 adult female Silver-haired Bats confirms SWH	Entire woodland or forest stand ELC ecosite containing colony is the SWH	Forested ecosites present in Study area with trees >25cm DBH.	Yes	Studies to be completed pre-construction if tree removal/damage to occur.	unknown
7	Turtle Wintering Area	- Areas with permanent water deep enough not to freeze, with mud/soft substrates	- 5 over-wintering Midland Painted Turtles, 1 or more Northern Map Turtle or Snapping Turtle confirms SWH	Mapped ELC ecosite, or deep pool element where turtles overwinter is the SWH	5 Candidate ponds identified in study area.	Yes	Basking surveys complete	Yes
8	Reptile Hibernaculum	- Sites below the frost line; rock barren, crevice and cave, talus, alvar, rock piles, slopes, stone fences and crumbling foundations	- Presence of hibernacula with minimum 5 individuals of 1 snake species/ individuals of 2 or more species confirms SWH - Congregations of a minimum of 5 snakes of 1 species/ individuals of 2 or more snake species, near potential hibernacula on sunny warm days in spring and fall confirms SWH	Feature hibernacula is located in, and 30m radius is the SWH	2 candidate hibernacula features identified in study area (rock piles in meadow openings-unknown depth)	yes	Snake basking transect surveys complete	No

APPENDIX 17. CANDIDATE SIGNIFICANT WILDLIFE HABITAT ASSESSMENT

Project: A12-137AA

Hillsburgh Dam Environmental Assessment, Natural Heritage – Existing Conditions

#	SIGNIFICANT WILDLIFE HABITAT (SWH)	CANDIDATE SWH CRITERIA	CRITERIA FOR SWH CONFIRMATION	SWH PROTECTED AREA	SITE ASSESSMENT DETAILS	CANDIDATE SWH	FIELD STUDIES REQUIRED/ COMPLETED	CONFIRMED SWH
9	Colonially-nesting Bird Habitat (cliff/bank)	- Eroding banks, sandy hills, borrow pits, steep slopes, sand piles, cliff faces, bridge abutments, silos, barns	- 1 or more nest sites with 8 or more Cliff Swallow or, 50 Bank Swallow and Rough-winged Swallow pairs during the breeding season.	Colony and 50m radius around peripheral nest is the SWH	No Habitat matching Criteria identified in Study Area	No	None required	No
10	Colonially-nesting Bird Habitat (Tree/shrub)	- Live or dead standing trees in wetlands, lakes, islands and peninsulas, occasionally shrubby and emergent vegetation	- 5 or more active Great-blue Heron or other listed species nests	Edge of the colony plus minimum 300m radius, or extent of the forest ecosite, or entire island <15ha is the SWH	No Habitat matching Criteria identified in Study Area	No	None required	No
11	Colonially-nesting Bird Habitat (Ground)	- Rocky islands or peninsulas within a lake or large river(natural or artificial)	- >25 active nests of Herring Gull, Ring-billed Gull, >5 active nests of Common Tern, or >2 active nests of Caspian Tern. 5 or more pairs of Brewer's Blackbird. Any active nesting colony of Little Gull, Great Black-backed Gull.	Edge of colony plus min 150m radius or extent of ELC ecosite, or island <3ha is the SWH	No Habitat matching Criteria identified in Study Area	No	None required	No
12	Migratory Butterfly Stopover Area	- At least 10ha, with undisturbed field/meadow and forest or woodland edge habitat present, within 5km of Lake Ontario.	- Presence of Monarch use days >5000 or >3000 where there is a mix of Monarch with Painted Ladies or White Admirals	Field/meadow and forest/woodland is the SWH	No Habitat matching Criteria identified in Study Area, >5km from any Great Lake	No	Fall migration survey completed.	No
13	Land bird Migratory Stopover Area	- Woodlots >5ha in size - within 5km of lake Ontario	- Use by >200 birds/day, with >35species, with at least 10sp recorded on 5 different survey dates.	Woodlot is the SWH	No Habitat matching Criteria identified in Study Area, >5km from any Great Lake	No	Fall migration survey completed.	No
14	Deer Yarding Areas	- ELC communities providing Thermal cover (FOM,FOC,SWM,SWC, CUP2, CUP3, FOD3, CUT)	- Deer yards are managed by MNRF, available through district offices and LIO.	LIO mapping	No Deer yarding areas identified on LIO Mapping	No	None required.	No
15	Deer Winter Congregation Areas	- All forested ecosites >100ha - Conifer Plantations <50ha may be used	- Deer management is the responsibility of the MNRF - Contact MNRF or LIO for known deer winter areas.	LIO mapping	No Deer Winter Congregation areas identified on LIO Mapping	No	None required.	No
RARE VEGETATION COMMUNITIES								
16	Cliffs & Talus Slopes	- Cliff: vertical to near vertical bedrock >3m in height - Talus slope: rock rubble at the base of a cliff made up of coarse rocky debris	- Confirm any ELC Vegetation Type for Cliffs or Talus Slopes	Area of ELC sites: TAO, TAS, TAT, CLO, CLS, CLT	No Habitat matching Criteria identified in Study Area	No	None required	No
17	Sand Barren	- Exposed, sparsely vegetated & caused by lack of moisture, fires and erosion.	- area >0.5ha in size - Confirm any ELC vegetation Type for Sand Barren - Not dominated by exotic or introduced species	Area of ELC ecosite is the SWH	No Habitat matching Criteria identified in Study Area	No	None required	No

APPENDIX 17. CANDIDATE SIGNIFICANT WILDLIFE HABITAT ASSESSMENT

Project: A12-137AA

Hillsburgh Dam Environmental Assessment, Natural Heritage – Existing Conditions

#	SIGNIFICANT WILDLIFE HABITAT (SWH)	CANDIDATE SWH CRITERIA	CRITERIA FOR SWH CONFIRMATION	SWH PROTECTED AREA	SITE ASSESSMENT DETAILS	CANDIDATE SWH	FIELD STUDIES REQUIRED/ COMPLETED	CONFIRMED SWH
18	Alvar	- Level, mostly un-fractured calcareous bedrock feature, overlain by a thin veneer or soil	- area >0.5ha in size - Field Studies that identify four of the five Alvar Indicator Species - Not dominated by exotic or introduced species	Area of ELC ecosite is the SWH	No Habitat matching Criteria identified in Study Area	No	None required	No
19	Old Growth Forest	- >30ha forests with at least 10ha interior habitat and multi-layered canopy	- Dominant Tree Species >140 years old - No recognizable signs forestry practices (old stumps)	Area of ELC ecosite is the SWH	No Habitat matching Criteria identified in Study Area	No	None required	No
20	Savannah	- Tall Grass Prairie Habitat with 25%-60% Tree cover - Remnant sites such as Railway Right of ways are not SWH	- No minimum size, and must be restored to a natural state. - Confirm one or more savannah indicator species - Not dominated by exotic or introduced species	Area of ELC ecosite is the SWH	No Habitat matching Criteria identified in Study Area	No	None required	No
21	Tallgrass Prairie	- Ground cover dominated by prairie grasses with <25% tree cover - Remnant sites such as Railway Right of ways are not SWH	- No minimum size, and must be restored to a natural state. - Confirm one or more prairie indicator species - Not dominated by exotic or introduced species	Area of ELC ecosite is the SWH	No Habitat matching Criteria identified in Study Area	No	None required	No
22	Other Rare Vegetation Communities	- All Provincially Rare S1, S2, S3 Vegetation Communities (Appendix M of SWHTG)	- Field Studies Confirming ELC vegetation type is a rare vegetation community	Area of ELC ecosite is the SWH	No Habitat matching Criteria identified in Study Area	No	None required	No
SPECIALIZED HABITAT FOR WILDLIFE								
23	Waterfowl Nesting Areas	- Upland Habitat, adjacent to Wetland ELC ecosites (except SWC, SWM) - Extends 120m from a wetland (>0.5ha) and any small wetlands (<0.5ha) within a cluster of at least 3 - Upland area at least 120m wide	- Presence of 3 or more nesting pairs of listed species excluding Mallards - Presence of 10 or more nesting pairs including mallards - Any active Black Duck nesting site	SWH may be greater than or less than 120m from the wetland edge and must provide enough habitat for waterfowl to successfully nest	Treed communities adjacent all wetlands/ponds, may provide nesting habitat	Yes	Breeding bird surveys completed	No
24	Bald Eagle or Osprey Nesting, Foraging and Perching Habitat	- Forest communities, adjacent to riparian areas - Osprey nests usually at top of tree - Bald Eagle nest usually in super canopy tree in a notch within canopy	- Studies confirm one or more active Bald Eagle or Osprey nest - Alternate nests included in SWH - Nests must be used annually, if found inactive, must be known inactive at least 3 years, or suspected unused for 5 years if unknown	Active nest plus 300m for Osprey Active nest plus 400-800m for Bald Eagle	Forested Habitat adjacent Hillsburgh pond may provide nesting opportunities for Osprey	Yes	Breeding Bird Surveys completed	No

APPENDIX 17. CANDIDATE SIGNIFICANT WILDLIFE HABITAT ASSESSMENT

Project: A12-137AA

Hillsburgh Dam Environmental Assessment, Natural Heritage – Existing Conditions

#	SIGNIFICANT WILDLIFE HABITAT (SWH)	CANDIDATE SWH CRITERIA	CRITERIA FOR SWH CONFIRMATION	SWH PROTECTED AREA	SITE ASSESSMENT DETAILS	CANDIDATE SWH	FIELD STUDIES REQUIRED/ COMPLETED	CONFIRMED SWH
25	Woodland Raptor Nesting Habitat	<ul style="list-style-type: none"> - Forested communities, forested swamp communities and cultural Plantations - Natural Forested/conifer plantations >30ha with >10ha interior habitat (200m buffer) 	<ul style="list-style-type: none"> - One or more active nest of listed species 	Nest protection radius: <ul style="list-style-type: none"> - Red-Shouldered Hawk, Northern Goshawk 400m - Barred Owl 200m - Broad-winged Hawk, Coopers Hawk 100m - Sharp-shinned Hawk 50 	Forested habitat may provide opportunities for woodland raptor nesting	Yes	No stick nests observed during SWH or Winter Wildlife Surveys	No
26	Turtle Nesting Areas	<ul style="list-style-type: none"> - Exposed Mineral soil (sand or gravel) adjacent (<100m) or within shallow marsh, shallow submerged, shallow floating, bog or fen communities - Located in open sunny areas, away from roads and less prone to predation - Municipal and provincial road shoulders are not SWH. 	<ul style="list-style-type: none"> - Confirm 5 or more nesting Midland Painted Turtles, 1 or more nesting Northern Map Turtle or Snapping Turtle 	Area or sites with exposed mineral soils, plus a radius of 30-100m around the nesting area is the SWH.	No Habitat matching Criteria identified in Study Area	No	None required	No
27	Seeps and Springs	<ul style="list-style-type: none"> - Areas where ground water comes to the surface - Any forested area within the headwaters of a stream or river system 	<ul style="list-style-type: none"> - Confirm site with 2 or more seeps/springs. 	Area of ELC forest ecosite containing seep/spring is the SWH	Seeps and springs possible within forested and wetland communities	Yes	ELC complete	No seeps or springs identified
28	Amphibian Breeding Habitat (woodland)	<ul style="list-style-type: none"> - Breeding pools within woodlands - Wetland, pond or pool >500m² within or adjacent (<120m) to a woodland. - Woodlands with permanent ponds, or those with water until mid-July more likely to be used. 	<ul style="list-style-type: none"> - Confirm Breeding population of 1 or more listed newt/salamander species, 2 or more of the listed frog species with at least 20 individuals (adults or egg masses), 2 or more of the listed frog species with call code levels of 3. - Wetland adjacent to woodlands includes travel corridor connecting features as SWH. 	Wetland area, plus 230m radius of woodland is the SWH.	Candidate habitat throughout study area, shallow ponds, woodland pools, marshes	yes	Amphibian Surveys complete	None confirmed as significant
29	Amphibian Breeding Habitat (Wetland)	<ul style="list-style-type: none"> - Swamp, marsh, fen, bog, open aquatic and shallow aquatic ELC communities. - Typically isolated from woodlands (>120m), but includes larger wetlands with primarily aquatic species (bull frogs) that are adjacent to woodlands. - Wetlands >500m² - Presence of shrubs & logs - Bullfrogs require permanent water bodies and abundant emergent vegetation. 	<ul style="list-style-type: none"> - Confirm Breeding populations of 1 or more listed newt/salamander species, or 2 or more listed frog/toad species with at least 20 individuals (adults or egg masses), or 2 or more listed frog/toad species with a call code level of 3 - Or any wetland with confirmed breeding Bullfrog. 	ELC ecosite and shoreline is the SWH Movement corridors (SWH) must be considered if this habitat is significant	No wetlands >120m from woodland habitat	No	Amphibian surveys complete	No

APPENDIX 17. CANDIDATE SIGNIFICANT WILDLIFE HABITAT ASSESSMENT

Project: A12-137AA

Hillsburgh Dam Environmental Assessment, Natural Heritage – Existing Conditions

#	SIGNIFICANT WILDLIFE HABITAT (SWH)	CANDIDATE SWH CRITERIA	CRITERIA FOR SWH CONFIRMATION	SWH PROTECTED AREA	SITE ASSESSMENT DETAILS	CANDIDATE SWH	FIELD STUDIES REQUIRED/ COMPLETED	CONFIRMED SWH
30	Area-sensitive Breeding Bird Habitat	<ul style="list-style-type: none"> - Habitats where interior breeding birds are breeding - Large mature(>60 years) forest stands or woodlots >30ha - Forest and swamp ELC communities - Interior habitat at least 200m from edge 	<ul style="list-style-type: none"> - Presence of nesting or breeding pairs of 3 or more of the listed species - Any site with Cerulean Warbler or Canada Warbler is SWH 	ELC ecosite is the SWH	No interior habitat identified in study area	no	None required	No
HABITATS OF SPECIES OF CONSERVATION CONCERN CONSIDERED SWH								
31	Marsh Bird Breeding Habitat	<ul style="list-style-type: none"> - Some meadow marsh, shallows submerged, shallow floating, mixed shallow floating, fen and bog communities (see SWH Ecoregion guide for specifics) - Nesting occurs in wetlands, all wetland habitat is considered with presence of shallow water with emergent aquatic vegetation - Green heron at edge of water sheltered by shrubs and trees. 	<ul style="list-style-type: none"> - 5 or more nesting pairs of Sedge Wren or Marsh Wren, 1 pair of Sandhill Crane, or breeding by any combination of 5 or more of the listed species - Any Wetland with 1 or more breeding pair Black Tern, Trumpeter Swan, Green Heron or Yellow Rail 	ELC ecosite is the SWH	Candidate habitat identified in study area.	Yes	Marsh Breeding Bird Surveys complete	No
32	Open Country Bird Breeding Habitat	<ul style="list-style-type: none"> - Grassland area >30ha (natural & cultural fields and meadows) - Grasslands not class 1 or 2 agriculture (no row crops or intensive hay or livestock pasturing) - Mature hayfields or pasture at least 5 years old 	<ul style="list-style-type: none"> - Nesting or breeding of 2 or more of the listed species - Field with 1 or more Short-eared Owls 	Contiguous ELC ecosite is the SWH	No Habitat matching Criteria identified in Study Area	No	None required	No
33	Shrub/Early Successional Bird Breeding Habitat	<ul style="list-style-type: none"> - Cultural thickets, savannah and woodland habitat - Large field area succeeding to shrub and thicket habitat >10ha in size - Patches of shrub ecosite may be complexed into larger old field ecosites for some species 	<ul style="list-style-type: none"> - Confirm nesting or breeding of 1 of the listed indicator species and at least 2 of the common species - Habitat with Yellow-breasted Chat Or Golden-winged Warbler is SWH 	SWH is contiguous ELC ecosite field/thicket area	No Habitat matching Criteria identified in Study Area	No	None required	No

APPENDIX 17. CANDIDATE SIGNIFICANT WILDLIFE HABITAT ASSESSMENT

Project: A12-137AA

Hillsburgh Dam Environmental Assessment, Natural Heritage – Existing Conditions

#	SIGNIFICANT WILDLIFE HABITAT (SWH)	CANDIDATE SWH CRITERIA	CRITERIA FOR SWH CONFIRMATION	SWH PROTECTED AREA	SITE ASSESSMENT DETAILS	CANDIDATE SWH	FIELD STUDIES REQUIRED/ COMPLETED	CONFIRMED SWH
34	Terrestrial Crayfish	<ul style="list-style-type: none"> - Meadow marsh, shallow marsh, swamp thicket, deciduous swamp and mixed swamp communities - Cultural meadow with inclusions of meadow marsh may be used - Wet edges of marshes and wet meadows should be surveyed for crayfish 	<ul style="list-style-type: none"> - Presence of 1 or more individuals of listed species or their chimneys in suitable habitat 	Area of ELC ecosite or Eco element area of meadow marsh or swamp within the larger ecosite area is the SWH	Candidate habitat identified in study area.	Yes	Incidental during ELC	No
35	Special Concern & Rare Wildlife Species	<ul style="list-style-type: none"> - All Special concern and Provincially Rare plant and animal species - Where an element occurrence is identified within a 1 or 10km grid for a species listed, linking candidate habitat on the site must be completed to ELC ecosites 	<ul style="list-style-type: none"> - Assessment/inventory of site for identified special concern or rare species completed during time of year when species is present or easily identifiable - Habitat must be easily mapped and cover an important life stage component (specific nesting habitat, foraging) 	SWH is the finest ELC scale that protects the form and function of the habitat	NHIC identified Carey's Sedge (<i>Carex careyana</i> S2) and Rugulose Grapefern (<i>Sceptridium rugulosum</i> S2) as occurring in the 1km square containing the study area. Habitat occurs in study area	Yes	Three season Botanical Survey	No
ANIMAL MOVEMENT CORRIDORS								
36	Amphibian Movement Corridor	<ul style="list-style-type: none"> - Corridors may occur in all ecosites associated with water - Presence of significant amphibian breeding indicates the requirement for identifying corridors - Movement corridors between breeding habitat and summer habitat 	<ul style="list-style-type: none"> - Corridors typically include areas with native vegetation, with several layers of vegetation, unbroken by roads, waterways or waterbodies are most significant - At least 15% of vegetation on both sides of the waterway or up to 200m wide of woodland habitat with gaps of <20m - Shorter corridors are more significant than longer, but amphibians must be able to get to and from their summer breeding habitat 	Corridor is the SWH	No Habitat matching Criteria identified in Study Area	No	None required	No
37	Deer Movement Corridor	<ul style="list-style-type: none"> - May occur in all forested ecosites - Determined when deer wintering habitat is confirmed as SWH 	<ul style="list-style-type: none"> - Corridors at least 200m wide with gaps <20m leading to wintering habitat - Unbroken by roads and residential areas - Shorter corridors are more significant 	Corridor is the SWH	No Habitat matching Criteria identified in Study Area	No	None required	No

APPENDIX 18. SPECIES WITH CONSERVATION STATUS ASSESSMENT
 Hillsburgh Dam Environmental Assessment, Natural Heritage – Existing Conditions

Project: AA12-137A

COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-RANK	BACKGROUND SOURCE	HABITAT REQUIREMENTS	SUITABLE HABITAT IN STUDY AREA	OBSERVED BY A & A
BUTTERFLIES									
Monarch	<i>Danaus plexippus</i>	SC	SC	SC	S2N,S4B	OBAO (2012)	Primarily found where milkweed and wildflowers exist, including abandoned farmland, along roadsides, and other open spaces (MNRF 2015).	Yes, large area of abundant milkweed occurs in the MEMM3 community	None observed during any surveys
West Virginia White	<i>Pieris virginiensis</i>	SC			S3	MNRF (Wellington list)	Generally prefer moist, deciduous woodlands. The larvae feed primarily on the leaves of the two-leaved toothwort (<i>Cardamine diphylla</i>), which is a small, spring-blooming plant of the forest floor (MNRF 2015).	Yes, host plant occurs in study area, in very small numbers in the SWMCM1-2 and SWMO1-1 communities	None observed during spring vegetation surveys
BIRDS									
Bald Eagle	<i>Haliaeetus leucocephalus</i>	SC	NAR		S2N,S4B	CVC (2013) ¹	Prefer deciduous and mixed-deciduous forest habitat close to large water bodies, including lakes and rivers; Nests in super canopy trees including Pine (MNRF 2015).	No, trees of sufficient size and species do not occur in study area. No suitably sized rivers or lakes in study area.	Observed during Winter Wildlife Survey
Bank Swallow	<i>Riparia riparia</i>	THR	THR		S4B	OBBA (2005)	Nesting occurs in a variety of natural and anthropogenic vertical banks, which often erode and change over time, including aggregate pits and the shores of large lakes and rivers	No, banks or aggregate pits of sufficient size, depth or texture, were not observed in the study area.	None observed during Breeding Bird season or incidentally
Barn Swallow	<i>Hirundo rustica</i>	THR	THR		S4B	OBBA (2005)	Farmland; lake/river shorelines; wooded clearings; urban populated areas; rocky cliffs; and wetlands. Nest inside or outside buildings; under bridges and in road culverts; on rock faces and in caves (MNRF 2015).	Yes, Bridges and dams in study area may provide suitable nesting habitat.	None observed during Breeding Bird season or incidentally
Bobolink	<i>Dolichonyx oryzivorus</i>	THR	THR		S4B	OBBA (2005)	Prefers open grasslands and hay fields. In migration and in winter uses freshwater marshes and grasslands (MNRF 2015).	No habitat of sufficient size or species composition occurs in the study area, habitat occurs outside the study area in agricultural fields	One male observed incidentally, not singing, flushed from MEMM3 community
Canada Warbler	<i>Wilsonia canadensis</i>	SC	THR	THR	S4B	OBBA (2005)	Prefers wet coniferous, deciduous and mixed forest types, with a dense shrub layer. Nests on the ground, on logs or hummocks, and uses dense shrub layer to conceal the nest (MNRF 2015).	Possible, areas of wet mixed coniferous occur throughout study area (SWCM3-2, SWCM1-2), no site access provided to suitable communities for further habitat suitability examination.	None observed during Breeding Bird season or incidentally
Canvasback	<i>Aythya valisineria</i>				S1B,S4N	CVC (2013) ¹	Canvasbacks are not known to breed in Ontario, occurring during spring and fall migration (MNRF 2015).	None	None observed incidentally

APPENDIX 18. SPECIES WITH CONSERVATION STATUS ASSESSMENT
 Hillsburgh Dam Environmental Assessment, Natural Heritage – Existing Conditions

Project: AA12-137A

COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-RANK	BACKGROUND SOURCE	HABITAT REQUIREMENTS	SUITABLE HABITAT IN STUDY AREA	OBSERVED BY A & A
Eastern Meadowlark	<i>Sturnella magna</i>	THR	THR		S4B	MNRF (Wellington List)	Generally prefers grassy pastures, meadows and hay fields. Nests are always on the ground and usually hidden in or under grass clumps (MNRF 2015).	No habitat of sufficient size or species composition occurs in the study area, habitat occurs outside the study area in agricultural fields	Observed singing outside study area during breeding bird surveys
Eastern Wood-pewee	<i>Contopus virens</i>	SC	SC		S4B	OBBA (2005)	Associated with deciduous and mixed forests. Within mature and intermediate age stands, prefers areas with little understory vegetation as well as forest clearings and edges (MNRF 2015).	Yes, Deciduous forest communities (FODM5-8) within study area provide breeding habitat for Eastern wood-pewee	Yes, at least two territories observed during Breeding Bird Surveys
Grasshopper Sparrow	<i>Ammodramus savannarum</i>		SC		S4B	OBBA (2005)	Prefers moderately open grasslands and prairies with patchy bare ground; avoids grasslands with extensive shrub cover (MNRF 2015).	No, grassland habitat in the study area has a high thatch cover, and abundant woody shrubs and forb cover.	None observed during Breeding Bird season or incidentally
Great Egret	<i>Ardea alba</i>				S2B	CVC (2013) ¹	Nests in woody vegetation, shrubs and trees; over water or on islands. Colony nester often mixed species aggregations, in lakes, ponds, marshes and estuaries (MNRF 2015).	Habitat observed is of insufficient size and low quality, no stick nests of wading birds observed in study area.	Observed incidentally during spring and fall migration period
Long-tailed Duck	<i>Clangula hyemalis</i>				S3B	CVC (2013) ¹	Breeds in subarctic and arctic wetlands. Nests adjacent to freshwater. Winters in coastal marine water and large freshwater lakes (MNRF 2015).	None	None observed incidentally
Wood Thrush	<i>Hylocichla mustelina</i>	SC	THR		S4B	OBBA (2005)	Nests in second-growth and mature deciduous and mixed forests, with saplings and well-developed understory layers. Prefers large forest mosaics, occasionally nests in small forest fragments (MNRF 2015).	Habitat observed is of insufficient size and low quality, with very low shrub cover	None observed during Breeding Bird season or incidentally
FISH									
Black Redhorse	<i>Moxostoma duquesnei</i>	THR	THR		S2	MNRF (Wellington List)	Generally lives in moderately sized rivers and streams, with generally moderate to fast currents (MNRF 2015).	No probable habitat in the study area, not known to occur in the Credit River.	None observed within the Study Area or identified through background review.
Redside Dace	<i>Clinostomus elongatus</i>	END	END		S2	MNRF (Wellington List)	Generally found in pools and slow-moving areas of small headwater streams with a moderate to high gradients (MNRF 2015).	Possible habitat in shaded areas of streams throughout study area.	None observed within the Study Area or identified through background review.

APPENDIX 18. SPECIES WITH CONSERVATION STATUS ASSESSMENT
 Hillsburgh Dam Environmental Assessment, Natural Heritage – Existing Conditions

Project: AA12-137A

COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-RANK	BACKGROUND SOURCE	HABITAT REQUIREMENTS	SUITABLE HABITAT IN STUDY AREA	OBSERVED BY A & A
Silver Shiner	<i>Notropis photogenis</i>	THR	TH	SC	S2S3	MNRF (Wellington List)	Generally prefer moderate to large, deep, relatively clear streams with swift currents, and moderate to high gradients (MNRF 2015).	No probable habitat in the study area, not known to occur in the Credit River.	None observed within the Study Area or identified through background review.
MAMMALS									
Eastern Small-footed Myotis	<i>Myotis leibii</i>	END	END	END		MNRF (Wellington list)	Overwinter in caves and mines that remain above 0 Maternity Roost primarily under loose rocks on exposed rock outcrops, crevices and cliffs, and occasionally in buildings, under bridges and highway overpasses and under tree bark (MNRF 2015).	Possible habitat in study area, FODM5-8 and FODM6 communities includes trees of sufficient size	None observed, no studies completed
Little Brown Myotis	<i>Myotis lucifugus</i>	END	END	END	S4	OMA (1994)	Overwinter in caves and mines that remain above 0 Maternal Roosts Often associated with buildings (attics, barns etc.). Occasionally found in trees (25-44 cm dbh) (MNRF 2015).	Possible habitat in study area, FODM5-8 and FODM6 communities includes trees of sufficient size	bats observed flying towards pond during evening bat banding observation conducted by MNRF
Northern Myotis	<i>Myotis septentrionalis</i>	END	END	END		MNRF (Wellington list)	Overwinter in caves and mines that remain above 0 Maternal Roosts: Often associated with cavities of large diameter trees (25-44 cm dbh). Occasionally found in structures (attics, barns etc.) (MNRF 2015).	Possible habitat in study area, FODM5-8 and FODM6 communities includes trees of sufficient size	None observed, no studies completed
MUSSELS									
Rainbow Mussel	<i>Villosa iris</i>	THR	END	END	S2S3	MNRF (Wellington List)	Abundant in shallow, well-oxygenated reaches of small- to medium-sized rivers and sometimes lakes, on substrates of cobble, gravel, sand and occasionally mud (MNRF 2015).	Possible habitat within streams throughout study area. Not known to occur in the Credit River Watershed.	None observed within the Study Area or identified through background review.
Wavy-rayed lampmussel	<i>Lampsilis fasciola</i>	THR	SC	SC	S1	MNRF (Wellington List)	Generally inhabit clear rivers and streams of a variety of sizes, where the water flow is steady and the substrate is stable (MNRF 2015).	Possible habitat within streams throughout study area. Not known to occur in the Credit River Watershed.	None observed within the Study Area or identified through background review.

APPENDIX 18. SPECIES WITH CONSERVATION STATUS ASSESSMENT
 Hillsburgh Dam Environmental Assessment, Natural Heritage – Existing Conditions

Project: AA12-137A

COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-RANK	BACKGROUND SOURCE	HABITAT REQUIREMENTS	SUITABLE HABITAT IN STUDY AREA	OBSERVED BY A & A
REPTILES									
Snapping Turtle	<i>Chelydra serpentina</i>	SC	SC	SC	S3	ORAA (2014)	Generally found in shallow waters with soft mud and leaf litter. Nesting occurs on gravelly or sandy areas along streams. Snapping Turtles often use man-made structures for nest sites, including roads (especially gravel shoulders), dams and aggregate pits (MNRF 2015).	All ponds in study area provide ideal habitat for snapping turtle, including MASO1-1, SAS1, and SAM1-8 communities	Yes, observed during turtle surveys, and incidentally.
Blanding's Turtle	<i>Emydonidea blandingii</i>	THR	THR	THR	S3	MNRF (Wellington List)	Found in freshwater lakes, permanent or temporary pools, slow-flowing streams, marshes and swamps. Preference for shallow water that is rich in nutrients, organic soil and dense vegetation (MNRF 2015).	Ponds, wetlands and streams in study area may provide habitat for Blanding's turtle.	None observed within study area during turtle surveys or incidentally.
Milksnake	<i>Lampropeltis triangulum</i>	SC	SC	SC	S3	MNRF (Wellington List)	Found in rural areas, frequently reported in and around buildings, especially old structures. Proximity to water, basking and nesting sites, and overwintering habitat is required (MNRF 2015).	May occur along farm field edges, and near older buildings in study area. Building foundations may provide overwintering habitat.	None observed within study area during snake surveys or incidentally.
Eastern Ribbonsnake	<i>Thamnophis sauritus</i>	SC	SC	SC	S3	MNRF (Wellington List)	Found along the edges of shallow ponds, streams, marshes, swamps, or bogs bordered by dense vegetation that provides cover (MNRF 2015).	Meadow marshes, and edges of ponds and streams may provide habitat,	None observed within study area during snake surveys or incidentally.
VASCULAR PLANTS									
American Chestnut	<i>Castanea dentata</i>	END	END	END	S2	MNRF (Wellington List)	Deciduous forest communities; this tree prefers arid forests with acid and sandy soils (MNRF 2015).	Deciduous forests of FODM5-8 and FOCM6 provide potential habitat for American Chestnut.	None observed during Botanical Survey.
American Ginseng	<i>Panax quinquefolius</i>	END	END	END	S2	MNRF (Wellington List)	Rich, moist, undisturbed and relatively mature deciduous woods in areas of neutral soil (such as over limestone or marble bedrock) (MNRF 2015).	Deciduous forests of FODM5-8 provide potential habitat for American Ginseng.	None observed during Botanical Survey.
Butternut	<i>Juglans cinerea</i>	END	END	END	S3?	MNRF (Wellington List)	Rich, moist, and well-drained soils often found along streams. May also occur on well-drained gravel sites, especially those made up of limestone. Seldom found on dry, rocky and sterile soils. In Ontario, the Butternut generally grows alone or in small groups in deciduous forests as well as in hedgerows (MNRF 2015).	Habitat present along mineral soil edge of stream communities SWMO3-3 and FODM7-7, as well as within communities SWDM2-1 and FODM8-1.	None observed during Botanical Survey.

APPENDIX 18. SPECIES WITH CONSERVATION STATUS ASSESSMENT
 Hillsburgh Dam Environmental Assessment, Natural Heritage – Existing Conditions

Project: AA12-137A

COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-RANK	BACKGROUND SOURCE	HABITAT REQUIREMENTS	SUITABLE HABITAT IN STUDY AREA	OBSERVED BY A & A
Hill's Pondweed	<i>Potamogeton hillii</i>	SC	SC		S2	MNRF (Wellington List)	Generally grows in clear, cold ponds and slow-moving streams where the water is alkaline (MNRF 2015).	All ponds in study area provide possible habitat, including SAS1, and SAM1-8, although temperature are likely too warm.	None observed during Botanical Survey.
Carey's Sedge	<i>Carex careyana</i>				S2	NHIC	Grows in dry to moist rich deciduous upland forests (NatureServe 2015).	Deciduous forests of FODM5-8 and FOCM6 provide potential habitat.	None observed during Botanical Survey.
Rugulose Grapefern	<i>Sceptridium rugulosum</i>				S2	NHIC	Grows in sandy to silty soil in open fields, young successional forests or at the edge of forests (Wagner and Wagner 1982).	The edge of deciduous forests of FODM5-8 and FOCM6 provide potential habitat.	None observed during Botanical Survey.

1-observed outside the breeding season

References:

MNRF, 2015. Wellington County Upper Tier Species at Risk. Ministry of Natural Resources and Forestry. Provided February, 2015.

Mccrimmon, Jr., Donald A., John C. Ogden and G. Thomas Bancroft. 2011. Great Egret (*Ardea alba*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/570>

NatureServe, 2015. *Carex careyana* – Torr. Ex Dewey. NatureServe Explore. An Online Encyclopedia of Life. Available at: <http://explorer.natureserve.org/servlet/NatureServe?searchName=CAREX+CAREYANA>

Vickery, Peter D. 1996. Grasshopper Sparrow (*Ammodramus savaanarum*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/239>

Robertson, Gregory J. and Jean-Pierre L. Savard. 2002. Long-tailed Duck (*Clangula hyemalis*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/651>

Wagner, W. H. and F. S. Wagner. 1982. *Botrychium rugulosum* (Ophioglossaceae), a newly recognized species of evergreen grapefern in the Great Lakes Area of North America. Contributions from the University of Michigan Herbarium.

SEGMENT 1		DESCRIPTION
Fish Community Classification ¹ :	Cold Water	The stream reach runs through a White Cedar Conifer Mineral Coniferous Swamp, and has a natural meander pattern with areas of flat water, gentle runs and small riffles. Substrate is mostly sand with a lesser component of fines and small amounts of gravel, cobble and boulders. The instream cover consists of aquatic vegetation and woody debris, but is predominantly open. The stream bank is mostly stable and vegetated, with some minor areas of undercut bank or bare soil. Small fish of approximately 7cm were seen within this stream segment on two occasions, species unknown. Minor log jams were observed.
Mean Channel Width ² (m):	2m	
Mean Channel Depth ³ (m):	0.7m	
Mean Water Depth ⁴ (m):	0.4m	
Stream Shading %:	90%	FISH SPECIES
		Brook Trout, Central Mudminnow, White Sucker, Creek Chub, Brook Stickleback, Mottled Sculpin, Slimy Sculpin,
SEGMENT 2		DESCRIPTION
Fish Community Classification ¹ :	Cold Water	The stream reach runs through a Mixed Willow Organic Thicket Swamp Type in a braided flow pattern, with a poorly defined channel throughout much of the segment length. The substrate is mostly fines (silt and organic) with minor components of sand, gravel cobble and boulders. The flow pattern is flat with areas of deeper pools. In stream cover is high, consisting of emergent vegetation, submergent vegetation, woody debris and algae. There are minor areas of channel hardening at the outflow into the Hillsburgh Pond. There are also potential barriers to fish passage at the outflow into the Hillsburgh Pond. Barriers consist of a presumed fish gate that is likely intended to keep common carp out of the upstream reach, log jams and poor culvert design could also make passage difficult for non-jumping fish.
Mean Channel Width ² (m):	Variable	
Mean Channel Depth ³ (m):	Variable	
Mean Water Depth ⁴ (m):	0.5m	
Stream Shading %:	60%	FISH SPECIES
		Cyprinid sp., Rock Bass, Large Mouth Bass, Brook Trout, Slimy Sculpin, Central Mudminnow, White Sucker, Creek Chub, Brook Stickleback, Bluntnose Minnow, Pumpkinseed, Round Goby,
SEGMENT 3		DESCRIPTION
Fish Community Classification ¹ :	Warm Water	This segment is the open water community of the Hillsburgh Pond. The Pond is fed by two upstream tributaries and flows out at the Hillsburgh dam, under the Station Street bridge. Substrate composition of the pond is unknown. A variety of emergent, submergent and floating plants are present within the segment, particularly around the edges of the pond.
Mean Channel Width ² (m):	Open Water	
Mean Channel Depth ³ (m):	Unknown	
Mean Water Depth ⁴ (m):	Unknown	
Stream Shading %:	5%	FISH SPECIES
		Rock Bass, Large Mouth Bass, Bluntnose Minnow, Pumpkinseed, Round Goby.
SEGMENT 4		DESCRIPTION
Fish Community Classification ¹ :	Cold Water	Segment 4 runs through a Fresh-Moist Manitoba Maple Lowland Deciduous Forest, within the Downtown area of Hillsburgh. The flow pattern within the segment is mostly flat, with areas of pools riffles and runs. The channel has a gentle meander with some channelization and channel hardening. Substrate is a mixture of primarily sand, cobble and boulders, with a lesser component of fines and gravel, and may provide trout spawning habitat. In stream cover consists of 20% woody debris 10% aquatic vegetation and 70% open. The bank is mostly stable and vegetated, with some areas of erosion evident. A single 8cm fish of unknown species was observed in the segment. There are numerous culverts and stream crossings within the segment, but none appear to be barriers to fish passage.
Mean Channel Width ² (m):	2.5m	
Mean Channel Depth ³ (m):	0.75m	
Mean Water Depth ⁴ (m):	0.4m	
Stream Shading %:	80%	FISH SPECIES
		Brook Trout, Brown Trout, Creek Chub, White Sucker, Eastern Blacknose Dace, Bluntnose Minnow, Brook Stickleback, Central Mudminnow, Longnose Dace, Rock Bass, Round Goby.
SEGMENT 5		DESCRIPTION
Fish Community Classification ¹ :	Warm Water	This is a small segment between the Hillsburgh Pond and the Ainsworth Pond, including the Station Street bridge and outflow structure from the Hillsburgh Pond. The segment is 30% channelized with concrete wall and a sloped concrete pad as substrate. The flow pattern in the segment consists of a 1m fall from the Hillsburgh Pond and two smaller 0.3m falls, as well as a series of riffles and runs. Substrate is primarily the concrete slab, as well as gravel, cobble and boulder further downstream. MNR records have recorded trout spawning redds in this location. The outflow structure from the Hillsburgh Pond creates a barrier to fish passage.
Mean Channel Width ² (m):	3.5m	
Mean Channel Depth ³ (m):	0.45m	
Mean Water Depth ⁴ (m):	0.4m	
Stream Shading %:	65%	FISH SPECIES
		Bluntnose Minnow, Brook Trout, Creek Chub, Eastern Blacknose Dace, Golden Shiner, Largemouth Bass, Pumpkinseed, Rock Bass, Round Goby.

SEGMENT 6		DESCRIPTION
Fish Community Classification ¹ :	Warm Water	This segment is the shallow open water community of the Ainsworth Pond. Substrate of the community is unknown; aquatic cover includes emergent vegetation, floating vegetation, submergent vegetation and woody debris. The banks around the pond appear stable.
Mean Channel Width ² (m):	Open Water	
Mean Channel Depth ³ (m):	Unknown	
Mean Water Depth ⁴ (m):	Unknown	
Stream Shading %:	15%	FISH SPECIES
		Bluntnose Minnow, Brook Trout, Creek Chub, Eastern Blacknose Dace, Largemouth Bass, Rock Bass, White Sucker.
SEGMENT 7		DESCRIPTION
Fish Community Classification ¹ :	Cold Water	This segment consists of the two outflows from the Ainsworth Pond, which reconnect in the downstream portion of the segment. The main outfall from the pond consists of a series of two 0.6m fall and represents a complete barrier to fish passage. The second outfall appears to be temporary in nature and is constructed or reinforced by sandbags and plastic lining. This outfall consists of a series of smaller drops of 0.2m or less and may be passable by jumping fish under certain water levels. The flow pattern below the outfalls consist of pools, riffles and runs and the channel morphology is a gentle meander with straight sections. Substrate is a mixture of sand, gravel, cobble and boulders and may represent possible spawning habitat for trout.
Mean Channel Width ² (m):	2.5m	
Mean Channel Depth ³ (m):	0.5m	
Mean Water Depth ⁴ (m):	0.35m	
Stream Shading %:	65%	FISH SPECIES
		Banded Killifish, Brook Trout, Common Shiner, Creek Chub, Central Mudminnow, Eastern Blacknose Dace, Golden Shiner, Largemouth Bass, Pumpkinseed, Rock Bass, Round Goby, White Sucker.
SEGMENT 8		DESCRIPTION
Fish Community Classification ¹ :	Cold Water	This segment runs along the edge of a Tamarack White Cedar Treed Fen. The segment has a gentle meander with minor amounts of in-stream cover consisting of emergent plants, submergent plants and woody debris. Substrate is a mixture of sand, gravel, cobble and boulders and may represent possible spawning habitat for trout.
Mean Channel Width ² (m):	2m	
Mean Channel Depth ³ (m):	0.5m	
Mean Water Depth ⁴ (m):	0.3m	
Stream Shading %:	40%	FISH SPECIES
		Bluntnose Minnow, Brook Trout, Central Mudminnow, Common Shiner, Creek Chub, Eastern Blacknose Dace, Golden Shiner, Largemouth Bass, Pumpkinseed, Rock Bass, Round Goby, White Sucker.
SEGMENT 9		DESCRIPTION
Fish Community Classification ¹ :	Cold Water	This segment is a wide stream section leading to the Rudd Pond. The channel morphology is meandering with a flow pattern of riffles and runs upstream, and flat in the downstream portion. There is a high coverage of submergent vegetation consisting of watercress and <i>vallisneria americana</i> . Substrate is a mixture of sand, gravel, cobble and boulders in the upstream portion, transitioning to mostly sand and fines in the downstream portion.
Mean Channel Width ² (m):	4m	
Mean Channel Depth ³ (m):	0.7m	
Mean Water Depth ⁴ (m):	0.5m	
Stream Shading %:	60%	FISH SPECIES
		Central Mudminnow, Common Shiner, Creek Chub, Rock Bass.
SEGMENT 10		DESCRIPTION
Fish Community Classification ¹ :	Cold Water	This segment is the shallow open water community of the Rudd Pond. Substrate of the community is unknown; aquatic cover includes emergent vegetation, floating vegetation, submergent vegetation and woody debris. The banks around the pond appear stable.
Mean Channel Width ² (m):	Open Water	
Mean Channel Depth ³ (m):	Unknown	
Mean Water Depth ⁴ (m):	Unknown	
Stream Shading %:	5%	FISH SPECIES
		Bluntnose Minnow, Largemouth Bass, Pumpkinseed, Rock Bass, White Sucker.
SEGMENT 11		DESCRIPTION
Fish Community Classification ¹ :	Cold Water	This segment consists of the outflows from the Rudd Pond, and the downstream watercourse to Wellington Rd. 22. The outfall from the pond consists of a series of two 0.30m falls and a sloped section of concrete slab. This outfall likely represents a complete barrier to fish passage, although jumping fish may be able to pass under certain water level. The flow pattern below the outfalls consists of pools, riffles and runs and the channel morphology is a gentle meander with straight sections. Substrate is a mixture of sand, gravel, cobble and boulders and may represent possible spawning habitat for trout.
Mean Channel Width ² (m):	2.5m	
Mean Channel Depth ³ (m):	0.8m	
Mean Water Depth ⁴ (m):	0.4m	
Stream Shading %:	85%	FISH SPECIES
		Bluntnose Minnow, Brook Trout, Common Shiner, Eastern Blacknose Dace, Golden Shiner, Largemouth Bass, Pumpkinseed, Rock Bass, White Sucker.
SEGMENT 12		DESCRIPTION

Fish Community Classification ¹ :	Cold Water	This segment is primarily outside the study area, upstream on the southwest tributary. The section of watercourse is ephemeral, drying during periods of the summer. The channel morphology is straight with stable vegetated banks.
Mean Channel Width ² (m):	1.5m	
Mean Channel Depth ³ (m):	0.25m	
Mean Water Depth ⁴ (m):	Dry	
Stream Shading %:	75%	FISH SPECIES
SEGMENT 13		DESCRIPTION
Fish Community Classification ¹ :	Cold Water	This segment is outside the study area, upstream on the northeast tributary. The section is mostly within a large forested area. The substrate is a mixture of fines, sand and gravel with minor components of cobble and boulder and may represent possible spawning habitat for trout.
Mean Channel Width ² (m):	2.5m	
Mean Channel Depth ³ (m):	0.75m	
Mean Water Depth ⁴ (m):	0.45m	
Stream Shading %:	90%	FISH SPECIES
SEGMENT 14		DESCRIPTION
Fish Community Classification ¹ :	Cold Water	This segment is outside the study area, downstream of the main tributary. The section is mostly within a large forested area, leading to a dammed pond. The substrate is a mixture of sand, gravel, cobble and boulder and may represent possible spawning habitat for trout. The banks are vegetated and stable. The culvert crossing at Wellington 22 has a concrete slab bottom and would allow fish passage under normal levels.
Mean Channel Width ² (m):	2.5m	
Mean Channel Depth ³ (m):	0.75m	
Mean Water Depth ⁴ (m):	0.4m	
Stream Shading %:	80%	FISH SPECIES
		Rock Bass, White Sucker

1. Fish Community Classification based in Erin Service and Settlement Master Plan: Phase 1 - Environmental Component (Erin SSMP 2011).
2. Mean Channel width measured as the width of the wetted bank.
3. Mean Channel Depth is the depth of the channel from the low point to the top of the wetted bank.
4. Mean Water Depth is the depth of the water at the time of observation (October 19th 2015).
5. Fish Species is compiled from data provided by MNRF and CVC.

DRAFT

APPENDIX 20. BACKGROUND WILDLIFE LIST
Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

SOURCE	CVC (most recent observation indicated)	COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-Rank	G-Rank	PIF (BCR 13)	CVC Tier (2010)	Wellington County (2008)	
		INSECTS AT RISK										
OBAO (2012)		Monarch	<i>Danaus plexippus</i>	SC	SC	SC	S2N,S4B	G4T3				
		AMPHIBANS										
ORAA (2009)	I (2012)	American Toad	<i>Anaxyrus americanus</i>				S5	G5		3		
ORAA (2009)	I (2003)	Gray Treefrog	<i>Hyla versicolor</i>				S5	G5				
ORAA (1994)		Spring Peeper	<i>Pseudacris crucifer</i>				S5	G5		3		
ORAA (1990)	I (2014)	Green Frog	<i>Lithobates clamitans</i>				S5	G5		3		
	I (2008)	Northern Leopard Frog	<i>Lithobates pipiens</i>	NAR	NAR		S5	G5		3		
		SNAKES AND LIZARDS										
ORAA (1999)		Eastern Gartersnake	<i>Thamnophis sirtalis sirtalis</i>				S5	G5T5		4		
		TURTLES										
ORAA (2014)		Snapping Turtle	<i>Chelydra serpentina</i>	SC	SC	SC	S3	G5T5		1	✓	
ORAA (2013)	I (2014)	Midland Painted Turtle	<i>Chrysemys picta marginata</i>				S5	G5T5		3		
		BIRDS										
	FM (2012)	Pied-billed Grebe	<i>Podilymbus podiceps</i>				S4B,S4N	G5		2	✓	
	FM (2012)	Double-crested Cormorant	<i>Phalacrocorax auritus</i>	NAR	NAR		S5B	G5		2	✓	
OBBA (2005)	I (2014)	Great Blue Heron	<i>Ardea herodias</i>				S4	G5		3	✓	
	I (2014)	Great Egret	<i>Ardea alba</i>				S2B	G5		1	✓	
OBBA (2005)	FM (2012)	Green Heron	<i>Butorides virescens</i>				S4B	G5		2	✓	
	SWH (2011)	Tundra Swan	<i>Cygnus columbianus</i>				S4	G5		1		
OBBA (2005)	FM (2012)	Trumpeter Swan	<i>Cygnus buccinator</i>	NAR	NAR		S4	G4		1	✓	
OBBA (2005)	BB (2009)	Canada Goose	<i>Branta canadensis</i>				S5	G5		4		
OBBA (2005)	FM (2012)	Wood Duck	<i>Aix sponsa</i>				S5	G5		2	✓	
OBBA (2005)	FM (2012)	Green-winged Teal	<i>Anas crecca</i>				S4	G5		2	✓	
	FM (2012)	American Black Duck	<i>Anas rubripes</i>				S4	G5		2	✓	
OBBA (2005)	NB (2014)	Mallard	<i>Anas platyrhynchos</i>				S5	G5		4		
	FM (2012)	Blue-winged Teal	<i>Anas discors</i>				S4	G5		2	✓	
	FM (2012)	Gadwall	<i>Anas strepera</i>				S4	G5		2	✓	
	SM (2012)	Canvasback	<i>Aythya valisineria</i>				S1B,S4N	G5		1	✓	
	SM (2012)	Ring-necked Duck	<i>Aythya collaris</i>				S5	G5		3	✓	
	SWH (2011)	Long-tailed Duck	<i>Clangula hyemalis</i>				S3B	G5		1		
	SWH (2011)	Bufflehead	<i>Bucephala albeola</i>				S4	G5				
	FM (2012)	Hooded Merganser	<i>Lophodytes cucullatus</i>				S5B,S5N	G5		2	✓	
	FM (2012)	Common Merganser	<i>Mergus merganser</i>				S5B,S5N	G5		2	✓	
	FM (2012)	Red-breasted Merganser	<i>Mergus serrator</i>				S4B,S5N	G5		2	✓	
OBBA (2005)	BB 2009	Turkey Vulture	<i>Cathartes aura</i>				S5B	G5		3	✓	
	SWH (2011)	Osprey	<i>Pandion haliaetus</i>				S5B	G5		2	✓	
		Bald Eagle	<i>Haliaeetus leucocephalus</i>	SC	NAR		S2N,S4B	G5	✓	1	✓	
OBBA (2005)	SWH (2011)	Northern Harrier	<i>Circus cyaneus</i>	NAR	NAR		S4B	G5	✓	2	✓	
OBBA (2005)		Sharp-shinned Hawk	<i>Accipiter striatus</i>	NAR			S5	G5		3	✓	
	SWH (2011)	Cooper's Hawk	<i>Accipiter cooperii</i>	NAR	NAR		S4	G5		2	✓	

APPENDIX 20. BACKGROUND WILDLIFE LIST
Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

Project: AA12-137A

SOURCE	CVC (most recent observation indicated)	COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-Rank	G-Rank	PIF (BCR 13)	CVC Tier (2010)	Wellington County (2008)
OBBA (2005)		Northern Goshawk	<i>Accipiter gentilis</i>	NAR	NAR		S4	G5T5		2	✓
OBBA (2005)		Broad-winged Hawk	<i>Buteo platypterus</i>				S5B	G5		2	✓
OBBA (2005)		Red-tailed Hawk	<i>Buteo jamaicensis</i>	NAR	NAR		S5	G5		4	
OBBA (2005)		American Kestrel	<i>Falco sparverius</i>				S4	G5	✓	3	✓
OBBA (2005)	NB (2009)	Ruffed Grouse	<i>Bonasa umbellus</i>				S4	G5		2	
OBBA (2005)	BB (2009)	Wild Turkey	<i>Meleagris gallopavo</i>				S5	G5		3	
OBBA (2005)		Virginia Rail	<i>Rallus limicola</i>				S5B	G5		2	
	BB (2009)	Semipalmated Plover	<i>Charadrius semipalmatus</i>				S4B,S4N	G5		4	
OBBA (2005)	FM (2012)	Killdeer	<i>Charadrius vociferus</i>				S5B,S5N	G5		3	
	SM (2012)	Greater Yellowlegs	<i>Tringa melanoleuca</i>				S4B,S4N	G5		4	
	SM (2012)	Lesser Yellowlegs	<i>Tringa flavipes</i>				S4B,S4N	G5		4	
	BB (2009)	Solitary Sandpiper	<i>Tringa solitaria</i>				S4B	G5			
OBBA (2005)	I (2013)	Spotted Sandpiper	<i>Actitis macularius</i>				S5	G5		3	
	SWH (2011)	Least Sandpiper	<i>Calidris minutilla</i>				S4B,S5N	G5		4	
OBBA (2005)	FM (2012)	Wilson's Snipe	<i>Gallinago delicata</i>				S5B	G5		2	
OBBA (2005)	I (2011)	American Woodcock	<i>Scolopax minor</i>				S4B	G5		2	
	I (2012)	Ring-billed Gull	<i>Larus delawarensis</i>				S5B,S4N	G5		2	✓
	SM (2012)	Herring Gull	<i>Larus argentatus</i>				S5B,S5N	G5		2	✓
OBBA (2005)		Rock Pigeon	<i>Columba livia</i>				SNA	G5		5	
OBBA (2005)	BB (2009)	Mourning Dove	<i>Zenaida macroura</i>				S5	G5		4	
OBBA (2005)		Black-billed Cuckoo	<i>Coccyzus erythrophthalmus</i>				S5B	G5	✓	2	✓
OBBA (2005)		Yellow-billed Cuckoo	<i>Coccyzus americanus</i>				S4B	G5		2	✓
OBBA (2005)		Eastern Screech-Owl	<i>Megascops asio</i>	NAR	NAR		S4	G5		3	
OBBA (2005)		Great Horned Owl	<i>Bubo virginianus</i>				S4	G5		3	
OBBA (2005)		Long-eared Owl	<i>Asio otus</i>				S4	G5		2	✓
OBBA (2005)		Ruby-throated Hummingbird	<i>Archilochus colubris</i>				S5B	G5		3	
OBBA (2005)	NB (2009)	Belted Kingfisher	<i>Megaceryle alcyon</i>				S4B	G5	✓	3	
OBBA (2005)		Yellow-bellied Sapsucker	<i>Sphyrapicus varius</i>				S5B	G5		2	✓
OBBA (2005)	BB (2009)	Downy Woodpecker	<i>Picooides pubescens</i>				S5	G5		4	
OBBA (2005)	BB (2009)	Hairy Woodpecker	<i>Picooides villosus</i>				S5	G5		3	✓
OBBA (2005)	BB (2009)	Northern Flicker	<i>Colaptes auratus</i>				S4B	G5	✓	3	
OBBA (2005)	BB (2009)	Pileated Woodpecker	<i>Dryocopus pileatus</i>				S5	G5		2	✓
OBBA (2005)		Eastern Wood-pewee	<i>Contopus virens</i>	SC	SC		S4B	G5	✓	1	✓
OBBA (2005)	BB (2009)	Alder Flycatcher	<i>Empidonax alnorum</i>				S5B	G5			
OBBA (2005)		Willow Flycatcher	<i>Empidonax traillii</i>				S5B	G5	✓	3	✓
OBBA (2005)		Least Flycatcher	<i>Empidonax minimus</i>				S4B	G5		3	✓
OBBA (2005)	BB (2009)	Eastern Phoebe	<i>Sayornis phoebe</i>				S5B	G5		3	
OBBA (2005)	BB (2009)	Great Crested Flycatcher	<i>Myiarchus crinitus</i>				S4B	G5		3	
OBBA (2005)	BB (2009)	Eastern Kingbird	<i>Tyrannus tyrannus</i>				S4B	G5	✓	3	✓
OBBA (2005)		Horned Lark	<i>Eremophila alpestris</i>				S5B	G5		3	
		Purple Martin	<i>Progne subis</i>				S4B	G5		2	
OBBA (2005)	BB (2009)	Tree Swallow	<i>Tachycineta bicolor</i>				S4B	G5		3	
OBBA (2005)	BB (2009)	Northern Rough-winged Swallow	<i>Stelgidopteryx serripennis</i>				S4B	G5		3	
OBBA (2005)		Bank Swallow	<i>Riparia riparia</i>	THR	THR		S4B	G5	✓	1	✓
OBBA (2005)		Cliff Swallow	<i>Petrochelidon pyrrhonota</i>				S4B	G5		3	✓

APPENDIX 20. BACKGROUND WILDLIFE LIST
Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

SOURCE	CVC (most recent observation indicated)	COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-Rank	G-Rank	PIF (BCR 13)	CVC Tier (2010)	Wellington County (2008)
OBBA (2005)		Barn Swallow	<i>Hirundo rustica</i>	THR	THR		S4B	G5		1	✓
OBBA (2005)	BB (2009)	Blue Jay	<i>Cyanocitta cristata</i>				S5	G5		4	
OBBA (2005)	BB (2009)	American Crow	<i>Corvus brachyrhynchos</i>				S5B	G5		2	
OBBA (2005)		Common Raven	<i>Corvus corax</i>				S5	G5			✓
OBBA (2005)	BB (2009)	Black-capped Chickadee	<i>Poecile atricapillus</i>				S5	G5		4	
OBBA (2005)	BB (2009)	Red-breasted Nuthatch	<i>Sitta canadensis</i>				S5	G5		3	✓
OBBA (2005)	BB (2009)	White-breasted Nuthatch	<i>Sitta carolinensis</i>				S5	G5		3	
OBBA (2005)		Brown Creeper	<i>Certhia americana</i>				S5B	G5		2	✓
OBBA (2005)	BB (2009)	House Wren	<i>Troglodytes aedon</i>				S5B	G5		4	
OBBA (2005)		Winter Wren	<i>Troglodytes troglodytes</i>				S5B	G5		3	✓
OBBA (2005)		Sedge Wren	<i>Cistothorus platensis</i>	NAR	NAR		S4B	G5		2	✓
OBBA (2005)		Marsh Wren	<i>Cistothorus palustris</i>				S4B	G5		2	✓
OBBA (2005)	I (2012)	Golden-crowned Kinglet	<i>Regulus satrapa</i>				S5B	G5		2	✓
	I (2012)	Ruby-crowned Kinglet	<i>Regulus calendula</i>				S4B	G5		2	✓
OBBA (2005)		Eastern Bluebird	<i>Sialia sialis</i>	NAR	NAR		S5B	G5		3	
OBBA (2005)		Veery	<i>Catharus fuscescens</i>				S4B	G5		3	✓
OBBA (2005)		Wood Thrush	<i>Hylocichla mustelina</i>	SC	THR		S4B	G5	✓	1	
OBBA (2005)	BB (2009)	American Robin	<i>Turdus migratorius</i>				S5B	G5		4	
OBBA (2005)	BB (2009)	Gray Catbird	<i>Dumetella carolinensis</i>				S4B	G5		3	
OBBA (2005)		Brown Thrasher	<i>Toxostoma rufum</i>				S4B	G5	✓	2	✓
OBBA (2005)	BB (2009)	Cedar Waxwing	<i>Bombycilla cedrorum</i>				S5B	G5		3	
OBBA (2005)	BB (2009)	European Starling	<i>Sturnus vulgaris</i>				SNA	G5		5	
	BB (2009)	Blue-headed Vireo	<i>Vireo solitarius</i>				S5B	G5		2	✓
OBBA (2005)	BB (2009)	Warbling Vireo	<i>Vireo gilvus</i>				S5B	G5		4	
OBBA (2005)	BB (2009)	Red-eyed Vireo	<i>Vireo olivaceus</i>				S5B	G5		4	
OBBA (2005)	BB (2009)	Nashville Warbler	<i>Vermivora fuficapilla</i>				S5B	G5		2	
OBBA (2005)	BB (2009)	Yellow Warbler	<i>Dendroica petechia</i>				S5B	G5		4	
OBBA (2005)		Chestnut-sided Warbler	<i>Dendroica pensylvanica</i>				S5B	G5		2	
OBBA (2005)		Magnolia Warbler	<i>Dendroica magnolia</i>				S5B	G5		2	✓
OBBA (2005)	BB (2009)	Yellow-rumped Warbler	<i>Dendroica coronata</i>				S5B	G5		2	
OBBA (2005)	BB (2009)	Black-throated Green Warbler	<i>Dendroica virens</i>				S5B	G5		2	✓
OBBA (2005)	BB (2009)	Pine Warbler	<i>Dendroica pinus</i>				S5B	G5		3	✓
OBBA (2005)		Black-and-white Warbler	<i>Mniotilta varia</i>				S5B	G5		3	✓
OBBA (2005)	BB (2009)	American Redstart	<i>Setophaga ruticilla</i>				S5B	G5		3	✓
OBBA (2005)		Ovenbird	<i>Seiurus aurocapilla</i>				S4B	G5		3	✓
OBBA (2005)	BB (2009)	Northern Waterthrush	<i>Seiurus noveboracensis</i>				S5B	G5		3	
OBBA (2005)		Mourning Warbler	<i>Opipornis philadelphia</i>				S4B	G5		3	
OBBA (2005)	BB (2009)	Common Yellowthroat	<i>Geothlypis trichas</i>				S5B	G5		4	
OBBA (2005)		Canada Warbler	<i>Wilsonia canadensis</i>	SC	THR	THR	S4B	G5		1	✓
OBBA (2005)		Scarlet Tanager	<i>Piranga olivacea</i>				S4B	G5		3	✓
OBBA (2005)	BB (2009)	Northern Cardinal	<i>Cardinalis cardinalis</i>				S5	G5		4	✓
OBBA (2005)	BB (2009)	Rose-breasted Grosbeak	<i>Pheucticus ludovicianus</i>				S4B	G5	✓	3	✓
OBBA (2005)	BB (2009)	Indigo Bunting	<i>Passerina cyanea</i>				S4B	G5			
OBBA (2005)		Eastern Towhee	<i>Pipilo erythrophthalmus</i>				S4B	G5	✓	3	✓
OBBA (2005)	BB (2009)	Chipping Sparrow	<i>Spizella passerina</i>				S5B	G5		4	

APPENDIX 20. BACKGROUND WILDLIFE LIST

Hillsburgh Dam Environmental Assessment, Natural Heritage - Existing Conditions

SOURCE	CVC (most recent observation indicated)	COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-Rank	G-Rank	PIF (BCR 13)	CVC Tier (2010)	Wellington County (2008)
OBBA (2005)	BB (2009)	Field Sparrow	<i>Spizella pusilla</i>				S4B	G5	<		<
OBBA (2005)		Vesper Sparrow	<i>Poocetes gramineus</i>				S4B	G5	✓	2	✓
OBBA (2005)		Savannah Sparrow	<i>Passerculus sandwichensis</i>				S4B	G5	✓	4	✓
OBBA (2005)		Grasshopper Sparrow	<i>Ammodramus savannarum</i>		SC		S4B	G5TU	✓	1	✓
OBBA (2005)	BB (2009)	Song Sparrow	<i>Melospiza melodia</i>				S5B	G5		4	
OBBA (2005)	BB (2009)	Swamp Sparrow	<i>Melospiza georgiana</i>				S5B	G5		4	
OBBA (2005)	BB (2009)	White-throated Sparrow	<i>Zonotrichia albicollis</i>				S5B	G5		3	
	SWH (2011)	Dark-eyed Junco	<i>Junco hyemalis</i>				S5B	G5		2	✓
OBBA (2005)		Bobolink	<i>Dolichonyx oryzivorus</i>	THR	THR		S4B	G5	✓	1	✓
OBBA (2005)	BB (2009)	Red-winged Blackbird	<i>Agelaius phoeniceus</i>				S4	G5		4	
OBBA (2005)		Eastern Meadowlark	<i>Sturnella magna</i>		THR		S4B	G5	✓	1	✓
OBBA (2005)	BB (2009)	Common Grackle	<i>Quiscalus quiscula</i>				S5B	G5		4	
OBBA (2005)	BB (2009)	Brown-headed Cowbird	<i>Molothrus ater</i>				S4B	G5		4	
OBBA (2005)	BB (2009)	Baltimore Oriole	<i>Icterus galbula</i>				S4B	G5	✓	3	✓
OBBA (2005)	BB (2009)	Purple Finch	<i>Carpodacus purpureus</i>				S4B	G5		2	
OBBA (2005)	BB (2009)	House Finch	<i>Carpodacus mexicanus</i>				SNA	G5		5	
		Pine Siskin	<i>Carduelis pinus</i>				S4B	G5		2	
OBBA (2005)	BB (2009)	American Goldfinch	<i>Carduelis tristis</i>				S5B	G5		4	
OBBA (2005)	BB (2009)	House Sparrow	<i>Passer domesticus</i>				SNA	G5		5	
		MAMMALS									
OMA (1994)		Virginia Opossum	<i>Didelphis virginiana</i>				S4	G5		4	
OMA (1994)		Star-nosed Mole	<i>Condylura cristata</i>				S5	G5		2	
OMA (1994)		Little Brown Myotis	<i>Myotis lucifugus</i>	END	END	END	S4	G3G4		1	
OMA (1994)		Big Brown Bat	<i>Eptesicus fuscus</i>				S5	G5		3	
OMA (1994)		Hoary Bat	<i>Lasiurus cinereus</i>				S4	G4		3	
OMA (1994)		Eastern Cottontail	<i>Sylvilagus floridanus</i>				S5	G5		4	
OMA (1994)		Snowshoe Hare	<i>Lepus americanus</i>				S5	G5		2	
OMA (1994)		European Hare	<i>Lepus europaeus</i>				SNA	G5		5	
OMA (1994)	I (2009)	Eastern Chipmunk	<i>Tamias striatus</i>				S5	G5		3	
OMA (1994)		Woodchuck	<i>Marmota monax</i>				S5	G5		3	
OMA (1994)		Eastern Gray Squirrel	<i>Sciurus carolinensis</i>				S5	G5		4	
OMA (1994)	I (2002)	Red Squirrel	<i>Tamiasciurus hudsonicus</i>				S5	G5		3	
OMA (1994)	I (2014)	Beaver	<i>Castor canadensis</i>				S5	G5		3	
OMA (1994)		Meadow Vole	<i>Microtus pennsylvanicus</i>				S5	G5		3	
OMA (1994)		Muskrat	<i>Ondatra zibethicus</i>				S5	G5		3	
OMA (1994)		Norway Rat	<i>Rattus norvegicus</i>				SNA	G5		5	
OMA (1994)	I (2003)	Porcupine	<i>Erethizon dorsatum</i>				S5	G5		2	
OMA (1994)		Coyote	<i>Canis latrans</i>				S5	G5		3	
OMA (1994)		Red Fox	<i>Vulpes vulpes</i>				S5	G5		3	
OMA (1994)		Northern Raccoon	<i>Procyon lotor</i>				S5	G5		4	
OMA (1994)		Ermine	<i>Mustela erminea</i>				S5	G5		2	
OMA (1994)		Long-tailed Weasel	<i>Mustela frenata</i>				S4	G5		3	✓
OMA (1994)	I (2012)	American Mink	<i>Mustela vison</i>				S4	G5		2	
OMA (1994)	I (2009)	Striped Skunk	<i>Mephitis mephitis</i>				S5	G5		4	

SOURCE	CVC (most recent observation indicated)	COMMON NAME	SCIENTIFIC NAME	SARO	COSEWIC	SARA	S-Rank	G-Rank	PIF (BCR 13)	CVC Tier (2010)	Wellington County (2008)
OMA (1994)	I (2008)	White-tailed Deer	<i>Odocoileus virginianus</i>				S5	G5		3	

Legend:

SARO: Species at Risk Ontario
 COSEWIC: Committee on the Status of Endangered wildlife in Canada
 SARA: Species at Risk Act
 ESA: Endangered Species Act
 END: Endangered
 THR: Threatened
 SC: special Concern
 NAR: Not At Risk
 NL: Not listed
 DD: Data Deficient
 I: Incidental Observation
 BB: Breeding Bird Survey Observation-Breeding Evidence noted
 NB: Breeding Bird Survey Observation-No Breeding evidence noted
 SWH: Significant Wildlife Habitat survey
 FM: Fall migration Survey
 SM: Spring Migration Survey

Source codes

OBAB: Ontario butterfly Atlas Online
 ORAA: Ontario Reptile and Amphibian Atlas
 OMA: Ontario Mammal Atlas
 OBBA: Ontario Breeding Bird Atlas
 CVC: Credit Valley Conservation Data, provided 2014

CVC Tiers:

- 1 - Species of Conservation Concern
- 2 - Species of Interest
- 3 - Species of Urban Interest
- 4 - Secure Species
- 5 - Non-native & Non-native Hybrid Species

Global Rank:

- G1: Extremely rare globally
- G1G2: Extremely rare to very rare globally
- G2: Very rare globally
- G2G3: Very rare to uncommon globally
- G3: Rare to uncommon globally
- G3G4: Rare to common globally
- G4: Common globally
- G4G5: Common to very common globally
- G5: Very common globally; demonstrably secure
- T: rank applies to a subspecies or variety

Provincial Rank:

- S1: Critically Imperiled—Critically imperiled in the province
- S2: Imperiled—Imperiled in the province, very few populations
- S3: Vulnerable—Vulnerable in the province, relatively few populations
- S4: Apparently Secure—Uncommon but not rare
- S5: Secure—Common, widespread, and abundant in the province
- SX: Presumed extirpated
- SH: Possibly Extirpated (Historical)
- SNR: Unranked
- SU: Unrankable—Currently unrankable due to lack of information
- SNA: Not applicable—A conservation status rank is not applicable
- S#S#: Range Rank— indicates range of uncertainty about the status
- S#B- Breeding status rank
- S#N- Non Breeding status rank
- ?: Indicates uncertainty in the assigned rank

Wellington County:

- ✓ Significant Species

PIF:

- ✓ Priority Species



CVC	MNRF	COMMON NAME	SCIENTIFIC NAME	COSARO	COSEWIC	SARA	S-Rank	G-Rank	CVC (2010)
✓		Slimy Sculpin	<i>Cottus cognatus</i>				S5	G5	2
✓		Banded Killifish	<i>Fundulus diaphanus</i>				S5	G5	2
✓	✓	Brook Trout	<i>Salvelinus fontinalis</i>				S5	G5	2
✓		Golden Shiner	<i>Notemigonus crysoleucas</i>				S5	G5	3
✓		Central Mudminnow	<i>Umbra limi</i>				S5	G5	3
✓	✓	Rock Bass	<i>Ambloplites rupestris</i>				S5	G5	4
✓	✓	White Sucker	<i>Catostomus commersonii</i>				S5	G5	4
✓		Brook Stickleback	<i>Culaea inconstans</i>				S5	G5	4
✓	✓	Pumpkinseed	<i>Lepomis gibbosus</i>				S5	G5	4
✓	✓	Common Shiner	<i>Luxilus cornutus</i>				S5	G5	4
✓	✓	Largemouth Bass	<i>Micropterus salmoides</i>				S5	G5	4
✓	✓	Bluntnose Minnow	<i>Pimephales notatus</i>				S5	G5	4
✓	✓	Eastern Blacknose Dace	<i>Rhinichthys atratulus</i>				S5	G5	4
✓		Creek Chub	<i>Semotilus atromaculatus</i>				S5	G5	4
✓	✓	Round Goby	<i>Neogobius melanostomus</i>				SNA	G5	5
✓	✓	Brown Trout	<i>Salmo trutta</i>				SNA	G5	5

Data Source:

CVC: Data collected within the study area between 1954 and 2013; method of collection unknown.

MNRF: Data collected within the study area between 2013 and 2014; Data collected through electrofishing, drift nets, minnow traps and incidental observations.

Legend:

COSARO: Committee on Species at Risk Ontario

COSEWIC: Committee on the Status of Endangered Wildlife in Canada

SARA: Species at Risk Act

ESA: Endangered Species Act

END: Endangered

THR: Threatened

SC: special Concern

NAR: Not At Risk

NL: Not listed

DD: Data Deficient

CVC Tiers:

1 - Species of Conservation Concern

2 - Species of Interest

3 - Species of Urban Interest

4 - Secure Species

5 - Non-native & Non-native Hybrid Species

G-Rank:

G1: Extremely rare globally

G1G2: Extremely rare to very rare globally

G2: Very rare globally

G2G3: Very rare to uncommon globally

G3: Rare to uncommon globally

G3G4: Rare to common globally

G4: Common globally

G4G5: Common to very common globally

G5: Very common globally; demonstrably secure

T: rank applies to a subspecies or variety

S-Rank:

S1: Critically Imperiled—Critically imperiled in the province

S2: Imperiled—Imperiled in the province, very few populations

S3: Vulnerable—Vulnerable in the province, few populations

S4: Apparently Secure—Uncommon but not rare

S5: Secure—Common, widespread, and abundant in the province

SX: Presumed extirpated

SH: Possibly Extirpated (Historical)

SNR: Unranked

SU: Unrankable—Currently unrankable due to lack of information

SNA: Not applicable—conservation status rank is not applicable

S#S#: Range Rank—range of uncertainty about the status

S#B- Breeding status rank

S#N- Non Breeding status rank

?: Indicates uncertainty in the assigned rank

Appendix 21
Curriculum Vitae

DRAFT

Aboud & Associate Inc. - Project staff.

Steven Aboud: Principal. Senior Ecologist

Ryan Hamelin: Terrestrial and Wetland Ecologist

Cheryl-Anne Ross: Wildlife Ecologist

Matt Iles: Biologist (No longer with Aboud & Associates Inc.)

DRAFT



Steven About

B.Sc. (Botany)

Principal . Senior Ecologist . ISA Certified Arborist

BIO

Steven has thirty five years of public and private sector experience in the disciplines of arboriculture and ecology. His considerable experience includes testifying as an expert witness before the Ontario Municipal Board, expert testimony on legal matters related to trees, urban forestry policy development and assessment of natural heritage features across southern Ontario. Steven is the author of several publications and documents related to woodland restoration, schoolyard naturalization and the status of rare tree species. He continues to lead a team of skilled and creative individuals developing practical and cost-effective solutions to urban forestry, ecology and landscape design issues using natural systems models.

EDUCATION

1978
Bachelor of Science (Botany)
University of Guelph

PROFESSIONAL AFFILIATIONS

International Society of Arboriculture
Society for Ecological Restoration
Tallgrass Ontario
Ontario Urban Forest Council

PROFESSIONAL CERTIFICATIONS

ISA Certified Arborist ON-03234
International Society of Arboriculture
Butternut Health Assessor No. 497
Ontario Ministry of Natural Resources
ISA Tree Risk Assessment Qualified
International Society of Arboriculture
Ontario Wetland Evaluator (OWES)
Ontario Ministry of Natural Resources

PROFESSIONAL EXPERIENCE

1991-1996
University of Guelph - The Arboretum
Coordinator of Interpretive Programs
1991-1996
The Seed to Seed (Guelph) . *Proprietor*
1978-1991
University of Guelph - The Arboretum
*Manager of Tree and Shrub
Collections and Plant Data*

RELEVANT PROJECT EXPERIENCE

URBAN FORESTRY

- Heritage (Bronte) White Oak Monitoring (Oakville)
- Downtown Brampton Street Tree Inventory (Brampton)
- Milton Urban Area Public Lands Tree Inventory (Milton)
- Whitby Tree Inventory Asset Management Project (Whitby)
- Allan Gardens Revitalization (Toronto)
- Graham Arboretum Renewal Master Plan (London)
- Aurthur Street Sanitary EA (Guelph)
- Gordon/Woolwich Streets (Guelph)
- Watson Parkway (Guelph)
- James Mountain Road (Hamilton)
- Red Hill Valley Parkway (Hamilton)
- Glen Abby Golf Club (Oakville)
- Lambton Golf Club (Toronto)
- Wrigley Canada Headquarters (Toronto)
- Canadian National Institute for the Blind Headquarters (Toronto)
- Parc Downsview Park (Toronto)
- Sanofi-Pasteur Pharmaceuticals Connaught Campus (Toronto)

EXPERT & WRITTEN TESTIMONY AND PEER REVIEW

- Residential Development OMB Hearings (Hamilton . Pickering . Toronto)
- Woodland Policy OMB (London)
- Hurontario Street Expropriation (Caledon)
- Tree Failure Assessment/Testimony (London . Stoney Creek)
- Street Tree Planting Deficiencies Review (Guelph)
- Township of Centre Wellington Tree Policy (Centre Wellington)
- Natural Environment Level 1 and 2 Report Reviews (Centre Wellington)
- Environmental Impact Study Report Reviews (Centre Wellington)



PROFESSIONAL DEVELOPMENT

- 2012
IML Resistograph Technical Workshop
Town of Oakville
- 2012
Tree Planting BMP's and Changes to
Development Manual Workshop
City of Kitchener
- 2012
Ottawa Heritage Tree Workshop
Ontario Urban Forest Council
- 2011
Erosion and Sediment Control Workshop
Toronto and Region Conservation Authority
- 2011
Soils and Urban Trees Conference
Toronto Botanical Gardens
- 2007
Structural Soil and Care, Selection and
Management of Urban Street Trees
Cornell University & City of Ithaca
- 2004
Statistics Intergrating Estimation
Method of Tree Risk Assessment
Arbormaster Training Canada

TEACHING

- 1999
Certified Arborist Program . Instructor
International Society of Arboriculture
- 1997
Resource Management Field Camp
Sessional Lecturer . *University of Guelph*
- 1997
Ecosystem Restoration Post Graduate
Program . Guest Lecturer
Niagara College
- 1995-1996
A Life Zone Approach to Schoolyard
Naturalization Series Workshops
University of Guelph
- 1978-1996
Natural Interpretation Workshops
University of Guelph

COMMUNITY SERVICE

- 1990-1991
Trees For Guelph

ECOLOGICAL RESTORATION

- Creditview Crossing Community Woodlot Management Plans (Brampton)
- Jefferson Forest Community Edge Management Plans (Richmond Hill)
- Wrigley Canada Ravine Stewardship Plan (Toronto)
- Private Residential Stewardship Plans (Toronto)
- The Rosewood Condominium Stewardship Plan (Toronto)
- Environment Canada - Downsview Campus Naturalization Master Plan (Toronto)
- Health Canada - Campus Naturalization Master Plan (Toronto)
- Canada Centre for Inland Waters - Campus Naturalization Master Plan (Burlington)
- Wellington Terrace Wetland Enhancement (Elora)
- A.M. Cunningham Public School Naturalization (Hamilton)
- Victory Public School Naturalization (Milton)
- Thornhill Woods Community Woodlot Edge Management Plans (Vaughan)
- Thornhill Ravines Community Valley Features Edge Management Plan (Vaughan)
- Upper Thornhill Estates Community Natural Systems Edge Management Plans (Vaughan)
- Vellore Village Woodlot Edge Management Plans (Vaughan)

ENVIRONMENTAL STUDIES

- Jefferson Forest Community EIS (Richmond Hill)
- Community Planning Area Subwatershed Study (Course Wellington)
- Wetland Boundary Delineation (Erin)
- Sunset Hills Estates EIS (Woolwich)
- Valley Road Estates EIS (Guelph)
- Bird Landing EIS (Guelph)
- River Systems Assessment Study (Guelph)
- Sawmill Valley EIS (Mississauga)
- Blue Heron Ridge EIS (Cambridge)
- Mount Pleasant GO Station EA (Brampton)
- ASECO Intergrated Systems EIS (Oakville)
- Functional Servicing Development Area Study (King)
- King City East Buffer Strategy (King)
- MTRCA Woody Plant Selection Guidelines (Greater Toronto Bio-Region)
- West Humber River Naturalization Plan (Toronto)
- Highway 407 ETR (Toronto)
- E.T. Seton Park Naturalization Plan (Toronto)
- Brampton Sports Parks (Brampton)
- Brampton Vegetation Assessment (Brampton)

PUBLICATIONS

- Aboud, S. W. and H. Kock. 1994 (1996 Rev. ed.) A life zone approach to school yard naturalization: the Carolinian life zone. University of Guelph. Guelph, Ontario. 86 pp.
- Waldron, G.E., S. W. Aboud, J. D. Ambrose and G.A. Meyers. 1987. Shumard Oak (*Quercus shumardii*) in Canada. *Can. Field Naturalist* 101: 532-538.
- Ambrose, J.D. and S. W. Aboud. 1985. Status report on *Castanea dentata*. COSEWIC, Ottawa.
- Ambrose, J.D. and S. W. Aboud. 1983. Status report on *Magnolia acuminata*. COSEWIC, Ottawa.
- Ambrose, J.D. and S. W. Aboud. 1982a. Status report on *Fraxinus quadrangulata* COSEWIC, Ottawa
- Ambrose, J.D. and S. W. Aboud. 1982b. Status report on *Ptelea trifoliata*. COSEWIC, Ottawa.



Ryan Hamelin

M.Sc., B.Sc.(Env)

Terrestrial and Wetland Ecologist

BIO

Ryan is an experienced ecologist with a diverse background in the public and private sectors. He has a proven track record of liaising with government agencies, engineers, landowners, and contractors to manage and complete multifaceted, complex projects. Ryan's strong project management and field ecology skills are successfully applied to wetland restoration, wetland evaluations, vegetation surveys, habitat surveys and water quality monitoring projects across southern Ontario.

EDUCATION

- 2012
M.Sc. Integrative Biology
University of Guelph
- 2008
B.Sc. Environmental Science
University of Guelph

PROFESSIONAL EXPERIENCE

- 2014-present
Aboud & Associates
Terrestrial and Wetland Ecologist
- 2013-2014
Ontario Ministry of Natural Resources
Coastal Wetland Biologist
- 2013
Toronto & Region Conservation Authority
Environmental Field Labourer
- 2006-2010
Severn Sound Environmental Association
Environmental Technician

SELECTED PROJECT EXPERIENCE

ECOLOGICAL RESTORATION

- **Rondeau Wetland Restoration and Watershed Buffer Program (Chatham-Kent)**
*with Ontario Ministry of Natural Resources
- **Carolinian Forest Tree Planting Project (Chatham-Kent)**
*with Ontario Ministry of Natural Resources

ENVIRONMENTAL STUDIES

- **Rondeau Vegetation Monitoring Surveys (Chatham-Kent)**
*with Ontario Ministry of Natural Resources
- **Near Shore Fish Habitat Evaluation (Simcoe County)** *with Severn Sound Environmental Association
- **Loon Surveys (Killarney)** *with Ontario Ministry of Natural Resources
- **Fish Habitat/Population Electrofishing surveys (Toronto Harbour)**
*with Toronto and Region Conservation Authority
- **Benthic Biomonitoring Surveys (Simcoe County)**
*with Severn Sound Environmental Association
- **Ladysmith Wetland Evaluation (Lambton County)**
*with Ontario Ministry of Natural Resources

WATER QUALITY MONITORING

- **Talbot Tract Water Quality Monitoring (Chatham-Kent)** *with Ontario Ministry of Natural Resources
- **Environ Property Water Quality Monitoring (St. Williams)** *with Ontario Ministry of Natural Resources
- **Provincial Water Quality Monitoring Network (Simcoe County)**
*with Severn Sound Environmental Association
- **Provincial Ground Water Monitoring Network (Simcoe County)**
*with Severn Sound Environmental Association



PROFESSIONAL CERTIFICATIONS

2013
Wetland Evaluation Training Course
Ontario Ministry of Natural Resources

2013
Ecological Land Classification Training
Course (ELC)
Ontario Ministry of Natural Resources

PROFESSIONAL DEVELOPMENT

- 2014 - Great Lakes Wetland Day
- 2013 - Wetland Graminoid Identification Workshop (Royal Botanical Gardens)
- 2013 - Ontario Fish Identification Workshop (Royal Ontario Museum)
- 2013 - Introduction to the Canadian Environmental Assessment Act (Canadian Environmental Assessment Agency)
- 2012 - Water Management & Wetland Restoration Training Course (OMNR)

DRAFT





Cheryl-Anne Ross

Fish and Wildlife Technologist Dip., B.Sc.
Wildlife Ecologist

EDUCATION

- 2007
B.Sc. Natural Resource Management
University of Northern British Columbia
- 2004
Fish and Wildlife Technologist Diploma
Sir Sanford Fleming College

PROFESSIONAL EXPERIENCE

- 2011-2014
Stantec
Terrestrial Ecologist
- 2008-2010
Natural Resources Solutions Inc.
Terrestrial and Wetland Ecologist
- 2006
Earl Rowe Provincial Park
Natural Heritage Educator
- 2004
Ministry of Natural Resources
Field Ecologist
- 2003
Wye Marsh Wildlife Centre
Outdoor Education/Naturalist

PROFESSIONAL CERTIFICATIONS

- Ecological Land Classification (ELC)
Ontario Ministry of Natural Resources
- ISA Certified Arborist
International Society of Arboriculture



ABOUD & ASSOCIATES INC.
Consulting Arborists • Ecologists • Landscape Designers

BIO

Cheryl-Anne Ross has a comprehensive understanding of the natural environment and the tools that are used to evaluate it. Cheryl-Anne completed her Undergraduate degree in NREM-Wildlife and Fisheries Biology at the University of Northern British Columbia, has a Technical Diploma in Fish and Wildlife Technology from Sir Sanford Fleming College and a decade of experience in the public and private sectors. In addition to working as a Natural Heritage Educator at provincial parks, her professional experience includes working on residential and industrial development and renewable energy projects throughout Ontario. Cheryl-Anne conducts a broad range of natural heritage inventory and assessments including botanical inventories, ELC, species at risk reports, environmental assessments, environmental impact statements, and monitoring for wildlife (avian, herptiles and mammals) and wildlife habitat.

SELECTED PROJECT EXPERIENCE

OIL AND GAS DEVELOPMENT

- TransCanada Energy East Pipeline (Ontario) *with Stantec
- Payne-Sarnia Pipeline (Sarnia) *with Stantec
- Natural Gas Development Project (Burlington/Oakville) *with Stantec
- Brantford Kirkwall Pipeline (Kirkwall) *with Stantec
- NOVA Genesis Pipeline Extension (Corunna) *with Stantec

RENEWABLE ENERGY

- Amherst Island Wind Farm (Amherst Island) *with Stantec
- Cedar Point Wind Farm (Forest) *with Stantec
- Bow Lake Wind Farm (Montreal River Harbour) *with Stantec
- Niagara Region Wind Centre (Niagara Region) *with Stantec

INDUSTRIAL DEVELOPMENT

- Dundalk Industrial Park Municipal Class Environmental Assessment (Southgate)
- Erin Pit Extension (Orangeville) *with Stantec
- Industrial Development Project (Milton) *with Stantec
- NOVA 2020 Plant Expansion (Corunna) *with Stantec

RESIDENTIAL DEVELOPMENT

- Westside, Vista Hills, Clair Creek Meadows Developments (Waterloo) *with Stantec

MUNICIPAL INFRASTRUCTURE

- Hillsburgh Dam Municipal Class Environmental Assessment (Erin)

COMMUNITY SERVICE

2009

Ontario Streams
Spawning Survey

2009

Ontario Public Interest Research Group
Tree Planter

2004

Florida Panther National Wildlife Refuge
Invasive Plant Species Control

ENVIRONMENTAL IMPACT STUDIES . WILDLIFE STUDIES

- *Sunset Hills Estates Wildlife Studies (Maryhill)*
- *Martin Street School Environmental Impact Study (Milton)*
- *Wellington Street Improvements Active Bird Nest Inventory (Guelph)*
- *Listowel Environmental Impact Study (North Perth)*
- *Private Property Severance, Environmental Impact Study (Puslinch)*
- *Block 18 Woodlot Active Bird Nest Inventory (Vaughan)*
- *Elora Development Active Bird Nest Inventory (Centre Wellington)*
- *Rockwood Commercial Development Amphibian Habitat Assessment (Guelph/Eramosa)*

DRAFT





Matthew Iles

M.Sc., B.Sc.
Biologist

BIO

Matthew is a well-rounded biologist with diverse experiences studying aquatic and terrestrial ecosystems in Europe, South America and Ontario. He has demonstrated an ability to design and implement research projects, in addition to working closely with indigenous communities, volunteers, landowners and other vested stakeholders, on conservation and ecology projects. An avid field ornithologist, Matthew has held positions and volunteered with the Canadian Wildlife Service, Bird Studies Canada and Long Point Bird Observatory.

EDUCATION

- 2007
M.Sc. of Restoration Ecology of
Terrestrial and Aquatic Environment
University of Liverpool
- 2006
B.Sc. Zoology (with honours)
University of Liverpool

PROFESSIONAL EXPERIENCE

- 2012
Long Point Waterfowl
Research Technician
- 2012
Bird Studies Canada
Research Technician
- 2012
Innis Point Bird Observatory
Band-in-charge
- 2008-2009
Global Vision International, Rainforest
Conservation and Community
Development Expedition
Field Staff and Biologist
- 2007-2008
The Environment Partnership
Ecologist

SELECTED PROJECT EXPERIENCE

ORNITHOLOGY

- Nanaksar Gurdwara Gursikh Temple Breeding Bird Survey (Brampton)
- Caledon Structure (Bridge/Culvert) Program Nest Searches (Caledon)
- Bird Landing, breeding bird monitoring and protection measures (Guelph)
- Caledon East Reservoir Nest Searches (Caledon)
- Courtice Road Breeding Bird Survey (Clarington)
- Stone Road Reconstruction Nest Search (Guelph)
- Britannia Rd. Woodlot, forest bird breeding survey (Halton Region)
- French's Bridge Rehabilitation Nest Search (Puslinch)
- Perth County Rd. 313 Bridge, nesting swallow inventory (Perth County)
- Private Property Severance, 87 acre breeding bird survey (Bowmanville)
- Waterfowl capture, observation and study (Prince Edward County, Toronto, Hamilton, Wolfe Island, Lake St. Clair) *with Long Point Waterfowl and Canadian Wildlife Service
- Population counts and behavioral observations (Manitoulin Island) *with Long Point Waterfowl
- Golden-winged Warbler (SAR - Thr.) capture, blood sampling, territory assessments (Elgin County) *with Bird Studies Canada
- Eastern Whip-poor-will (SAR - Thr.) point count surveys (Elgin County) *with Bird Studies Canada
- Migration monitoring, including capture, banding, daily census (Kanata, Long Point) *with Innis Point Bird Observatory and Long Point Bird Observatory
- Grey Jay nest searching (Algonquin Provincial Park) *with University of Guelph
- Breeding ecology of Tree Swallows (Long Point) *with Long Point Bird Observatory/University of Guelph
- Inventorying rainforest birds (Ecuador) *with Global Vision International
- Hen Harrier nest searching (Isle of Man, UK) *with The Environmental Partnership



COMMUNITY SERVICE

2011-2012
Canadian Wildlife Service
Waterfowl Research Technician

2012
Norris Lab University of Guelph
Avian Technician

2012
Carleton University
Entomology Lab Technician

2011-2012
Long Point Bird Observatory
Avian Biologist

2010
Halton Region Conservation Authority
Casual Volunteer

2009
Nazca Institute for Marine Research
Marine Biologist

ENVIRONMENTAL IMPACT STUDIES (EIS)

- Sunset Hills Estates, Anuran call survey and Breeding bird survey, GIS, report (Maryhill)
- Lakeshore Boulevard West Development Natural Heritage Impact Study (Toronto)
- Hill Street Bridge Environmental Impact Study (Woolwich)
- Wellington Terrace Service Road Environmental Impact Statement (Centre Wellington)
- Private Property Severance, Anuran Call Survey, breeding bird survey, GIS, report (Puslinch)
- Woodlawn Road Development Environmental Impact Study (Guelph)
- Brock Road Development Environmental Impact Study (Puslinch)
- Longyards Community Trail Impact Study (Vaughan)
- Gordon Street Development Natural Heritage Peer Review (Guelph)

BOTANY, STEWARDSHIP, HERPETOLOGY, MAMMALS

- Eden Park Butternut Restoration Monitoring Program (Hamilton)
- Woodlawn Road Development Environmental Impact Study (Guelph)
- Elmira Road Industrial Expansion Vegetation Inventory (Guelph)
- 11th Concession Road Development Vegetation Inventory (Hamilton)
- Habitat and vegetation assessments for Golden-winged Warbler and Blue-winged Warbler (Elgin County, Norfolk County) *with Bird Studies Canada
- Habitat stewardship and restoration, including tree sapling planting and invasive species control *with Long Point Bird Observatory
- Reptile and Amphibian Inventory (Ecuador) *with Global Vision International
- Surveying and stewardship for protected amphibian species, including great-crested newt (UK) *with The Environmental Partnership
- Surveying protected mammal species, including bats and water vole (UK) *with The Environmental Partnership

ENTOMOLOGY

- Dung beetle and benthic invertebrate communities in fragmented rainforest habitats (Ecuador) *with Global Vision International
- Butterfly Inventory (Ecuador) *with Global Vision International
- Effect of species-rich grassland translocation on invertebrate communities, with particular reference to Carabid beetles (Manchester, UK) *M.Sc. Research, University of Liverpool

MARINE BIOLOGY

- Willoughby Road Bridge Fish Rescue (Caledon)
- Arkell Dam Fish Rescue (Guelph)
- Artisanal lobster fisheries and by-catch (Ecuador) *with Nazca Institute for Marine Research
- Colonization of man-made substrates by an invasive barnacle species (Liverpool, UK) *with University of Liverpool
- Transition rates in sessile inter-tidal organisms (Liverpool, UK) *with University of Liverpool

PUBLICATIONS

Bracewell SA, Spencer M, Marrs RH, Iles M, Robinson LA (2012). Cleft, Crevice, or the Inner Thigh: 'Another Place' for the Establishment of the Invasive Barnacle *Austrominius modestus* (Darwin, 1854). PLoS ONE 7(11)



DRAFT

- Urban Forestry
- Ecological Restoration
- Landscape Architecture
- Environmental Studies
- Expert Opinion

ABOUD & ASSOCIATES INC.
Consulting Arborists • Ecologists • Landscape Designers



591 Woolwich Street . Guelph . Ontario . N1H 3Y5 . T:519.822.6839 . F:519.822.4052 . info@aboudtng.com . www.aboudtng.com